

BRVKENTHAL. ACTA MVSEI

VII. 3

MINISTERUL CULTURII ȘI PATRIMONIULUI NAȚIONAL

MUZEUL NAȚIONAL BRUKENTHAL

BRVKENTHAL

ACTA MVSEI

VII. 3

EDITOR IN CHIEF: Prof. Univ. Dr. Sabin Adrian LUCA

SECRETARIAL REDACTION: Dr. Anca NIȚOI
Dr. Iulia MESEA
Ioan TĂUȘAN
Iulia-Maria PASCU

EDITORIAL BOARD: Dr. Raluca-Maria TEODORESCU

Alexandru SONOC
Dr. Constantin ITTU
Dr. Rodica CIOBANU
Ana Maria MESAROȘ
Cecilia HĂRĂSTAȘAN
Dr. Dorin BARBU
Dr. Dana HRIB

ASSOCIATED MEMBERS TO THE EDITORIAL BOARD:

Prof. Dr. Docent Theodor Anton NEAGU (Member of the Romanian Academy)
Prof. Univ. Dr. Ioan-Aurel POP (Member of the Romanian Academy)
Prof. Dr. Paul NIEDERMAIER (Correspondent Member of the Romanian Academy)
Prof. Dr. Conrad GÜNDISCH (Universität Oldenburg – Germany).
Prof. Dr. Erika SCHNEIDER – BINDER (Universität Karlsruhe, Bereich
WWF Auen Institut – Germany)
Prof. Dr. Zeno-Karl PINTER (“Lucian Blaga” University, Sibiu)
Prof. Dr. Rudolf GRÄF (“Babeș-Bolyai” University, Cluj Napoca)
Prof. Dr. Nicolae SABĂU (“Babeș-Bolyai” University, Cluj Napoca)
Prof. Dr. Alexandru AVRAM (“Lucian Blaga” University, Sibiu)

ISSN 2285 - 9470 ISSN-L 1842 - 2691



Editura Muzeului Național Brukenthal

Orice corespondență referitoare la această publicație rugăm a se adresa la:

Muzeul Național Brukenthal - Piața Mare 4-5, 550163, Sibiu. **Tel:** 004/369/101782; **Fax:** 004/ 269 /211545.

E-mail: ioan.tausan@brukenthalmuseum.ro; Website: www.brukenthalmuseum.ro

Autorii își vor asuma întreaga responsabilitate pentru informația de specialitate din materialele trimise, care vor fi supuse procesului de peer review, ale cărui detalii pot fi consultate la <http://www.brukenthalmuseum.ro/publicatii/01.htm>

Ghidul pentru autori se regăsește pe website: <http://www.brukenthalmuseum.ro/publicatii/01.htm>

Please send any mail or messages regarding this publication at:

Brukenthal National Museum – Piața Mare 4–5, 550163, Sibiu. **Phone number:** 004/369/101782; **Fax:** 004/ 269 /211545.

E-mail: ioan.tausan@brukenthalmuseum.ro; Website: www.brukenthalmuseum.ro

The entire responsibility for the specialized information of the article's content is to be assumed by the author; all materials will be submitted to a peer review process. The details can be found at http://www.brukenthalmuseum.ro/publicatii_en/01.htm.

The guide for the authors can be found at: http://www.brukenthalmuseum.ro/publicatii_en/01.htm

TABLE OF CONTENTS

ENTOMOMOLOGY

Sergiu-Cornel TÖRÖK, <i>Contributions to the study of Heterocera (Lepidoptera) from Mediaş and Surroundings (Sibiu County).....</i>	441
Sergiu-Cornel TÖRÖK, Gabriela CUZEPAN, <i>Data regarding genus Parnassius Latreille, 1804 (Lepidoptera: Papilionidae) preserved in Natural History Museum collections from Sibiu.....</i>	459
Gabriel LUPU, Mihai DOROFTEI, <i>Asiotmethis limbatus (Charpentier, 1845) (Orthoptera: Pamphagidae) in Romania – specific habitats.....</i>	473
Ionuț Ștefan IORGU, <i>First record of Isophya pienensis in the Western Romanian Carpathians (Orthoptera: Phaneropteridae).....</i>	481
Alexandru Ioan TATU, Ioan TĂUȘAN, <i>Armoured crickets (Orthoptera: Tettigoniidae, Bradyporinae) in the Natural History Museum Collections Of Sibiu (Romania).....</i>	487
Ioan TĂUȘAN, Mădălina Maria JERPEL, Ioana Roxana PUȘCAȘU, Roberta Elena BRUTARU, Cătălin SADEANU, Lavinia Maria RĂDUȚIU, Valeria GIURESCU, <i>Ant Fauna (Hymenoptera: Formicidae) of Sibiu County (Transylvania, Romania).....</i>	499

MALACOLOGY

Ana-Maria MESAROȘ, Raluca STOICA, <i>The Bielz Malacological Slug Collection from the Natural History Museum in Sibiu</i>	521
Voichița GHEOCA, <i>Helicids in land snail communities from secondary grasslands of Southern Transylvania.....</i>	533
Voichița GHEOCA, <i>Considerations regarding land snails diversity in urban areas. Case study - The city of Sibiu.....</i>	541

VERTEBRATES

Alexandru Nicolae STERMIN, Liviu Răzvan PRIPON, Alin DAVID, <i>The importance of homogeneous vs. heterogenous wetlands in rallid (Rallidae) phenological seasons.....</i>	549
Ana Maria LAZĂR, Ana Maria BENEDEK, Georgiana MĂRGINEAN, <i>Small mammals (Mammalia: Rodentia et Insectivora) from the Upper Tisa river basin (Romania).....</i>	555

BOTANY

Ghizela VONICA, Aurelia HOROTAN, <i>Herbarium vivum: A historical collection of medicinal plants from the Natural History Museum (Sibiu, Romania).....</i>	569
Enikő LACZI, Alexandru Silviu APAHIDEAN, <i>Protected culture study of chinese cabbage (Brassica campestris var. pekinensis) varieties and hybrids collection grown in the Transylvanian Tableland specific conditions.....</i>	579
Kinga ÖLLERER, <i>The flora of the Breite wood-pasture (Sighișoara, Romania).....</i>	589

GEOLOGY & GEOMORPHOLOGY

Marioara COSTEA, Rodica CIOBANU, <i>Geological and hydrological considerations in the Secașul Mare Basin (South – Western part of the Transylvanian Depression).....</i>	605
Marioara COSTEA, <i>Morphological and hydrological dysfunctions in the urban space of Sibiu municipality.....</i>	621

PALEONTOLOGY

Rodica CIOBANU, Nicolae TRIF, <i>Diodontidae (Osteichthyes) from „The Turnu Roşu (Romania) Eocene Limestone” Reserve</i>	631
--	-----

SHORT COMMUNICATIONS

Rodica CIOBANU, <i>Scientist of his time – Johann Michael Ackner (1782-1862)</i>	643
Rodica CIOBANU, <i>Marioara Costea – Mapping and geomorphologic analysis – Review</i>	647
Ana-Maria MESAROŞ, <i>The Miklos Szekeres donation to the Malacological collection from the Natural History Museum in Sibiu</i>	649

CONTRIBUTIONS TO THE STUDY OF HETEROCERA (LEPIDOPTERA) FROM MEDIAȘ AND SURROUNDINGS (SIBIU COUNTY)

Sergiu-Cornel TÖRÖK*

Abstract. 222 *Heterocera* (Lepidoptera) species were recorded from the surroundings of Mediaș locality, a writable white spot in the Romanian lepidopterological picture. This paper also offers an ecological and zoogeographical description of the Lepidoptera from this area. In addition, it points out the endangered species, with the use of Romanian Red List study. From the species present in this area the most remarkable are: *Phragmatobia luctifera* (Denis & Schiffermüller, 1775), *Dichagyris signifera* (Denis & Schiffermüller, 1775), *Thyris fenestrella* (Scopoli, 1763), *Proserpinus proserpina* (Pallas, 1772); *Acherontia atropos* (Linnaeus, 1758); *Hyles gallii* (Rottemburg 1775), *Eucarta virgo* (Treitschke, 1835) and *Eucarta amethystina* (Hubner, 1803).

Keywords: *Macrolepidoptera, Heterocera, Mediaș, Curciu, Bazna, faunistics, ecology, zoogeography*

Rezumat. 222 specii aparținând subordinului *Heterocera* (Lepidoptera) au fost identificate din zona localității Mediaș, această zonă reprezentând o veritabilă “pată albă” în tabloul lepidopterologic românesc. Lepidopterofauna acestei zone este caracterizată din punct de vedere zoogeografic și ecologic. Speciile periclitare sunt scoase în evidență prin analiza speciilor cuprinse în Lista Roșie a României. Astfel, printre speciile remarcabile prezente în această zonă se numără: *Phragmatobia luctifera* (Denis & Schiffermüller, 1775), *Dichagyris signifera* (Denis & Schiffermüller, 1775), *Thyris fenestrella* (Scopoli, 1763), *Proserpinus proserpina* (Pallas, 1772), *Acherontia atropos* (Linnaeus, 1758), *Hyles gallii* (Rottemburg 1775), *Eucarta virgo* (Treitschke, 1835) și *Eucarta amethystina* (Hubner, 1803).

Cuvinte cheie: *Macrolepidoptera, Heterocera, Mediaș, Curciu, Bazna, faunistică, ecologie, zoogeografie*

Introduction

Mediaș and his surrounding villages are located in the middle basin of Târnava Mare River, in the southern part of Târnavelor Plain. This region is characterized by average height hills (250 m – 500 m) divided by wide valleys with well-developed terraces (Sorocovschi 1992).

The soils of The Târnavelor Plain are very diverse, varying from brown forest soils to alluvial clay soils found on the riverbanks (Badea *et al.* 1981).

The climate of this region is temperate-continental with an average annual temperature spreading from 8.4°C at Dumbrăveni weather station (at 318 m altitude) to 9.2°C at Blaj weather station (at 334 m altitude) and an average amount of precipitations of 600 – 700 mm (Sorocovschi 1992).

The vegetation presents a great diversity of habitats. In the valley we encounter alder and willow associations, as well as hygrophilous grasslands. Going up, well-developed terraces are covered by mesohygrophilous and mesophilous

grasslands. In a small proportion secondary steppe-like lawns are spread. At the tops of the hills, oak and beech forests and shrub associations as *Pruno spinosae-Crataegetum* Hueck, 1931 (Badea *et al.* 1981) are spread.

The first documented collection of Lepidoptera made in present day Mediaș was a series of Lepidoptera specimens collected by Daniel Czekelius, who at the end of the 19th century, in The First Catalogue of the Transylvanian Lepidoptera, cited two endangered species, *Daphnis nerii* (Linnaeus, 1758) and *Orgyia recens* (Hübner, 1819) (Czekelius 1897). He also made in 1912, 1913, 1917 and 1921 several trips to Bazna, citing from here, 11 species of Lepidoptera (Burnaz 1993).

After Czekelius, Ostrogovich outlook a field trip in 1921 also at Bazna, and cited the presence of 19 species of Lepidoptera species (Popescu-Gorj 1964).

The most recent data is reported by Rákossy who sampled from Târnava Mare Valley, including Mediaș and Bane. Among the rare species from

* Babeș-Bolyai University of Cluj-Napoca, Romania, ser.torok@yahoo.com

this region, he mentioned: *Nola cristatula* (Hübner, 1793), *Yigoga signifera* (Denis & Schiffermüller, 1775), *Athetis furvula* (Hübner, 1808), *Sideridis turbida* Esper, 1790 and many others (Rákossy 1996).

This study also contains information regarding the Lepidoptera from Mediaş and Moşna found in Eckbert Schneider Entomological Collection, part of the Entomological Collection of the Natural History Museum from Sibiu.

Overall, this region's lepidopterofauna is insufficiently investigated; as a result, one of the aims of this study is to add supplementary data to the Lepidoptera fauna from Mediaş area. Another purpose is to characterize this region's moth fauna from the zoogeographical and the ecological point of view.

Material and methods

The study of Macroheterocera species was carried out during 2008-2011. We used an entomological net for some day-flying moths and 125 W light traps and 8V portable traps for the night-flying Lepidoptera. Additional sugar traps (early in the spring and late in the autumn) were used in order to attract moths that are not (well) attracted to light. Forest edges (Mediaş: Vewern, Greweln, Binderbubi, Izvorul Dorului), clearings, damp meadows (Târnava Mare riverbank), ruderal areas (Curciu, village) and mezophilous grasslands (Curciu, Cărairi Hill), were the main studied habitats.

The identification of the lepidopterological material was made using several identification keys: Rákossy (1996), Koch (1958, 1961), Malicky *et al.* (2000) and Székely (2010). The systematic list was compiled according to the one present in The Catalogue of the Romanian Lepidoptera (Rákossy *et al.* 2003) and Karsholt and Nieuwerkerken (2011). The zoogeographical and ecological character was written using Romanian faunistics studies (Burnaz 1995, Rákossy 1996, Cremene *et al.* 2003, Dincă 2006, Székely 2006, Burnaz, 2008, 2009, Székely 2010).

The red list was made following the one proposed by Rákossy *et al.* (2003) according to IUCN 2001, which point out the conservation status of Lepidoptera species at the country's level.

Results and discussions

As the result of the Macroheterocera investigation, a number of 222 species were found at Mediaş and its surroundings (Bazna, Curciu, Moşna). The checklist of the recorded Macrolepidoptera with their zoogeographical, ecological and red list status is given in table 1.

The family spectrum of species collected by the light traps is shown in figure 1. Noctuidae (88 species) and Geometridae (68 species) are represented with the highest numbers of species, but the species of Erebididae, and Sphingidae showed a significant proportion as well.

The other aspect of the high species diversity is the occurrence of several faunal types. Their composition showed that 82.59% of the collected species are Eurasiatic (185 species, from the total number of species), 9.38% West-Asiatic Mediterranean, 2.68% Holarctic, 2.23% Subtropical, Cosmopolit and 1.34% Ponto-Mediterranean (Fig. 2).

The dominance of Eurasiatic elements represents the consequence of the geographical position of the study site and of Romania in general. This can also explain the presence of 21 West-Asiatic Mediterranean species (9.38%) which is a relative small number in comparison with the proportion from the Danube Delta, 28% (Székely 2006). In fact Dobrogea contains the highest proportion of West-Asiatic Mediterranean and West-Asiatic elements in Romania (Rákossy, Székely 1996, Dincă 2006).

The majority of species prefer the mezophilous habitats (47%). Many species prefer the deciduous forest or deciduous forest scrub associations from Vewern, Greweln, Binderbubi and Izvorul Dorului, like: *Agria tau* (Linnaeus, 1758), *Cyclophora annularia* (Fabricius, 1775), *Selenia lunularia* (Hübner, 1788), *Lithophane ornitopus* (Hufnagel, 1766), *Orthosia cerasi* (Fabricius, 1775), *Orgyia antiqua* (Linnaeus, 1758) or *Spatalia argentina* (Denis & Schiffermüller, 1775). Others species such as *Macrothylacia rubi* (Linnaeus, 1758), *Laothoe populi* (Linnaeus, 1758), *Archanara neurica* (Hübner, 1808), *Archanara dissoluta* (Treitschke, 1825), *Furcula bifida* (Brahm, 1787), *Clostera curtula* (Linnaeus, 1758), *Eucarta amethystina* (Hubner, 1803), *Eucarta virgo* (Treitschke, 1835) are characteristic for the mesohygrophilous or hygrophilous pastures

(13.56%), found on the Târnava Mare riverbanks (Fig. 3).

On the south facing slopes, such as Viile Pusti (Curciu), we can find a high number of xerothermophilous, thermophilous, meso - thermophilous or mezoxerophilous species; these species represent 22.43% from the total species identified.

In this region are also present 15 species of migratory Lepidoptera. From these the most notable is the presence of subtropical sphingids, *Acherontia atropos* (Linnaeus, 1758), *Hyles gallii* (Rottemburg, 1775), *Hyles euphorbiae* (Linnaeus, 1758) and *Daphnis nerii* L. This last species has not been seen in the region from more than 50 years, the last sighting was made by Rákossy and Weber in Sighișoara in 1966 (Rákossy, Weber 1986).

The study of the Romanian Red List reveals (Rákossy *et al.* 2003) a high number of protected species, 33 of them being endangered at some level (Fig. 4).

The most important aspect is represented by the presence of two endangered species: *Phragmatobia luctifera* (Denis & Schiffermüller,

1775) and *Dichagyris signifera* (Denis & Schiffermüller, 1775), both found at Bazna and six vulnerable species, they are *Thyris fenestrella* (Scopoli, 1763); *Proserpinus proserpina* (Pallas, 1772); *Acherontia atropos* (Linnaeus, 1758); *Hyles gallii* (Rottemburg 1775); *Eucarta virgo* (Treitschke, 1835) and *Eucarta amethystina* (Hubner, 1803).

24 near threatened species are as well present in this area (Table 1, Fig. 1).

Acknowledgements

The author wants to thank to the anonymous referees for their valuable and helpful comments on the manuscript. The author is grateful for the comments of Prof. Dr. László Rákossy and Ioan Tăușan for the linguistic support. Special thanks to Petre Crețu, Nicolae Crețu and Cornel Török for their help in the collecting campaigns.

This work was possible with the financial support of the Sectoral Operational Programme for Human Resources Development 2007-2013, cofinanced by the European Social Fund, under the project number POSDRU/107/1.5/S/76841 with the title „Modern Doctoral Studies: Internationalization and Interdisciplinarity”.

REFERENCES

- Badea *et al.* 1981 Badea Lucian, Caloianu Nicolae, Dragu Gheorghe, Tufescu Victor, Gâstescu Petre, Velcea Ioan, *Județul Sibiu: Monografie*. In *Editura Academiei Republicii Socialiste România* (1971) București.
- Burnaz 1993 Burnaz Silvia, *Catalogul colecției de lepidoptere a Muzeului județean Deva*. In: *Sargetia. Acta Musei Deva, Seria Științele Naturii* 14-15 (1993) Deva, p. 157-302.
- Burnaz 1995 Burnaz Silvia, *Considerații ecologice și zoogeografice privind fauna de macrolepidoptere a Cheilor Crivădiei*. In: *Buletin informativ Societatea lepidopterologică română* 6(1-2) (1995) Cluj-Napoca, p. 30-50.
- Burnaz 2008 Burnaz, Silvia, *Macrolepidoptere din Munții Șureanului*. In *Editura Tipografia Deva* (2008) Deva.
- Burnaz 2009 Burnaz Silvia, *Macrolepidoptere din Munții Poiana Ruscă (Carpații Occidentali, România)*. In: *Brukenthal Acta Musei* IV.3 (2009) Sibiu, p. 645-660.
- Cremene *et al.* 2003 Cremene Cristina, Rákósy László, Erhardt Andreas, *Diversity of Macrolepidoptera in steppe habitats of Caianu Mic (Cluj, Romania)*. In: *Entomologica Romanica* 7 (2003) Cluj-Napoca, p. 5-14.
- Czekelius 1897 Czekelius Daniel, *Kritisches Verzeichnis der Schmetterlinge Siebenbürgens*. In: *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* 47 (1897) Sibiu, p. 1-78.
- Dincă 2006 Dincă Vlad, *The Macrolepidoptera (Insecta: Lepidoptera) from Istrița Hill (Buzău County, Romania)*. *Entomologica Romanica* 10 (2006) Cluj-Napoca, p. 5-24.
- Koch 1958 Koch, Manfred, Heinicke, Wolfgang, Müller, Bernd, *Wir bestimmen schmetterlinge, Band 3, Eulen*. In *Neumann Verlag Radebeul* (1958) Leipzig.
- Koch 1961 Koch, Manfred, *Wir bestimmen schmetterlinge, Band 4, Spinner*. In *Neumann Verlag Radebeul* (1961) Leipzig.
- Malicky *et al.* 2000 Malicky Michael, Hauser Ervin, Huemer Peter, Wieser Christian, *Verbreitungsatlas der Tierwelt Österreichs: Noctuidae sensu classico*. In *Staphia* 74 (2000) Linz.
- Karsholt, Nieuwerkerken 2011 Karsholt Ole, Nieuwerkerken Erik van, *Lepidoptera, Moths. Fauna Europaea version 2.4*, <http://www.faunaeur.org> (2011).
- Popescu-Gorj 1964 Popescu-Gorj, Aurelian, *Catalogue de la collection de lépidoptères „Prof. A. Ostrogoșich” du Museum d’Histoire Naturelle „Grigore Antipa”, Bucharest*. In *Editura Muzeului „Grigore Antipa”* (1964) Bucharest.
- Rákósy 1996 Rákósy, László, *Die Noctuiden Rumäniens*. In *Staphia* 46 (1996) Linz.
- Rákósy *et al.* 2003 Rákósy László, Goia Marin, Kovács Zoltan, *Verzeichnis der Schmetterlinge Rumäniens. Catalogul lepidopterelor din România*. In: *Editura Societății Lepidopterologice Române* (2003) Cluj-Napoca.
- Rákósy, Székely 1996 Rákósy László, Székely Levente, *Die Makrolepidopteren der Süddobrudscha (Lepidoptera, Südostrumänien)*. In: *Entomologica Romanica* 1 (1996) Cluj-Napoca, p. 17 - 62.
- Rákósy, Weber 1986 Rákósy László, Weber Wilhelm, *Die Großschmetterlinge von Sighisoara (Schässburg) und Umgebung (Siebenbürgen, Rumänien)*. In: *Atalanta* 16 (1986) Würzburg, p. 315-392.
- Sorocovschi 1992 Sorocovschi Victor, *Podișul Târnavelor., Studiu Hidrogeografic*. In *Cetib* (1992) Cluj-Napoca.
- Székely 2006 Székely Levente, *Fluturii din Delta Dunării - The Butterflies and moths of the Danube Delta*. In *Editura Disz – Tipo* (2006) Săcele.
- Székely 2010 Székely Levente, *Moths of Romania 1 – Fluturi de noapte din Romania 1, - Hepialidae, Limacodidae, Cossidae, Thyrididae, Lasiocampidae, Endromidae, Saturniidae, Lemoniidae, Sphingidae, Drepanidae, Thaumetopoeidae, Notodontidae, Pantheidae, Lymantriidae, Arctiidae*. In *Editura Disz – Tipo* (2010) Săcele.

LIST OF ILLUSTRATIONS

Fig. 1. The representation of the suborder Heterocera from Mediaş and surroundings

Fig. 2. Zoogeographical spectrum of the Lepidoptera species identified in Mediaş and surroundings

Fig. 3. Ecological spectrum of the Lepidoptera species identified in northern part of Sibiu County

Fig. 4. The representation of IUCN categories after Rákósy *et al.* (2003) for the Mediaş region

Tab.1. Systematic list of the Macrolepidoptera from Mediaş and surroundings (Sibiu County, Romania)/

Abbreviations head of the table – Loc. = Localities; Z. El. = zoogeographical element; E. C. = ecological character; R. L. = red list; **Z. El.** – Eua = Eurasiatic; Vam = West-Asiatic Mediterranean; E = European; Hol = Holarctic; Str = Subtropical; Cosm = Cosmopolit; Pm = Ponto-Mediterranean. **E. C.** – M = mezophilous; Mx = mezoxerophilous; Mxt = mezoxerothermophilous; Xt = xerothermophilous; T = thermophilous; Mt = mezothermophilous; Mh = mezohygrophilous; Mht = mezohygrothermophilous; Hg = hygrophilous; Eu = euribiont; Mg = migratory; **L.** – NT = near threatened; VU = vulnerable; EN = endangered ER = erratic

LISTA ILUSTRAȚIILOR

Fig. 1. Reprezentarea familiilor de lepidoptere din subordinul Heterocera din Mediaş și împrejurimi

Fig. 2. Spectrul distribuției geografice pentru Heterocerele din Mediaş și împrejurimi

Fig. 3. Spectrul cerințelor ecologice ale speciilor de lepidoptere din zona investigată (Mediaş și împrejurimi)

Fig. 4. Categoriile IUCN pentru Heterocerele din zona Mediaş după Rákósy *et al.* (2003)

Tab.1. Lista sistematică a speciilor de macrolepidoptere identificate în localitatea Mediaş și împrejurimi (județul Sibiu, Romania)

Abrevieri: cap tabel; Loc. = Localități; Z. El. = element zoogeografic; E. C. = caracter ecologic; R. L. = lista roșie. **Z. El.** – Eua = Eurasiatic; Vam = West-Asiatic Mediterranean; E = European; Hol = Holarctic; Str = Subtropical; Cosm = Cosmopolit; Pm = Ponto-Mediterranean. **E. C.** – M = mezofil; Mx = mezoxerofil; Mxt = mezoxerotermofil; Xt = xerotermofil; T = termofil; Mt = mezotermofil; Mh = mezohigrofil; Mht = mezohigrotermofil; Hg = higrofil; Eu = euribiont; Mg = migrator. **R. L.** – NT = taxon potențial amenințat; VU = taxon vulnerabil; EN = taxon periclitat; ER = eratic.




Fig. 1. The representation of the suborder Heterocera from Mediaș and surroundings




Fig. 2. Zoogeographical spectrum of the Lepidoptera species identified in Mediaș and surroundings




Fig. 3. Ecological spectrum of the Lepidoptera species identified in northern part of Sibiu County




Fig. 4. The representation of IUCN categories after Rákossy *et al.* (2003) for the Mediaș region

Table 1 – Systematic list of the Macrolepidoptera from Mediaş and surroundings (Sibiu County, Romania)

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
Superfamily Hepialoidea					
Family Hepialidae					
1	<i>Triodia sylvina</i> (Linnaeus, 1758)	Curciu (Viile Pusti),	Eua	M	
Superfamily Zygaenoidea					
Family Zygaenidae					
Subfamily Procridinae					
2	<i>Adscita statices</i> (Linnaeus, 1767)	Mediaş (Izvolul Dorului),	Eua	M	
Subfamily Zygaeninae					
3	<i>Zygaena carniolica</i> (Scopoli, 1763)	Mediaş (Izvolul Dorului, Târnava Mare bank), Curciu (Căări Hill),	Eua	Mxt	
4	<i>Zygaena loti</i> (Denis & Schiffermüller, 1775)	Curciu (Căări Hill),	Eua	M	
5	<i>Zygaena ephialtes</i> (Linnaeus, 1767)	Mediaş (Târnava Mare bank), Curciu (Căări Hill),	Eua	Mxt	
6	<i>Zygaena purpuralis</i> (Brünnich, 1763)	Bazna (Burnaz, 1993),	Eua	Mxt	
7	<i>Zygaena filipendulae</i> (Linnaeus, 1758)	Bazna (Burnaz, 1993), Mediaş (Târnava Mare bank, Izvolul Dorului), Curciu (Căări Hill),	Eua	M	
Family Limacodidae					
Subfamily Limacodinae					
8	<i>Apoda limacodes</i> (Hufnagel, 1766)	Mediaş (Binderbubi),	Eua	M	
Superfamily Cossoidea					
Family Cossidae					
Subfamily Zeuzerinae					
9	<i>Zeuzera pyrina</i> (Linnaeus, 1761)	Curciu (Căări Hill, village),	Eua	Mh	
Superfamily Thyridoidea					
Family Thyrididae					
Subfamily Thyridinae					
10	<i>Thyris fenestrella</i> (Scopoli, 1763)	Mediaş (Târnava Mare bank),	Eua	Mt	VU
Superfamily Bombycoidea					
Family Lasiocampidae					
Subfamily Lasiocampinae					
11	<i>Macrothylacia rubi</i> (Linnaeus, 1758)	Bazna (Burnaz, 1993), Mediaş (Târnava Mare bank),	Eua	M; Mh	NT
12	<i>Lasiocampa quercus</i> (Linnaeus, 1758)	Curciu (Căări Hill),	Eua	M	
Subfamily Pinarinae					
13	<i>Gastropacha quercifolia</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	NT
Family Saturniidae					
Subfamily Agliinae					
14	<i>Aglia tau</i> (Linnaeus, 1758)	Mediaş (Binderbubi),	Eua	M	
Superfamily Sphingoidea					
Family Sphingidae					

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
Subfamily Smerinthinae					
15	<i>Laothoe populi</i> (Linnaeus, 1758)	Curciu (village),	Eua	Mh	
Subfamily Sphinginae					
16	<i>Agrius convolvuli</i> (Linnaeus, 1758)	Mediaş (Binderbubi), Curciu (village, Cărați Hill),	Str	Mg, Mx	
17	<i>Acherontia atropos</i> (Linnaeus, 1758)	Curciu (potato plantation, village),	Str	Mg, Mx	VU
Subfamily Macroglossinae					
18	<i>Macroglossum stellatarum</i> (Linnaeus, 1758)	Mediaş (town, Izvorul Dorului, Târnava Mare bank), Curciu (village),	Eua	Mx, Mg	
19	<i>Daphnis nerii</i> (Linnaeus, 1758)	Mediaş (Czekelius, 1897),	Str	Mg	ER
20	<i>Proserpinus proserpina</i> (Pallas, 1772)	Curciu (garden),	Vam	Xt	VU
21	<i>Hyles gallii</i> (Rottemburg 1775)	Curciu (village),	Hol	Mxt, Mg	VU
22	<i>Hyles euphorbiae</i> (Linnaeus, 1758)	Curciu (village),	Eua	Mx, Mg	NT
23	<i>Deilephila elpenor</i> (Linnaeus, 1758)	Curciu (village),	Eua	Mh	NT
24	<i>Deilephila porcellus</i> (Linnaeus, 1758)	Mediaş (Târnava Mare bank), Curciu (Cărați Hill, village),	Eua	M	
Superfamily Drepanoidea					
Family Drepanidae					
Subfamily Thyatirinae					
25	<i>Thyatira batis</i> (Linnaeus, 1758)	Curciu (village), Mediaş (Târnava Mare bank, Binderbubi),	Eua	Mh	
26	<i>Habrosyne pyritoides</i> (Hufnagel, 1766)	Mediaş (Târnava Mare bank, Binderbubi), Curciu (Cărați Hill),	Eua	M, Mh	
27	<i>Tethea or</i> (Denis & Schiffermüller, 1775)	Curciu (village), Mediaş (Greweln),	Eua	Mh	NT
Subfamily Drepaninae					
28	<i>Cilix glaucata</i> (Scopoli, 1763)	Mediaş (Greweln, Târnava Mare bank), Curciu (village, Cărați Hill, Viile Pusti),	Eua	Xt	
29	<i>Sabra harpagula</i> (Esper, 1786)	Mediaş (Târnava Mare bank),	Eua	Mht	NT
Superfamily Noctuoidea					
Family Notodontidae					
Subfamily Notodontinae					
30	<i>Furcula bifida</i> (Brahm, 1787)	Curciu (village),	Vam	Mh	
31	<i>Ptilodon cucullina</i> (Denis & Schiffermüller, 1775)	Mediaş (Binderbubi),	Eua	Mh	
32	<i>Clostera pigra</i> (Hufnagel, 1766)	Curciu (village),	Eua	Mh	
33	<i>Clostera curtula</i> (Linnaeus, 1758)	Curciu (Cărați Hill),	Eua	Mh	
Subfamily Phalerinae					
34	<i>Phalera bucephala</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	
Subfamily Heterocampinae					

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
35	<i>Spatalia argentina</i> (Denis & Schiffermüller, 1775)	Curciu (village),	Eua	Mt	
Family Erebiidae					
Subfamily Lymantriinae					
36	<i>Calliteara pudibunda</i> (Linnaeus 1758)	Curciu (village),	Eua	M	
37	<i>Orgyia antiqua</i> (Linnaeus, 1758)	Mediaş (Greweln),	Hol	M	
38	<i>Euproctis chrysorrhoea</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	
39	<i>Penthopha morio</i> (Linnaeus, 1758)	Curciu (Cărați Hill), Mediaş (Izvorul Dorului),	Eua (Pm)	M	NT
Subfamily Arctiinae					
40	<i>Miltochrista miniata</i> (Forster, 1771)	Curciu (village), Mediaş (Binderbubi, Târnava Mare bank),	Eua	M	
41	<i>Atolmis rubricollis</i> (Linnaeus, 1758)	Bazna (Burnaz, 1993),	Eua	M	
42	<i>Lithosia quadra</i> (Linnaeus, 1758)	Curciu (Cărați Hill, village), Mediaş (Târnava Mare bank),	Eua	M	
43	<i>Eilema sororcula</i> (Hübner, 1808)	Curciu (village), Mediaş (Greweln),	Eua	Mh	
44	<i>Eilema pygmaeola</i> (Doubleday, 1847)	Curciu (Viile Pusti),	Eua	M	
45	<i>Eilema complana</i> (Linnaeus, 1758)	Curciu (village),	Eua	Mt	
46	<i>Amata phegea</i> (Linnaeus, 1758)	Curciu („Râpi” forest edge), Mediaş (Izvorul Dorului, Târnava Mare bank),	Eua	M	
47	<i>Phragmatobia fuliginosa</i> (Linnaeus, 1758)	Curciu (Cărați Hill, village), Mediaş (Izvorul Dorului, Târnava Mare bank),	Eua	M	
48	<i>Phragmatobia luctifera</i> (Denis & Schiffermüller, 1775)	Bazna (Burnaz, 1993),	Eua	Mt	EN
49	<i>Spilosoma lubricipeda</i> (Linnaeus, 1758)	Curciu (Cărați Hill, village), Mediaş (Târnava Mare bank, Binderbubi, Greweln),	Eua	M	
50	<i>Spilosoma lutea</i> (Hufnagel, 1766)	Mediaş (Târnava Mare bank),	Eua	M	
51	<i>Diaphora mendica</i> (Cerck, 1759)	Curciu (village), Mediaş (Târnava Mare bank),	Eua	Mh	
52	<i>Diacrisia sannio</i> (Linnaeus, 1758)	Curciu (village), Mediaş (Târnava Mare bank),	Eua	M, Mh	
53	<i>Rhyparia purpurata</i> (Linnaeus, 1758)	Curciu (village),	Eua	Mxt	
54	<i>Arctia caja</i> (Linnaeus, 1758)	Bazna (Popescu-Gorj, 1964),	Eua	M	
55	<i>Euplagia quadripunctaria</i> (Poda, 1761)	Curciu (garden),	Eua	M	
Subfamily Erebiinae					
56	<i>Catocala elocata</i> (Esper, 1787)	Curciu (village),	Vam	Mht	NT
57	<i>Euclidia glyphica</i> (Linnaeus, 1758)	Curciu (Viile Pusti, Cărați Hill), Mediaş (Târnava Mare bank),	Eua	Mt	

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
		Izvorul Dorului, Binderbubi, Greweln),			
58	<i>Lygephila pastinum</i> (Treitschke, 1826)	Mediaş (Târnava Mare bank),	Eua	T	
Subfamily Herminiinae					
59	<i>Polypogon tentacularia</i> (Linnaeus, 1758)	Bazna (Burnaz, 1993), Curciu, Mediaş (Târnava Mare bank, Izvorul Dorului),	Eua	M	
60	<i>Zanclognatha tarsipennalis</i> Treitschke, 1835	Bazna (Popescu-Gorj, 1964),	Eua	Mh	
Subfamily Aventiinae					
61	<i>Laspeyria flexula</i> (Germar, 1810)	Curciu (village), Mediaş (Greweln),	Eua	M	
Subfamily Hypeninae					
62	<i>Hypena proboscidalis</i> (Linnaeus, 1758)	Bazna (Popescu-Gorj, 1964), Curciu (village), Mediaş (Târnava Mare bank),	Eua	Mh	
63	<i>Hypena rostralis</i> (Linnaeus, 1758)	Bazna (Popescu-Gorj, 1964), Curciu (village, Că rări Hill), Mediaş (Vewern),	Eua	M	
Family Nolidae					
Subfamily Nolinae					
64	<i>Nola aerugula</i> (Hübner, 1808)	Mediaş (Târnava Mare bank),	Eua	Mh	
Subfamily Chloephorinae					
65	<i>Earias clorana</i> (Linnaeus, 1761)	Mediaş (Târnava Mare bank),	Eua	Mh	
66	<i>Pseudoips prasinana</i> (Linnaeus, 1758)	Curciu (village, Că rări Hill), Mediaş (Târnava Mare bank),	Eua	M	
Family Noctuidae					
Subfamily Pantheinae					
67	<i>Colocasia coryli</i> (Linnaeus, 1758)	Curciu (village)	Eua	M	
Subfamily Noctuinae					
68	<i>Agrotis segetum</i> (Denis & Schifferrmüller, 1775)	Curciu (village, Că rări Hill),	Eua	M	
69	<i>Agrotis exclamationis</i> (Linnaeus, 1758)	Curciu (village, Că rări Hill), Mediaş (Târnava Mare bank, Binderbubi, Greweln),	Eua	Eu, Mg	
70	<i>Agrotis ipsilon</i> (Hufnagel, 1766)	Curciu (village),	Cosm	Eu, Mg	
71	<i>Axylia putris</i> (Linnaeus, 1761)	Curciu (village, Că rări Hill), Mediaş (Târnava Mare bank, Binderbubi),	Eua	M	
72	<i>Ochropleura plecta</i> (Linnaeus, 1761)	Curciu (Că rări Hill), Mediaş (Binderbubi, Târnava Mare bank),	Hol	M	
73	<i>Dichagyris signifera</i> (Denis & Schifferrmüller, 1775)	Bazna (Rakosy, 1996),	Eua	Mxt, Xt	EN
74	<i>Noctua pronuba</i> (Linnaeus, 1758)	Curciu (village), Mediaş (Greweln),	Eua	M, Mg	
75	<i>Noctua fimbriata</i> (Schreber, 1759)	Curciu (village),	Vam	M, Mg	
76	<i>Xestia c-nigrum</i>	Curciu (village),	Cosm	Eu,	

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
	(Linnaeus, 1758)			Mg	
77	<i>Xestia ditrapezium</i> (Denis & Schiffermüller, 1775)	Curciu (village), Mediaş (Binderbubi),	Eua	M	
78	<i>Cerastis rubricosa</i> ([Denis & Schiffermüller], 1775)	Curciu (village),	Eua	M	
79	<i>Anarta trifolii</i> (Hufnagel, 1766)	Curciu (village),	Hol	Eu	
80	<i>Mamestra brassicae</i> (Linnaeus, 1758)	Curciu (village),	Eua	M, Eu	
81	<i>Melanchra persicariae</i> (Linnaeus 1761)	Curciu (Căzări Hill),	Eua	M	
82	<i>Thalpopbila matura</i> (Hufnagel, 1766)	Curciu (village),	Vam	M	
83	<i>Lacanobia contigua</i> (Denis & Schiffermüller, 1775)	Curciu (Căzări Hill),	Eua	M	
84	<i>Lacanobia suasa</i> (Denis & Schiffermüller, 1775)	Curciu (village), Mediaş (Târnava Mare bank),	Eua	Mh	
85	<i>Lacanobia oleracea</i> (Linnaeus, 1758)	Curciu (village),	Eua	Mht	
86	<i>Hadena perplexa</i> ([Denis & Schiffermüller], 1775)	Târnava (Rakosy, 1996),	Eua	Xt	
87	<i>Tholera decimalis</i> (Poda, 1761)	Curciu (village),	Eua	M	
88	<i>Anorthoa munda</i> (Denis & Schiffermüller, 1775)	Curciu (village), Mediaş (Vewern, Greweln),	Eua	M	
89	<i>Orthosia incerta</i> (Hufnagel, 1766)	Curciu (village),	Eua	M	
90	<i>Orthosia cruda</i> (Denis & Schiffermüller, 1775)	Curciu (village),	Vam	M	
91	<i>Orthosia gracilis</i> (Denis & Schiffermüller, 1775)	Curciu (village),	Eua	M	
92	<i>Orthosia cerasi</i> (Fabricius, 1775)	Curciu (village),	Eua	M	
93	<i>Orthosia gothica</i> (Linnaeus, 1758)	Curciu (village), Mediaş (Greweln),	Eua	M	
94	<i>Mythimna turca</i> (Linnaeus, 1758)	Bazna (Popescu-Gorj, 1964), Curciu (village, Căzări Hill), Mediaş (Târnava Mare bank, Binderbubi),	Eua	Mh	
95	<i>Mythimna pudorina</i> ([Denis & Schiffermüller], 1775)	Mediaş (Târnava Mare bank),	Eua	Mx	
96	<i>Mythimna ferrago</i> (Fabricius, 1787)	Curciu (village),	Eua	M, Mh	
97	<i>Mythimna vitellina</i> (Hübner, 1808)	Curciu (village),	Vam	Mxt, Xt, Mg	
98	<i>Mythimna l-album</i> (Linnaeus, 1767)	Curciu (village),	Eua	Eu, Mg	
99	<i>Mythimna pallens</i>	Curciu (village, Căzări Hill),	Vam	M,	

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
	(Linnaeus, 1758)	Mediaş (Târnava Mare bank),		Mh	
100	<i>Mythimna albipuncta</i> (Denis & Schiffermüller, 1775)	Curciu (village, Viile Pusti), Mediaş (Binderbubi, Târnava Mare bank),	Vam	M	
101	<i>Lithophane ornitopus</i> (Hufnagel, 1766)	Curciu (village),	Eua	M	
102	<i>Eupsilia transversa</i> (Hufnagel, 1766)	Curciu (village), Mediaş (Vewern, Greweln),	Eua	M	
103	<i>Agrochola helvola</i> (Linnaeus, 1758)	Curciu (village),	Eua	Mt	
104	<i>Agrochola circumcellaris</i> (Hufnagel, 1766)	Curciu (village),	Eua	M	
105	<i>Conistra vaccinii</i> (Linnaeus, 1761)	Mediaş (Vewern, Greweln),	Eua	M	
106	<i>Tiliacea aurago</i> ([Denis & Schiffermüller], 1775)	Curciu (village),	Eua	M	
107	<i>Polyphaenis sericata</i> (Esper, 1787)	Curciu (village),	Vam	M	
108	<i>Trachea atriplicis</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	
109	<i>Euplexia lucipara</i> (Linnaeus, 1758)	Curciu (village), Mediaş (Târnava Mare bank),	Eua	M, Mh	
110	<i>Phlogophora meticulosa</i> (Linnaeus, 1758)	Curciu (village),	Vam	M, Mh	
111	<i>Cosmia trapezina</i> (Linnaeus, 1758)	Bazna (Col. Schneider),	Vam	M	
112	<i>Actinotia polyodon</i> (Clerck, 1759)	Curciu (village),	Eua	Mxt	
113	<i>Apamea monoglyphia</i> (Hufnagel, 1766)	Curciu (village),	Eua	Eu	
114	<i>Apamea sordens</i> (Hufnagel, 1766)	Mediaş (Târnava Mare bank),	Eua	Eu	
115	<i>Oligia strigilis</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	
116	<i>Oligia latruncula</i> (Denis & Schiffermüller, 1775)	Curciu (village),	Eua	Mh	
117	<i>Oligia versicolor</i> (Borkhausen, 1792)	Curciu, Mediaş (Târnava Mare bank),	Eua	M	
118	<i>Mesoligia furuncula</i> ([Denis & Schiffermüller], 1775)	Bazna (Popescu-Gorj, 1964),	Pm	M	
119	<i>Elaphria venustula</i> (Hübner, 1790)	Curciu (Cărări Hill), Mediaş (Târnava Mare bank, Binderbubi),	Eua	Mt	
120	<i>Mesapamea secalis</i> (Linnaeus, 1758)	Curciu (village), Mediaş (Târnava Mare bank),	Eua	M	
121	<i>Charanyca trigrammica</i> (Hufnagel, 1766)	Curciu (Cărări Hill), Mediaş (Greweln),	Vam	M	
122	<i>Athetis furvula</i> (Hübner, 1808)	Mediaş (Rakosy, 1996),	Eua	Mxt	
123	<i>Archanara neurica</i> (Hübner, 1808)	Mediaş (Czekelius, 1917),	Eua	H	
124	<i>Archanara dissoluta</i> (Treitschke, 1825)	Mediaş (Binderbubi),	Pm	H	

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
125	<i>Sideridis turbida</i> Esper, 1790	Mediaş (Rakosy, 1996),	Eua	Mxt	
126	<i>Hoplodrina ambigua</i> (Denis & Schiffermüller, 1775)	Curciu (Cărați Hill),	Vam	Mt	
127	<i>Hoplodrina octogenaria</i> (Goeze, 1781)	Curciu (village),	Eua	M	
128	<i>Egira conspicillaris</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	
Subfamily Condiinae					
129	<i>Eucarta virgo</i> (Treitschke, 1835)	Curciu (village, Cărați Hill), Mediaş (Târnava Mare bank),	Eua	Mh	VU
130	<i>Eucarta amethystina</i> (Hubner, 1803)	Curciu (Cărați Hill), Mediaş (Târnava Mare bank, Grewel),	Eua	T-Hg	VU
Subfamily Cuculliinae					
131	<i>Shargacucullia verbasci</i> (Linnaeus, 1758)	Curciu (village),	Eua	Mt	
Subfamily Amphipyrae					
132	<i>Valeria oleagina</i> (Denis & Schiffermüller, 1775)	Mediaş (town),	Vam	Mxt	
133	<i>Amphipyra pyramidea</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	
134	<i>Amphipyra livida</i> (Denis & Schiffermüller, 1775)	Curciu (village),	Eua	Mt	
Subfamily Oncocnemidinae					
135	<i>Lamprosticta culta</i> (Denis & Schiffermüller, 1775)	Curciu (village),	Vam	T	
Subfamily Acronictinae					
136	<i>Acronicta euphorbiae</i> ([Denis & Schiffermüller], 1775)	Curciu (village),	Eua	Xt	
137	<i>Acronicta rumicis</i> (Linnaeus, 1758)	Curciu (Cărați Hill)	Eua	M	
138	<i>Craniophora ligustri</i> (Denis & Schiffermüller, 1775)	Curciu (village),	Eua	Mth	
Subfamily Heliothinae					
139	<i>Heliothis peltigera</i> (Denis & Schiffermüller, 1775)	Curciu (village),	Cosm, Str	T	
140	<i>Helicoverpa armigera</i> (Hübner, 1808)	Curciu (village),	Cosm, Str	T	
141	<i>Pyrrhia umbra</i> (Hufnagel, 1766)	Curciu (Cărați Hill),	Hol	M	
Subfamily Acontiinae					
142	<i>Acontia trabealis</i> (Scopoli, 1763)	Mediaş (Târnava Mare bank, Binderbubi, Grewel), Curciu (Cărați Hill),	Eua	Mt	
143	<i>Acontia lucida</i> (Hufnagel, 1766)	Mediaş (Târnava Mare bank),	Eua	Xt	
Subfamily Eustrotiinae					
144	<i>Deltote bankiana</i> (Fabricius, 1775)	Mediaş (Târnava Mare bank, Binderbubi),	Eua	Mh	
145	<i>Deltote (Protodeltote) pygarga</i> (Hufnagel 1766)	Mediaş (Târnava Mare bank),	Eua	Mh	
Subfamily Plusiinae					
146	<i>Abrostola asclepiadis</i>	Curciu (Cărați Hill),	Vam	Mt	

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
	(Denis & Schiffermüller, 1775)				
147	<i>Abrostola triplasia</i> (Linnaeus, 1758)	Bazna (Popescu-Gorj, 1964),	Eua	M	
148	<i>Diachrysia chrysitis</i> (Linnaeus, 1758)	Curciu (village, Cărbări Hill), Mediaş (Târnava Mare bank),	Eua	M	
149	<i>Macdunnoughia confusa</i> (Stephens, 1850)	Curciu (Cărbări Hill),	Eua	Mt, Mg	
150	<i>Autographa gamma</i> (Linnaeus, 1758)	Curciu, Mediaş (Târnava Mare bank, Binderbubi),	Eua	Eu, Mg	
Subfamily Metoponiinae					
151	<i>Tyta luctuosa</i> (Denis & Schiffermüller, 1775)	Curciu (Cărbări Hill), Mediaş (Târnava Mare bank),	Eua	Xt	
Subfamily Scoliopteryginae					
152	<i>Scoliopteryx libatrix</i> (Linnaeus, 1758)	Curciu (village), Mediaş (Vewern),	Hol	M	
Subfamily Rivulinae					
153	<i>Rivula sericealis</i> (Scopoli, 1763)	Bazna (Popescu-Gorj, 1964), Curciu (village), Mediaş (Binderbubi),	Eua	Mh	
Subfamily Phytometrinae					
154	<i>Phytometra viridaria</i> (Clerck, 1759)	Curciu (Cărbări Hill), Mediaş (Binderbubi),	Eua	M	
Superfamily Geometroidea					
Family Geometridae					
Subfamily Geometrinae					
155	<i>Pseudoterpna pruinata</i> (Hufnagel, 1767)	Curciu (Viile Pusti),	Eua	Xt	
156	<i>Thetidia smaragdaria</i> (Fabricius, 1787)	Bazna (Popescu-Gorj, 1964), Mediaş (Târnava Mare bank), Curciu (village),	Eua	Xt	
157	<i>Chlorissa viridata</i> (Linnaeus, 1758)	Curciu (Cărbări Hill),	Eua	T	
158	<i>Chlorissa cloraria</i> (Hübner, 1813)	Mediaş (Târnava Mare bank), Curciu (Cărbări Hill),	Eua	Mt	NT
159	<i>Thalera fimbrialis</i> (Scopoli, 1763)	Bazna (Popescu-Gorj, 1964), Mediaş (Târnava Mare bank), Curciu (Cărbări Hill),	Eua	Mt, Xt	
160	<i>Hemistola chrysoprasaria</i> (Esper, 1795)	Curciu (village),	Eua	Mt	
Subfamily Sterrhinae					
161	<i>Cyclophora annularia</i> (Fabricius, 1775)	Curciu (village),	Eua	Mt	NT
162	<i>Cyclophora linearia</i> (Hübner, 1799)	Curciu (village),	Eua	M	
163	<i>Timandra comae</i> (Schmidt, 1931)	Mediaş (Târnava Mare bank), Curciu (village, Cărbări Hill),	Eua	Mt	
164	<i>Scopula immorata</i> (Linnaeus, 1758)	Curciu (Cărbări Hill), Mediaş (Târnava Mare bank, Binderbubi),	Eua	Mt, Mht	
165	<i>Scopula ornata</i> (Scopoli, 1763)	Mediaş (Târnava Mare bank, Binderbubi), Curciu (village, Cărbări Hill),	Eua	Mt	
166	<i>Scopula rubiginata</i> (Hufnagel, 1767)	Mediaş (Greweln, Târnava Mare bank), Curciu (village),	Eua	Xt	

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
167	<i>Scopula immutata</i> (Linnaeus, 1758)	Bazna (Popescu-Gorj, 1964), Mediaş (Târnava Mare bank), Curciu (Cărați Hill),	Eua	Mh	
168	<i>Scopula nemoraria</i> (Hübner, 1799)	Mediaş (Binderbubi),	Eua	M	
169	<i>Scopula virgulata</i> (Denis & Schiffermüller, 1775)	Curciu (Cărați Hill, Viile Pusti),	Eua	Xt	
170	<i>Idaea serpentata</i> (Hufnagel, 1767)	Mediaş (Târnava Mare bank), Curciu (Cărați Hill),	Eua	Xt	NT
171	<i>Idaea ochrata</i> (Scopoli, 1763)	Curciu (Cărați Hill),	Eua	Mxt, Xt	
172	<i>Idaea aureolaria</i> (Denis & Schiffermüller, 1775)	Mediaş – Moşna (Col. Schneider),	Eua	Mx	
173	<i>Idaea biselata</i> (Hufnagel, 1767)	Mediaş (Binderbubi),	Eua	Mht	
174	<i>Idaea humiliata</i> (Hufnagel, 1767)	Mediaş (Binderbubi),	Eua	Mxt	NT
175	<i>Idaea politaria</i> (Hübner, 1799)	Mediaş (Binderbubi),	Vam	Xt	NT
176	<i>Idaea rusticata</i> (Denis & Schiffermüller, 1775)	Curciu (Cărați Hill),	Eua	Mt	NT
177	<i>Idaea aversata</i> (Linnaeus, 1758)	Bazna (Col. Schneider), Mediaş, Curciu (Cărați Hill),	Eua	M	
178	<i>Idaea straminata</i> (Borkhausen, 1794)	Curciu (Cărați Hill),	Eua	Mt	
179	<i>Idaea seriata</i> (Schrank, 1802)	Bazna (Popescu-Gorj, 1964)	Vam	Xt	NT
180	<i>Idaea muricata</i> (Hufnagel, 1767)	Curciu (Cărați Hill),	Eua	Mht	NT
181	<i>Rhodostrophia vibicaria</i> (Clerck, 1759)	Curciu (village), Mediaş (Binderbubi, Târnava Mare bank),	Eua	Xt	
182	<i>Lythria purpuraria</i> (Linnaeus, 1758)	Curciu (Viile Pusti),	Eua	Mth, Mt	
183	<i>Scotopteryx mucronata</i> (Scopoli, 1763)	Curciu (Viile Pusti),	Eua	Mxt	
Subfamily Larentiinae					
184	<i>Xanthorhoe ferrugata</i> (Clerck, 1759)	Bazna (Popescu-Gorj, 1964), Curciu (Cărați Hill), Mediaş (Târnava Mare bank, Binderbubi),	Eua	M, Mh	
185	<i>Xanthorhoe quadrifasiata</i> (Clerck, 1759)	Bazna (Popescu-Gorj, 1964),	Eua	M	
186	<i>Xanthorhoe fluctuata</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	
187	<i>Epirrhoe alternata</i> (Müller, 1764)	Mediaş (Târnava Mare bank),	Eua	Mht	
188	<i>Camptogramma bilineata</i> (Linnaeus, 1758)	Curciu (village, Cărați Hill), Mediaş (Izvorul Dorului),	Eua	M, Mh	
189	<i>Pelurga comitata</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	NT
190	<i>Thera variata</i> (Denis & Schiffermüller, 1775)	Bazna (Popescu-Gorj, 1964),	Eua	M	
191	<i>Colostygia pectinataria</i> (Knoch, 1781)	Curciu (village, Cărați Hill),	Eua	M	

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
192	<i>Horisme vitalbata</i> (Denis & Schiffermüller, 1775)	Curciu (Cărați Hill),	Eua	M	
193	<i>Melanthia procellata</i> (Denis & Schiffermüller, 1775)	Bazna (Popescu-Gorj, 1964), Curciu (village, Cărați Hill), Mediaş (Târnava Mare bank),	Eua	M	
194	<i>Perizoma flavofasciata</i> (Thunberg, 1792)	Bazna (Burnaz, 1993),			NT
195	<i>Perizoma alchemillata</i> (Linnaeus, 1758)	Curciu (village),	Eua	Mx	
196	<i>Eupithecia centaureata</i> (Denis & Schiffermüller, 1775)	Bazna (Popescu-Gorj, 1964),	Eua	M	
197	<i>Minoa murinata</i> (Scopoli, 1763)	Curciu (village),	Eua	M	
Subfamily Ennominae					
198	<i>Abraxas grossulariata</i> (Linnaeus, 1758)	Mediaş (Izvorul Dorului), Curciu (village),	Eua	M	NT
199	<i>Lomaspilis marginata</i> (Linnaeus, 1758)	Mediaş (Târnava Mare bank), Curciu (village, Izvorul Dorului),	Eua	M	
200	<i>Ligdia adustata</i> (Denis & Schiffermüller, 1775)	Bazna (Popescu-Gorj, 1964), Curciu (village), Mediaş (Binderbubi),	Eua	M	
201	<i>Macaria alternata</i> ([Denis & Schiffermüller], 1775)	Curciu, Mediaş (Târnava Mare bank),	Eua	M	
202	<i>Chiasmia clathrata</i> (Linnaeus, 1758)	Mediaş (Târnava Mare bank, Binderbubi), Curciu (village, Cărați Hill),	Eua	M	
203	<i>Tephрина arenacearia</i> ([Denis & Schiffermüller], 1775)	Curciu (village), Mediaş (Târnava Mare bank),	Eua	Xt	NT
204	<i>Heliomata glarearia</i> (Denis & Schiffermüller, 1775)	Curciu (Cărați Hill), Mediaş (Binderbubi),	Vam	Xt	
205	<i>Plagodis pulveraria</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	
206	<i>Plagodis dolabraria</i> (Linnaeus, 1758)	Mediaş (Buzd forest edge),	Eua	M	
207	<i>Therapis flavicaria</i> ([Denis & Schiffermüller], 1775)	Curciu (Cărați Hill),	Vam	Mt	NT
208	<i>Pseudopanthera macularia</i> (Linnaeus, 1758)	Mediaş (Izvorul Dorului), Curciu (Cărați Hill),	Eua	M	
209	<i>Ennomos erosaria</i> (Denis & Schiffermüller, 1775)	Curciu (village),	Eua	M	NT
210	<i>Selenia lunularia</i> (Hübner, 1788)	Curciu (village),	Eua	M	NT
211	<i>Apeira syringaria</i> (Linnaeus, 1758)	Curciu (village),	Eua	Mxt	NT
212	<i>Colotois pennaria</i> (Linnaeus, 1716)	Curciu (village),	Eua	M	
213	<i>Angerona prunaria</i> (Linnaeus, 1758)	Curciu, Mediaş (Binderbubi)	Eua	M	
214	<i>Biston betularia</i> (Linnaeus, 1758)	Curciu (village), Mediaş (Izvorul Dorului),	Eua	M	

No.	TAXON	LOC.	Z. El.	E. C.	R. L.
215	<i>Agriopis marginaria</i> (Fabricius, 1776)	Mediaş (Greweln, Binderbubi)	Eua	M	
216	<i>Hypomecis punctinalis</i> (Scopoli, 1763)	Curciu (village), Mediaş (Binderbubi)	Eua	M	
217	<i>Ascotis selenaria</i> (Denis & Schiffermüller, 1775)	Curciu (village, Cărați Hill), Mediaş (Târnava Mare bank, Binderbubi),	Eua	M	
218	<i>Ectropis crepuscularia</i> (Denis & Schiffermüller, 1775)	Moşna (Col. Schneider), Curciu, Mediaş (Binderbubi)	Eua	M	
219	<i>Ematurga atomaria</i> (Linnaeus, 1758)	Mediaş (Izvorul Dorului, Târnava Mare bank), Curciu (village, Cărați Hill),	Eua	M	
220	<i>Cabera pusaria</i> (Linnaeus, 1758)	Curciu (village),	Eua	M	
221	<i>Campaea margaritata</i> (Linnaeus, 1767)	Mediaş (Izvorul Dorului), Curciu (village, Cărați Hill)	Eua	M	
222	<i>Siona lineata</i> (Scopoli, 1763)	Mediaş (Izvorul Dorului, Târnava Mare bank), Curciu (village, Cărați Hill),	Eua	M	

**DATA REGARDING GENUS *PARNASSIUS* LATREILLE, 1804
(LEPIDOPTERA: PAPILIONIDAE) PRESERVED IN
NATURAL HISTORY MUSEUM COLLECTIONS FROM SIBIU**

Sergiu-Cornel TÖRÖK*
Gabriela CUZEPAN**

Abstract. The aim of this study is to present the specimens belonging to the genus *Parnassius* Latreille, 1804 preserved in the collections of the Natural History Museum from Sibiu. The material analyzed counts 244 specimens belonging to the following species: *Parnassius apollo* (Linnaeus, 1758) with 2 subspecies: *Parnassius apollo jaraensis* Kertész, 1922 and *Parnassius apollo transsylvanicus* Schweitzer, 1912; and *Parnassius mnemosyne* (Linnaeus, 1758) with 2 subspecies: *Parnassius mnemosyne distincta* Bryk & Eisner, 1930 and *Parnassius mnemosyne transsylvanica* Schmidt, 1930. The four subspecies are endemic to Romania. Their distribution in Romania is given according to the collecting sites from the museum collections. The conservation status in Romania it is also presented for both.

Keywords: *Parnassius* sp., distribution, faunistics, museum collections.

Rezumat. Acest studiu prezintă genul *Parnassius* Latreille, 1804 din colecția Muzeului de Istorie Naturală din Sibiu. Materialul analizat cuprinde 244 de specimene încadrate sistematic în 2 specii: *Parnassius apollo* (Linnaeus, 1758) cu 2 subspecii: *Parnassius apollo jaraensis* Kertész, 1922; *Parnassius apollo transsylvanicus* Schweitzer, 1912 și *Parnassius mnemosyne* (Linnaeus, 1758) cu 2 subspecii: *Parnassius mnemosyne distincta* Bryk & Eisner, 1930; *Parnassius mnemosyne transsylvanica* Schmidt, 1930. Cele 4 subspecii sunt endemice pentru România. Locurile de colectare împreună cu statutul din cadrul Listei Roșii a Fluturilor din România sunt de asemenea prezentate.

Cuvinte cheie: *Parnassius* sp., răspândire geografică, faunistică, colecții muzeale.

Introduction

The first article on Lepidoptera published in Romania by Carl Fuss, was based on the material collected by Joseph von Franzenau from Săcărâmb (Hunedoara County, Romania). This paper appeared in the first edition of the Transylvanian Society of Natural Sciences Journal (*Verhandlungen und Mitteilungen des Siebenbürgische Vereins für Naturwissenschaften zu Hermannstadt*) from Sibiu (Popescu-Gorj 1970). The material included a list of 964 species and forms of Macro- and Microlepidoptera. The same paper confirms for the first time the presence in Romania of the species *Doritis* (*Parnassius*) *apollo* (Linnaeus, 1758) and *Doritis mnemosyne* Linnaeus, 1758 (Fuss, 1850).

Doritis is also mentioned by Josef von Franzenau in his work published in the third number of the

Transylvanian Society Journal (1859), the specimens being collected from Kisbánya (today's Băișoara village, Cluj County, Romania) (Franzenau 1852, 1859).

During 1860 – 1866 Josef Mann from Vienna studied the fauna from the northern part of Dobrudja (at that time it was included in the Ottoman Empire) and he published several species, which remain the only citing in the Romania fauna (Mann 1866, Székely 2008).

Noticeable is the citing of *Doritis* (*Parnassius*) *apollo* (Linnaeus, 1758) in this region, considered uncertain, being the only mention of the species in the northern part of Dobrudja until today.

Regarding the distribution of *P. apollo* and *P. mnemosyne* in Romania, Constantin Hormuzachi and Daniel Czekelius wrote about these species in their articles published in the beginnings of the 20th century.

These two species are also mentioned in the first Catalogue of Lepidoptera from Romania, published by Aristide Caradja in 1895 (Caradja 1895), followed by Edmund Fleck's catalogue in

* Babeș-Bolyai University of Cluj-Napoca,
ser.torok@yahoo.com

**Brukenthal National Museum, Natural History Museum,
gabrielacuzepan@gmail.com,
gabriela.cuzepan@brukenthalmuseum.ro

1904 (Fleck, 1904), and by Franz Salay's catalogue in 1910 (Salay 1910).

At the beginning of the 20th century the most important contributions to the distribution of *P. apollo* (Linnaeus, 1758) and *P. mnemosyne* (Linnaeus, 1758) were made by Daniel Czekelius in his catalogues of the Transylvanian Lepidoptera (Czekelius 1897, Czekelius, 1917). In the Transylvanian Society Journal from 1920, Daniel Czekelius dedicates an entire article to *Parnassius apollo* (Linnaeus, 1758) introducing new data concerning the subspecies and variations from Transylvania and their distribution (Czekelius 1922).

The Transylvanian Society Journal included a series of papers related to the distribution of *P. mnemosyne* and *P. apollo*: in number 62 (1908) Baron N. Charles Rothschild (London) identified *P. mnemosyne* at Mezőség (today's Transylvanian Plain, Romania) and in number 83-84, László Diószeghy made remarks about the presence of *P. apollo* in the Retezat Mountains (Rotschild 1912, Diószeghy 1933-1934).

In the post war period there were relatively few publications regarding these two species: the *Catalogue of Adriano Ostrogovich* written by Aurelian Popescu-Gorj in the year 1964 (Popescu-Gorj 1964), *Family Papilionidae* by Eugen Niculescu published in Fauna R. S. Romania (Niculescu 1961), in the *Catalogue of the Lepidoptera collection from The Banat Museum (Timișoara)* by Frederic König (König 1975) and in the *Catalogue of Nicolae Delvig from Brașov* by Victor Ciochia and A. Barbu (Ciochia, Barbu 1980).

In the last decades, The Red List of the Romanian butterflies shows the presence of two geographical distinct subspecies of *Parnassius apollo*, ssp. *transsylvanicus* and ssp. *jaraensis* (Rákossy 2003). Other studies suggest that in reality in Romania only one subspecies of *Parnassius* (*Parnassius a. transsylvanicus*) is present (Niculescu 1961, Ruști, Dragomirescu 1991).

P. a. jaraensis seem to have been extinct from the Western Carpathians, while populations of *P. a. transsylvanicus* have also disappeared (Mihuț, Dincă 2006).

The decline was much accentuated during the last century, when the species seem to have disappeared from dozen of places where it had been recorded during the 19th century. Over-collecting might have been a significant role in the disappearance of the unique population of *P. a.*

jaraensis from Răcățău (Western Carpathians), it is more than 20 years since this subspecies was cited (Mihuț, Dincă 2006).

Victor Weindel presents new information regarding the distribution of *P. apollo* in Transylvania, and elaborates two hypotheses about the dispersion of *P. apollo* in Romania (Weindel 2000).

This study is intended to present data regarding *Parnassius* sp. preserved in the Lepidoptera Collection of the Natural History Museum from Sibiu.

Material and methods

The study is based on the Lepidoptera Collection of Natural History Museum from Sibiu. The investigated material was from: Daniel Czekelius Lepidoptera Collection from Transylvania, Victor Weindel Lepidoptera Collection from Transylvania, Eugen Worell Entomological Collection, Heinrich Hann von Hannenheim Entomological Collection, Rolf Weyrauch Entomological Collection and Eckbert Schneider Entomological Collection.

From these 6 collections the oldest is Daniel Czekelius Lepidoptera Collection, the collection includes specimens from 1887 (Pascu, Schneider 1998), he began sampling from 1887. This collection is also the largest Lepidoptera Collection of the Natural History Museum from Sibiu, containing 7.162 specimens of Macrolepidoptera and Microlepidoptera from Transylvania, as well as 6.929 other Palearctic Lepidoptera specimens.

Daniel Czekelius received material from Emil von Silbernagel from Avrig, Sachsenheim and Karl Petri from Sighișoara, Friedrich Deubel from Brașov, Köntzei Geró, Ștefan Peterfi, Eugen Worell, Arnold Müller, Josef v. Mallász, Adriano Ostrogovich, Béla Kiss, Rolf Weyrauch and many others. In one of his field trips to Retezat Mountains he met the famous lepidopterologist László Diószeghy, with whom he started collaborating. Other important collaboration, during his work as an entomologist, was the one with Hans Rebel from the Museum of Vienna, who showed interest in the fauna from Transylvania and who took part in several trips in this region (Pascu, Schneider 1998).

Daniel Czekelius last papers, published between 1922 and 1943, brought new data on the distribution of the genus *Parnassius* (syn=*Apollo*), a genus in which he was very interested studying the species in detail.

Heinrich Hann von Hanneenheim Entomological Collection contains over 1.902 lepidoptera specimens. Heinrich Hann von Hanneenheim collected during 1920-1964, especially from Transylvania region, particularly from the following sites: Păltiniș, Bărcăciu, Poiana Neamțului, Avrig Lake, Podragu and Sâmbăta Valley (Schneider 2003).

Eugen Worell Lepidoptera Collection, part of the Eugen Worell general Entomological Collection (93.897 specimens), includes 10.923 specimens of Macrolepidoptera and Microlepidoptera. The foundations of this collection were set in 1920; the items were collected from Romania, and from other different areas such as Europe, Africa, Asia, North and South America. He collected materials from all over the country, and in Transylvania he collected specimens from Sibiu and its surroundings.

Victor Weindel Transylvanian Lepidoptera Collection contains 4.322 specimens of Macrolepidoptera and Microlepidoptera, which were collected by the author between 1900-1959, particularly from Southern Transylvania (Sibiu and its surroundings), but also from other geographical regions from Romania (Schneider 1984).

Rolf Weyrauch Lepidoptera Collection contains over 5.500 Lepidoptera specimens out of 16.436 items from his entire entomological collection, the Noctuidae and the Geometridae families being best represented. The majority of these items were collected from around Sibiu (Pascu, Schneider 1998).

The most recent collection preserved in the Natural History Museum of Sibiu is Eckbert Schneider Entomological Collection. His collection contains over 20.000 specimens. In 30 years of activity, he collected specimens from the whole country, especially from the southern part of Transylvania.

The material from these 6 collections is presented in the following pages. Each species is followed by the number of specimens, the collecting site, date of collecting and the abbreviation of the collector. For an easy identification of the collecting sites, the original citing is mentioned followed by the present name and is included in the specific region of Romania or county. The species synonyms are mentioned in square brackets according with the labels from collections.

Abbreviations used in this paper:

alt. = altitude; ANONYMOUS = anonymous collector; n. – number of the specimen according to Daniel Czekelius Lepidoptera Collection from Transylvania; spec. = specimen/s; leg. = collected by; det. = determined in; Mt. / Mts. = mountain/s; / = or.

Collectors' name:

Béla Kiss = B.K.; Csab Béla = C.B.; Diószeghy László = D.L.; Adriano. Ostrogovich = A.O.; Arnold Müller = A.M.; Daniel Czekelius = D.C.; Eugen Worell = E.W.; Köntzei Geró = K.G.; Ștefan Peterfi = S.P.; Eckbert Schneider = E.S.; Emil v. Silbernagel = E.Silb.; H. Pelits = H.P.; Heinrich Hann von Hanneenheim = H.H.; Josef v. Mallász = J.M.; Kuales = K.; Ladislaus v. Dobay = L.D.; N. Falmtius = N.F.; Rolf Weyrauch = R.W.

Results and discussions

Taxonomic Hierarchy (Rákósy 2003; Rákósy *et al.* 2003):

Genus *Parnassius* Latreille, 1804

(*Doritis* Fabricius, 1807)

Species *apollo* Linnaeus, 1758

Subspecies *jaraensis* Kertész, 1922

Subspecies *transsylvanicus* Schweitzer, 1912

syn. *rosenius* Fruhstorfer, 1923

Species *mnemosyne* Linnaeus, 1758

Subspecies *wagneri* Bryk, 1925

Subspecies *transsylvanica* Schmidt, 1930

syn. *dioszeghyi* Bryk, 1930

Subspecies *distincta* Bryk & Eisner, 1930

syn. *hungaricus* Rothschild, 1911

syn. *bischoffi* Bryk & Eisner, 1930

syn. *cibinensis* Dannehl, 1933

syn. *czekeli* Bryk & Eisner, 1934

syn. *distinctus* Bryk & Eisner, 1935

Daniel Czekelius Lepidoptera Collection from Transylvanian

Parnassius apollo (Linnaeus, 1758)

1 spec. (n.37) Ratosnya [Răstolița, Mureș County], leg. ANONYMOUS; 1 spec. (n.39/28) Borgo Bistritza [Bistrița Bărgăului, Bistrița Năsăud County], leg. ANONYMOUS; 1 spec. (n.40) Békás / Békás patak [Bicaz, Neamț County], 16.07.1937, leg. ANONYMOUS; 3 specs. (n.41/9, n.42/10, n.43/11) Tölgyes – Putanapatak [Tulgheș, Harghita County], 23.07.1909 (?), leg. ANONYMOUS; 1 spec. (n.44/4) Gyewgyo st. Miklos [Gheorghieni, Harghita County], 28.07.1910, leg. ANONYMOUS; 5 specs. (n.45, n.46/5, n.47/8, n.48/7, n.49/6) Gyewgyo st. Miklos

[Gheorghieni, Harghita County], 24.07.1909, leg. ANONYMOUS; 1 spec. (n.72/97) Broos [Orăștie, Hunedoara County], leg. ANONYMOUS; 2 specs. (n.51, n.55) Bórnék [Borsec, Harghita County], 18.07.1911, leg. E.W.; 3 specs. (n.52, n.53, n.54) Bórnék [Borsec, Harghita County], 18.07.1911, leg. ANONYMOUS; 1 spec. (n.56) Bórnék [Borsec, Harghita County], 1.08.1921, leg. ANONYMOUS; 1 spec. (n.57) Bórnék [Borsec, Harghita County], 15.07.1922, leg. ANONYMOUS; 1 spec. (n.58/22) Bórnék [Borsec, Harghita County], 18.07.1911, leg. ANONYMOUS.

Parnassius apollo jaraensis Kertész, 1922

2 specs. (n.59, n.60) leg. Mărișel [Mărișelul, Gilăului Mts.], 03.07.1927, leg. S.P.; 1 spec. (n.61) without location, 24.07.1929, leg. ANONYMOUS; 3 specs. (n.62, n.63, n.64) Réketo [Răcățau, Cluj County], 14.07.1929, leg. ANONYMOUS; 1 spec. (n.50/47) Borgo-Bistritza [Bistrița Bârgăului, Bistrița Năsăud County], 12.07., leg. ANONYMOUS; 3 specs. (n.65, n.66, n.67) Colibitza [Colibița Lake, Bistrița Năsăud County], 14.07.1930, leg. A.M.; 1 spec. (n.68) Colibitza [Colibița Lake, Bistrița Năsăud County], 14.07.1930, leg. D.C.; 3 specs. (n.69, n.70/45, n.71/44) Retezat (Netiș?) [Retezat Mts.], 29.08.1930, leg. C.B.; 1 spec. (n.314) - has only the right pair of wings stuck on a label, Retezat (Nyekistok?) [Retezat Mts.], 15.08.1930, leg. A.M.

Parnassius mnemosyne distincta Bryk & Eisner, 1930

[*Parnassius mnemosyne bischoffi* Bryk & Eisner, 1930]

1 spec. (n.115) Kuhhorn [Ineu Peak, Rodna Mts.] 03.06.1911, leg. ANONYMOUS.

[*Parnassius mnemosyne czekelii* (?) Bryk & Eisner, 1934]

1 spec. (n.74) Riu Sadu [Râu Sadu, Sibiu County], 06.1935, leg. D.C.; 1 spec. (n.75) Riu Sadu [Râu Sadu, Sibiu County], 06.1936, leg. A.O.; 1 spec. (n.76) Riu Sadu [Râu Sadu, Sibiu County], ??07.1935, leg. A.O.; 1 spec. (n.77) Prejba [Prejba, Lotru Mts.], 27.07.1910, leg. H.H.; 2 specs. (n.78, n.79) Küküllövar [Cetatea de Baltă, Alba County], 09.05.1921, leg. K.G.; 1 spec. (n.80) Küküllövar [Cetatea de Baltă, Alba County], 19.05.1932 leg. K.G.; 2 specs. (n.81, n.82) Kukullburg [Kokelburg – Cetatea de Baltă, Alba County], 29.05.1932(?), leg. D.C.; 1 spec. (n.83) Kukullburg [Kokelburg – Cetatea de Baltă, Alba County], 29.05.1932(?), leg. K.G.; 1 spec. (n.84) Kukullburg [Kokelburg - Cetatea de Baltă, Alba County], 20.05.1921, leg. ANONYMOUS; 2 specs.

(n.85, n.86) Schässburg [Sighișoara, Mureș County], 30.06., leg. D.C.; 1 spec. (n.87) Hadad [Hodod, Satu Mare County], 29.08.1932, leg. B.K.; 1 spec. (n.88) Kolosvar, Klausenburg [Cluj Napoca, Cluj County], 18.04.1919, leg. ANONYMOUS; 2 specs. (n.89, n.90) Nagyág [Săcărâmb, Hunedoara County], 2.06.1928, leg. J.M.; 2 specs. (n.91, n.92) Deva(?) [Deva, Hunedoara County], 1.06.1924, leg. J.M.; 2 specs. (n.93, n.94) Tușnad [Harghita County], 21.04., det. 1932, leg. L.D.; 1 spec. (n.95) Kronstadt [Brașov, Brașov County], 04.1908, det. 15.05.1932, leg. E.Silb.; 1 spec. (n.96) Kronstadt [Brașov, Brașov County], 04.1908, det. 16.04.1932, leg. E.Silb.; 2 specs. (n.97, n.98) without data, leg. E.Silb.; 1 spec. (n.99) Hermannstadt, Michelsberg [Sibiu, Cisnădioara, Sibiu County], 24.05.1916, leg. D.C.

[*Parnassius mnemosyne distinctus* Bryk & Eisner, 1930]

1 spec. (n.100) Transilvania Butschetch [Transylvania, Bucegi Mts.], without data, leg. D.L.; 1 spec. (n.101) Honanstein, Kronstadt [Brașov, Brașov County], without data, leg. ANONYMOUS; 1 spec. (n.102) Schuler [Postăvarul Mts.], 14.06.1908, leg. D.C.; 2 specs. (n.103, n.104) Fedeleesch [Chica Fedeleșului Peak, Făgăraș Mts.], 31.05.1909, leg. D.C.; 3 specs. (n.105, n.106, n.107) Götzenberg [Măgura Cisnădiei, Sibiu County], 05.1932, leg. A.M.; 1 spec. (n.108) Riu Sadu [Râu Sadu, Sibiu County], 06.1932, leg. A.O.; 2 specs. (n.109, n.110) Riu de Mori, Malomvíz [Râu de Mori, Hunedoara County], 30.06.1924, leg. J. M.; 1 spec. (n.111) Surul Berg [Suru Mts., Făgăraș Mts.], 11.06.1922, leg. H.H.; 1 spec. (n.112) Zibinsgeb., Valea Ștezii [Cindrel Mts., Valea Șteaza], 09.06.1930, leg. A.M.; 1 spec. (n. 113) Tălmach, Kromwald Lands [Tălmăci, Sibiu County], 26.05., leg. D.C.; 1 spec. (n.114) Kronstadt [Brașov, Brașov County], without data, leg. E.Silb.

[*Parnassius mnemosyne hungaricus* Rothschild, 1911]

1 spec. (n.73) Nagy-Kagya, Bihar [Cadea Mare, Bihor County], 04.06.1913, leg. ANONYMOUS.

Parnassius mnemosyne transsylvanica Schmidt, 1930

1 spec. (n.116) Retezat [Retezat Mts.], 03.06.1923, leg. D.L.; 3 specs. (n.117, n.118, n.119) Retezat, Campanula Schultz [Retezat Mts., Campanula Shelter], 06.1929, leg. J.M.; 2 specs. (n.120, n.121) Riu Sadu [Râu Sadu, Sibiu County], ??06.1932, leg. A.O.

Victor Weindel Transylvanian Lepidoptera Collection

Parnassius apollo transsylvanicus Schweitzer,
1912

2 specs. Bârgău-Bistrița [Bistrița Năsăud County], 12.07., leg. D.C.; 1 spec. Borsec [Harghita County], 08.1914, leg. A.M.; 1 spec. Borsec [Harghita County], 29.07.1921, leg. ANONYMOUS; 6 specs. Borsec [Harghita County], 1.08.1921, leg. ANONYMOUS; 2 specs. Borsec [Harghita County], 2.08.1921, leg. ANONYMOUS; 1 spec. without data, leg. ANONYMOUS.

Parnassius mnemosyne distincta Bryk & Eisner,
1930

[*Parnassius mnemosyne distinctus* Bryk & Eisner,
1930]

5 specs. Măgura Cisnădiei [Sibiu County], 22.05.1904, leg. ANONYMOUS; 3 specs. Măgura Cisnădiei [Sibiu County], 22.05.1921, leg. ANONYMOUS; 1 spec. Brașov, Tâmpa [Brașov County], 15.05.1921, leg. ANONYMOUS; 1 spec. Ocna Sibiului [Sibiu County], 04.06.1924, leg. ANONYMOUS; 5 specs. Postăvaru [Postăvaru Mts], leg. ANONYMOUS; 5 specs. Măgura Cisnădiei – Schweinswiese [Sibiu County], 13.06.1954, leg. ANONYMOUS.

Eugen Worell Entomological Collection

Parnassius apollo (Linnaeus, 1758)

2 specs. Bucovina [Bucovina Region], leg. ANONYMOUS; 1 spec. Retezat(?), Netiș [Retezat Mts.], 29.08.1930(?), leg. C.B.(?); 1 spec. without location and data, leg. ANONYMOUS; 1 spec. Hășmașu Mare [Hășmașu Mare Mts.], 07.1936, leg. K.; 2 specs. Mții. Bârgău [Bârgău Mts.], 07.1036, leg. K.

Parnassius apollo jaraensis Kertész, 1922

4 specs. Reketo bei Cluj [Răcățiu, Cluj County], 14.07.1929, leg. D.C.; 1 spec. Borsec [Harghita County], 4.08.1911, leg. D.C.

Parnassius apollo transsylvanicus Schweitzer,
1912

1 spec. Borsec [Harghita County], 15.07.1922, leg. ANONYMOUS; 2 specs. without location and data, leg. ANONYMOUS.

Parnassius mnemosyne transsylvanica Schmidt,
1930

3 specs. Götzenberg [Măgura Cisnădiei, Sibiu County], 14.05.1947 leg. E.W.; 2 specs. Götzenberg [Măgura Cisnădiei, Sibiu

County], ??.06.1930, leg. E.W.; 1 spec. Götzenberg [Măgura Cisnădiei, Sibiu County], 06.09.1938, leg. E.W.; 2 specs. Götzenberg [Măgura Cisnădiei, Sibiu County], 17.06.1948, leg. E.W.

Heinrich Hann von Hanneheim Entomological Collection

Parnassius apollo transsylvanicus Schweitzer,
1912

1 spec. Rarău, M. Aramei [Rarău Mts., Aramei Mt.], 26.07.1959, leg. ANONYMOUS.

Parnassius mnemosyne distincta Bryk & Eisner,
1930

[*Parnassius mnemosyne distinctus* Bryk & Eisner,
1930]

1 spec. Götzenberg bei Hermannstadt, 1300m [Măgura Cisnădiei, Sibiu County], 14.05.1947, leg. E.W.; 1 spec. Podragu [Podragu Mt., Făgăraș Mts.], 30.06.1954, leg. ANONYMOUS; 1 spec. Podragu [Podragu Mt., Făgăraș Mts.], 10.07.1954, leg. ANONYMOUS; 1 spec. Hammersdorf [Gușterița, Sibiu County], 31.05.1964, leg. ANONYMOUS; 1 spec. Mănăstirea Sâmbăta [Sâmbăta Monastery, Brașov County], 08.07.1964, leg. ANONYMOUS.

Rolf Weyrauch Entomological Collection

Parnassius apollo jaraensis Kertész, 1922

7 specs. Valea Someș Rece [Cluj County], 1974, Răcățiu [Cluj County], 25-28.07, leg. R.W.; 2 specs. Valea Someș Rece [Cluj County], 18.07.1971, leg. H.P.; 2 specs., Valea Someș Rece [Cluj County], 20-22.07.1976, leg. R.W.

Parnassius apollo transsylvanicus Schweitzer,
1912

1 spec. Borsec [Harghita County], 25.07.1921, leg. A.M.; 1 spec. Borsec [Harghita County], 1.08.1921, leg. ANONYMOUS; 1 spec. Borsec [Harghita County], 2.08.1921, leg. ANONYMOUS.

[*Parnassius apollo rosenius* Fruhstorfer, 1923]

1 spec. Zugreni [Bistrița Năsăud County], 04.08.1974, leg. ANONYMOUS.

Parnassius mnemosyne (Linnaeus, 1758)

1 spec. without data, 05.05.1949, leg. R.W.; 6 specs. Hammersdorfer Berg (H.B.) [Dealul Gușteriței, Sibiu County], 05.06.1965, leg. R.W.; 1 spec. without data, 05.1950, leg. R.W.; 1 spec. Hammersdorfer Berg (H.B.) [Dealul Gușteriței, Sibiu County], 04.06.1967, leg. R.W.; 7 specs. Hammersdorfer Berg (H.B.) [Dealul Gușteriței,

Sibiu County], 03.06.1964, leg. R.W.; 4 specs. Surul [Suru Mt., Făgăraș Mts.], 12.06.1966, leg. R.W.; 4 specs. Vânturarița [Buila-Vânturarița, Mts., Căpățanii Mts.], 16/18.07.1960, leg. R.W.; 1 spec. Cozia [Cozia Mts.], 20.05.1966, leg. R.W.; 1 spec. Cozia [Cozia Mts.], 13.05.1967, leg. R.W.; 1 spec. Valea Sâmbăta [Brașov County], 12.5/6.1963 leg. R.W.; 2 specs. Tușnad [Harghita County], 22.07.1954, leg. R.W.; 1 spec. Tușnad [Harghita County], 01.07.1954, leg. R.W.; 2 specs. Domogled [Mehedinți Mts.], 26.06.1964, leg. R.W.; 7 specs. Domogled [Mehedinți Mts.], 30.06.1956, leg. R.W.; 5 specs. Domogled [Mehedinți Mts.], 09.06.1970, leg. R.W.; 4 specs. Dobrogea, Canaraua Fetii [Dobrogea de Sud Plateau], 15-18.05.1975, leg. R.W..

Eckbert Schneider Entomological Collection

Parnassius apollo (Linnaeus, 1758)

2 specs. without location, 30.VI.1962, leg. E.S.

Parnassius mnemosyne (Linnaeus, 1758)

6 specs. Hoia, Cluj [Cluj County], 06.1956, leg. E.S.; 1 spec. Valea Vinului, Rodna [Rodna Mts.], 29.06.1955 leg. E.S.; 1 spec. Hammersdorf [Gușterița, Sibiu County], 09.06.1954, leg. E.S.; 1 spec. Fântâna Rece, Gușterița, Sibiu [Sibiu County], 16.05.1973, leg. E.S.; 1 spec. Mții. Făgărașului, Poiana Neamțului [Făgăraș Mts.], 02.06.1978, leg. E.S.; 3 specs. Șeica Mare [Sibiu County], 22.05.1978, leg. E.S.; 1 spec. Saschiz [Mureș county], 29.05.1982, leg. E.S.; 1 spec. Mții. Făgărașului, Poiana Neamțului [Făgăraș Mts.], 17.05.1981 leg. E.S.; 6 specs. Mții. Făgărașului, Poiana Neamțului [Făgăraș Mts.], 29.05.1981, leg. E.S.; 1 spec. Rivasco, 20.06.1963, leg. E.S.; 2 specs. Retezat 1200m [Retezat Mts., 1200 alt.], 23.06.1957, leg. E.S.; 1 spec. Domogled [Mehedinți Mts.], 29.07.1957, leg. N.F.; 1 spec. Butschetsch [Bucegi Mts.], 06.1949, leg. ANONYMOUS; 1 spec. Hammersdorf Berg [Dealul Gușteriței, Sibiu County], 09.06.1954, leg. E.S.; 1 spec. Mții. Făgărașului, Poiana Neamțului [Făgăraș Mts.], 24.06.1964, leg. E.S.; 1 spec. Sibiel [Sibiu County], 30.05.1982, leg. E.S.

The material analyzed consists of 244 specimens belonging to the following species: *Parnassius apollo* (Linnaeus, 1758) with 2 subspecies: *Parnassius apollo jaraensis* Kertész, 1922 and *Parnassius apollo transsylvanicus* Schweitzer, 1912; and *Parnassius mnemosyne* (Linnaeus, 1758) with 2 subspecies: *Parnassius mnemosyne distincta* Bryk & Eisner, 1930 and *Parnassius mnemosyne transsylvanica* Schmidt, 1930, of the Natural History Museum from Sibiu.

Parnassius apollo is represented in the Lepidoptera Collection by 67 specimens, from which 16 specimens are represented by the subspecies *Parnassius apollo jaraensis* and 21 specimens by the subspecies *Parnassius apollo transsylvanicus* (Fig. 1). The species *Parnassius mnemosyne* counts 177 specimens, having the highest percentage in the investigated Lepidoptera collections, followed by the subspecies *Parnassius mnemosyne distincta* with 84 specimens and the subspecies *Parnassius mnemosyne transsylvanica* with 16 specimens (Fig. 1).

P. apollo is present in the following collections: Daniel Czekelius Lepidoptera Collection from Transylvania, Eugen Worell Lepidoptera Collection and Eckbert Schneider's Entomological Collection. *P. a. jaraensis* appear only in the following collections: Daniel Czekelius Lepidoptera Collection from Transylvania, Eugen Worell Lepidoptera Collection, Rolf Weyrauch Lepidoptera Collection, and *P. a. transsylvanicus* appears in: Victor Weindel Lepidoptera Collection from Transylvania, Eugen Worell Lepidoptera Collection, Heinrich Hann von Hanenheim Lepidoptera Collection and Rolf Weyrauch Lepidoptera Collection.

P. mnemosyne is present only in the collections of Rolf Weyrauch Lepidoptera Collection and Eckbert Schneider Entomological Collection. The subspecies of *P. mnemosyne* respectively *P. m. distincta* is found in: Daniel Czekelius Lepidoptera Collection from Transylvania, Victor Weindel Lepidoptera Collection from Transylvania, Rolf Weyrauch Lepidoptera Collection; respectively *P. m. transsylvanica* is found only in: Daniel Czekelius Lepidoptera Collection from Transylvania and Rolf Weyrauch Lepidoptera Collection.

The collecting sites of *P. apollo* are: Răstolița [Mureș County], Bistrița Bârgăului [Bistrița Năsăud County], Bicaz [Neamț County], Borsec; Tulgheș & Gheorghieni [Harghita County], Orăștie [Hunedoara County], Retezat Netiș [Retezat Mts.], Hășmașu Mare Mts., Bârgău Mts. and also from other locations that weren't noted on the labels. The largest numbers of specimens of *P. apollo* where sampled from Borsec region (Harghita County) (Fig. 2, Fig. 3).

The sampling sites of *P. a. jaraensis* are: Bistrița Bârgăului and Colibița Lake [Bistrița Năsăud County], Borsec [Harghita County], Retezat, Netiș [Retezat Mts.], Hășmașu Mare Mts., Răcățau & Someș Rece Valley [Cluj County], his distribution

range is in the north-east and south-west part of Transylvania (Fig. 3).

P. a. transsylvanicus was sampled from Bistrița Bârgăului and Zugreni [Bistrița Năsăud County], Borsec [Harghita County], Rarău Mts., Aramei Mt., showing that his distribution is restricted to the north-east part of Transylvania, especially in the north area of the Oriental Carpathians (Fig. 3).

Parnassius apollo (Linnaeus, 1758) specimens, which appear in the collections, were sampled between 1909 and 1976. More specifically, *P. apollo* was sampled for the first time in the year 1909, and in the years 1910, 1911, 1921, 1922, 1930, 1936, 1937, 1962. In 1962 was the last time when this species has been sampled, being collected by E. Schneider.

P.a. jaraensis was sampled in: 1911, 1927, 1929, 1971, 1974 and 1976 by R. Weyrauch from Someș Rece Valley [Cluj County]. Regarding the subspecies *P. a. transsylvanicus* the following years are mentioned: 1914, 1921, 1922, 1959 and 1974 collected by R. Weyrauch from Zugreni [Bistrița Năsăud County].

There are two high points of sampling, one in 1921 and another in 1930 both with 13 specimens. A great number of specimens were collected in the years 1909, 1910 (7-8 specs.) and 1974 (8 specs.) (Fig. 4).

Dividing the period 1909-1976 in two parts, we concluded that the majority of specimens are collected before 1937, after this year only 16 specimens from the total of 67 specimens were collected. This may possible suggest that the number of specimens in the wild was starting to decline. The decline of this species in Romania is also mentioned and confirmed by different authors (Mihuț, Dincă, 2006); this is why *Parnassius apollo* was included in *The Red Book of Romanian Lepidoptera*, with the status of Critically Endangered (Rákossy 2003).

This butterfly is also included in the Habitats Directive, Annex 3A and Annex 4A (Rákossy 2005).

At national level were established threats that had brought *Parnassius apollo* to his decline, like: climatic change; over collecting (killing or taking); isolation and fragmentation of habitat; deforestation on non-woodland habitats; overgrazing; abandonment and inappropriate habitat management, and others (Van Sway, Warren 1999).

In Romania only one conservation measure is taken that of the legal protection of habitats. Other countries have implemented many measures like: begin or improve monitoring, ecological research on species requirements, reintroduction, restrict recreational activities, avoid overgrazing, avoid natural deforestation, improve habitat management, stop development of mountain areas with sensitive populations and enforcement of measures proscribed by law (Van Sway, Warren 1999).

The specimens of *Parnassius mnemosyne* (Linnaeus, 1758) (Fig. 5), were sampled from Transylvania, especially from Sibiu County but also from Cluj County, Brașov County and Harghita County. A part of the *P. mnemosyne* specimens come from Rodna Mts., Făgăraș Mts., Retezat Mts., Căpățâni Mts., Cozia Mts. and Mehedinți Mts.

From Dobrogea, in the Schneider Entomological Collection, 4 specimens from Canaraua Fetii Canyon sampled in 1975, are present.

The subspecies *P. m. distincta* was sampled from whole Transylvania, but the biggest part of it comes from Sibiu County (Fig. 5).

Concerning the endemic subspecies *Parnassius mnemosyne transsylvanica* the collecting sites were: Retezat Mts., Campanula Shelter also from Retezat Mts., Râu Sadu and Măgura Cîsnădiei from Sibiu County. Therefore we can observe that the distribution range of this endemic subspecies of butterfly seems to be restricted to Transylvania (Fig. 5) (Rákossy 1997).

The analysis of collection years (Fig. 6) reveals that the specimens of *P. mnemosyne* were collected in the period between 1904 and 1982.

The earliest specimens were collected by V. Weindel in 1904 (5 specs.), more then one hundred years ago, after this date the number of specimens collected fluctuates homogeneously until 1982, when the last specimens of *P. mnemosyne* was sampled by E. Schneider.

In the 74 years, representing the collecting period of the *P. mnemosyne* specimens, there are three high points in 1932, 1956 (13 specs.) and 1964 (11 specs).

The numbers of specimens per year are equally distributed suggesting that there is stability in this butterfly population (Fig. 6).

Concerning the conservation status of *P. mnemosyne* it is included in *The Red Book of Romanian Butterflies* with the Status: Near

Threatened (Rákósy 2003). This species is also included in Habitats Directive, Annex 3A and Annex 4A (Rákósy 2005).

The threats in Romania are: destruction of woodland; isolation and fragmentation of habitat; change of woodland management; overgrazing and others (Van Sway, Warren 1999).

Although *P. mnemosyne* lives in isolated populations (Niculescu, 1961), it is a rather common butterfly, for this reason, there are no conservation measures proposed.

Acknowledgements

We would like to kindly thank Dr. Cristina Moise, Dr. Cornelia Chimişliu, Dr. Silvia Burnaz and Ioan Tăușan for the help in reviewing the paper.

REFERENCES

- Caradja 1895 Caradja Aristide, *Die Grossschmetterlinge des Königreiches Rumänien*. In: *Deutsche Entomologische Zeitschrift, Iris* 8, Dresden (1895), p. 1-112.
- Ciochia, Barbu 1980 Ciochia Victor, Barbu Alexandru, *Catalogul colecției de lepidoptere „Nicolae Delvig” a Muzeului Județean Brașov*. In: *Cumidava* 12(4), Brașov (1980), p. 7-99.
- Czekelius 1897 Czekelius Daniel, *Kritisches Verzeichnis der Schmetterlinge Siebenbürgens*. In: *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* 47, Sibiu (1897), p. 1-78.
- Czekelius 1917 Czekelius Daniel, *Beiträge zur Schmetterlingsfauna Siebenbürgens*. In: *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* 67, Sibiu (1917), p. 1-57.
- Czekelius 1922 Czekelius Daniel, *Beiträge zur Schmetterlingsfauna Siebenbürgens*. In: *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* 71, Sibiu (1922), p. 71-77.
- Diószeghy 1933-1934 Diószeghy László, *Die Lepidopterenfauna des Retezatgebirges. Nachtrag I*. In: *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* 83-84, Sibiu (1933-1934), p. 107-126.
- Fleck 1904 Fleck Eduard, *Die Makrolepidopteren Rumäniens. Nachtrag II*. *Bulletin de la Société des Sciences București* 13 (3-4), București (1904), p. 288-303.
- Franzenau 1852 Franzenau Joseph, *Lepidopterologische Mitteilungen*. In: *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* 3, Sibiu (1852), p. 181-186.
- Franzenau 1859 Franzenau Joseph, *Verzeichniss der bis jetzt Siebenbürgen aufgefunden Lepidopteren*. In: *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* 10, Sibiu (1859), p. 25-27.
- Fuss 1850 Fuss Carl, *Verzeichniss der bis jetzt in Siebenbürgen aufgefundenen Lepidopteren*. In: *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* 1, Sibiu (1850), p. 54-64.
- König 1975 König Frederich, *Catalogul colecției de lepidoptere a Muzeului Banatului*, In: *Editura Muzeului Banatului*, Timișoara (1975).
- Mann 1866 Mann Josef, *Aufzählung der im Jahre 1865 in der Dobruđa gesammelten Schmetterlinge*. In: *Verhandlungen der kaiserlich-königlich zoologisch-botanischen Gesellschaft Wien* 16, Viena (1866), p. 1-40.
- Mihuț, Dincă 2006 Mihuț Sergiu, Dincă Vlad, *Important areas for lepidoptera in Romania*. In: *Centrul Focal pentru Monitorizarea și Conservarea Biodiversității (CFMBCB)*, Cluj-Napoca (2006).
- Niculescu 1961 Niculescu Eugen Victor, *Lepidoptera, Familia Papilionidae. Fauna Republicii Populare Române, Insecta* 11 (5). In: *Editura Academiei Republicii Populare Române*, București (1961).
- Pascu, Schneider 1998 Pascu Mariana, Schneider Eckbert, *Colecțiile entomologice ale Muzeului de Istorie Naturală din Sibiu*, In: *Studii și Comunicări, Științe Naturale Muzeul Bruckenthal Sibiu* 27, Sibiu (1998), p. 201-218.
- Popescu-Gorj 1964 Popescu-Gorj Aurelian, *Catalogue de la collection de lépidoptères „Prof. A. Ostrogovich” du Museum d'Histoire Naturelle „Grigore Antipa”*, Bucarest. In: *Editura Muzeului „Grigore Antipa”*, Bucharest (1964).
- Popescu-Gorj 1970 Popescu-Gorj Aurelian, *100 de ani de cercetări lepidopterologice în cadrul Societății Ardelene de Științe Naturale*. In: *Studii și Comunicări, Științe Naturale Muzeul Bruckenthal Sibiu* 15, Sibiu (1970), p. 85-97.
- Rákósy 1997 Rákósy László, *Die endemischen Lepidopteren Rumäniens*. In: *Entomologica Romanica* 2, Cluj-Napoca (1997), p. 59-81.
- Rákósy 2003 Rákósy László, *Lista roșie pentru fluturii diurni din România (Rote Liste der Tagfalter Rumäniens)*. In: *Buletin informativ Societatea lepidopterologică*

- română 13(1-4), Cluj-Napoca (2003), p. 7-18.
- Rákósy 2005 Rákósy László, U.E. și legislația pentru protecția lepidopterelor din România. In: *Buletinul de Informare Entomologică* 16(3-4), Cluj-Napoca (2005), p. 89-96.
- Rákósy et al. 2003 Rákósy László, Goia Marin, Kovács Zoltan, *Verzeichnis der Schmetterlinge Rumäniens. Catalogul lepidopterelor din România*. In: *Editura Societății Lepidopterologice Române*, Cluj-Napoca (2003).
- Rothschild 1912 Rothschild Charles Nathaniel, *Beitrag zur Lepidopteroфаuna der Mezőség*. In: *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* 62, Sibiu (1912), p. 1-32.
- Ruști, Dragomirescu 1991 Ruști Dorel, Dragomirescu Liviu, *A revision of Parnassius apollo (Lepidoptera, Papilionidae) in Romania using numerical taxonomy*. In: *Travaux du Muséum d'Histoire Naturelle "Grigore Antipa"* 31, Bucharest (1991), p. 201-218.
- Salay 1910 Salay Franz, *Katalog der Makrolepidopteren Rumäniens*. In: *Bulletin de la Société des Sciences București* 19, București (1910), p. 76-206; p. 453-616.
- Schneider 1984 Schneider Eckbert, *Die Groß-Schmetterlinge der Sammlung Dr. V. Weindel. Ein Beitrag zur Faunistik der Lepidopteren Sudsiebenbürgens und angrenzender Gebiete*. In: *Studii și Comunicări, Științe Naturale Muzeul Brukenthal Sibiu* 26, Sibiu (1984), p. 289-316.
- Schneider 2003 Schneider Eckbert, *Die Schmetterlingssammlung Heinrich von Hanneheim im Naturwissenschaftlichen Museum in Hermannstadt*. In: *Studii și Comunicări, Științe Naturale Muzeul Brukenthal Sibiu* 28, Sibiu (2003), p. 201-230.
- Székely 2008 Székely Levente, *The Butterflies of Romania – Fluturii de zi din România*. In: *Brastar Print*, Brașov (2008).
- Van Sway, Warren 1999 Van Sway Chris., Warren Martin, *Red Data Book of European Butterflies*. In: *Nature and Environment 99, Council of Europe Publishing*, Strassbourg (1999).
- Weindel 2000 Weindel Victor, *Der Apollofalter (Parnassius apollo Linnaeus, 1758) in Siebenbürgen*, In: *Entomologica Romanica* 5, Cluj-Napoca (2000), p. 5-10.
- Internet source [HTTP://ro.wikipedia.org/](http://ro.wikipedia.org/)

LIST OF ILLUSTRATIONS

- Fig. 1.** The number of specimens from the genus *Parnassius* Latreille, 1804 preserved in the Natural History Museum Collection from Sibiu
- Fig. 2.** The collecting sites of *Parnassius apollo* (Linnaeus, 1758) according to Natural History Museum Collection from Sibiu
- Fig. 3.** *Parnassius apollo* (Linnaeus, 1758) distribution according to Natural History Museum Collections from Sibiu
- Fig. 4.** The sample dates of *Parnassius apollo* (Linnaeus, 1758) according to the Natural History Museum Collection from Sibiu
- Fig. 5.** *Parnassius mnemosyne* (Linnaeus, 1758) distribution according to the Natural History Museum Collection from Sibiu
- Fig. 6.** The sample dates of *Parnassius mnemosyne* (Linnaeus, 1758) in the Natural History Museum Collection from Sibiu.

LISTA ILUSTRAȚIILOR

- Fig. 1.** Numărul specimenelor aparținând genului *Parnassius* Latreille, 1804 conservate la Muzeul de Istorie Naturală din Sibiu
- Fig. 2.** Punctele de colectare pentru specia *Parnassius apollo* (Linnaeus, 1758) după speciemenele conservate la Muzeul de Istorie Naturală din Sibiu
- Fig. 3.** Răspândirea speciei *Parnassius apollo* (Linnaeus, 1758) după exemplarele conservate la Muzeul de Istorie Naturală din Sibiu
- Fig. 4.** Datele colectării exemplarelor de *Parnassius apollo* (Linnaeus, 1758) de la Muzeul de Istorie Naturală din Sibiu
- Fig. 5.** Punctele de colectare pentru specia *Parnassius mnemosyne* (Linnaeus, 1758) după speciemenele conservate la Muzeul de Istorie Naturală din Sibiu
- Fig. 6.** Datele colectării exemplarelor de *Parnassius mnemosyne* (Linnaeus, 1758) de la Muzeul de Istorie Naturală din Sibiu




Fig. 1. The number of specimens from the genus *Parnassius* Latreille, 1804 preserved in the Natural History Museum Collection from Sibiu.




Fig. 2. The collecting sites of *Parnassius apollo* (Linnaeus, 1758) according to Natural History Museum Collection from Sibiu



Fig. 3. *Parnassius apollo* (Linnaeus, 1758) distribution according to Natural History Museum Collections from Sibiu; representations: square - *Parnassius apollo*; pentagon – *P. a. jaraensis*; circle - *P. a. transsylvanicus* (map source: <http://ro.wikipedia.org/>).




Fig. 4. The sample dates of *Parnassius apollo* (Linnaeus, 1758) according to the Natural History Museum Collection from Sibiu.




Fig. 5. *Parnassius mnemosyne* (Linnaeus, 1758) distribution according to the Natural History Museum Collection from Sibiu; representations: circle – *Parnassius mnemosyne*, square – *P. m. distincta*, pentagon – *P. m. transsylvanica* (map source: <http://ro.wikipedia.org/>).




Fig. 6. The sample dates of *Parnassius mnemosyne* (Linnaeus, 1758) in the Natural History Museum Collection from Sibiu.

**ASIOTMETHIS LIMBATUS (CHARPENTIER, 1845) (ORTHOPTERA: PAMPHAGIDAE)
IN ROMANIA – SPECIFIC HABITATS**

Gabriel LUPU*
Mihai DOROFTEI**

Abstract. Observed for the first time in south-eastern Romania by Ramme in 1951 and described as a new subspecies from Cogelac (Constanța County), *Asiotmethis limbatus motasi* (Ramme, 1951) is endemic for the Romanian territory, being known only from central and southern Dobrogea. The species was discovered in – situ by its calling song and body colour, two major elements in identifying this very mobile insect. There are three major NATURA 2000 habitat types where this taxa have been identified: Subpannonic steppe grasslands described as xerophilous feathergrass steppe grasslands, dry grasslands described as hill and plateau xero-mesophilous grassland, and limestone few fallow steppe grasslands from Dobrogea described as steppe grassland on hill limestone, with an average altitude oscillating between 100 and 150 meters a.s.l.
Keywords: *Asiotmethis limbatus*, Orthoptera, Romania, distribution, specific habitats.

Rezumat. Observată pentru prima dată în sud-estul României de Ramme în 1951 și descrisă ca subspecie nouă din Cogelac (județul Constanța), *Asiotmethis limbatus motasi* (Ramme, 1951) este endemică pentru teritoriului României, fiind cunoscută numai din Dobrogea centrală și sudică. Specia a fost identificată în teren pe baza stridulației și a desenului corporal, două elemente majore în identificarea acestei insecte foarte mobile. Există trei tipuri de habitate NATURA 2000 unde acest taxon a fost identificat: pajiști de stepă sub-panonice descrise ca pajiști de stepă xerofile cu colilie, pajiști xerofile seminaturale descrise ca pajiști xero-mezofile de deal și podiș, și pajiști seminaturale calcifile de stepă din Dobrogea descrise ca pajiști de stepă pe dealuri calcaroase, cu altitudine medie ce oscilează între 100 și 150 metri.
Cuvinte cheie: *Asiotmethis limbatus*, Orthoptera, România, distribuție, habitate specifice.

Introduction

Dobrogea territory is characterized by the presence of large mass of water, in the northern part – Danube Delta, and in the same time the presence of an important continental area represented by thermal extremes. The influence of the Black Sea proximity is significantly important both for climate and for influencing the presence and the distribution of the species and their characteristic habitats. Biologically the area is relatively isolated from the other Romanian regions, the presence of the *Bradyporus dasypus* (Illiger 1800) (Iorgu 2009, p.144) and *Asiotmethis limbatus motasi* (Ramme 1951) (Iorgu 2008, p.217) almost exclusively here being the evidence of this fact.

The eastern Balkan species, *Asiotmethis limbatus* (Charpentier, 1845), have been known as being present in Mangalia – Frey-Gessner, 1900, Tulcea – Frey-Gessner, 1902 and Murfatlar – Zotu, 1903 (Kis 1967, p.387). In the data published by the abovementioned author, the species was named as *Eremobia cisti* Brunner 1882 (Kis 1967, p.387).

At the same time, the author indicates a new record point in Agigea, the specimen being collected by C. Mândru.

In Romania, *Asiotmethis limbatus* is represented by the subspecies *motasi*, described by Ramme in 1951 and collected from the surroundings of Cogelac, Constanța County. Apparently this taxa is endemic to Romania. Previous records from literature indicate the subspecies presence in the southern part of Constanța County in the Natural Reserve Hagieni Forest (Iorgu 2008); in the north-eastern part of the same County, we found a strong population near the Istria village (unpublished). In Tulcea County, the subspecies was reported from the southern part of Babadag forest (Iorgu 2008, p.217) while the northern point of signalization has been recorded in the surroundings of Agighiol village (unpublished), this being the northern known limit of distribution for these subspecies.

In 2005, Togănel and Chimișliu mentioned *Asiotmethis limbatus* as part of the entomological collection of Oltenia Museum's Natural Science section in Craiova. The authors indicate one male

* Danube Delta National Institute, Tulcea
*glupu@indd.tim.ro; **doroftei@indd.tim.ro

specimen as being collected from Ciuperceni village, Dolj County, by Ioan Fîru in August 1976. It is worth mentioning that the subspecies has not been determined at taxonomic level. Furthermore, in Dolj County there are two localities with Ciuperceni name, Ciupercenii Noi and Ciupercenii Vechi villages. For the moment, this record remains unclear for both taxonomical information and localization, being the only observation of this taxa outside of the Dobrogea territory.

Dobrogea's Orthoptera fauna was intensely studied during the past century, Kis being the one who performed complex investigations. He provided the most complete descriptions for *Asiotmethis limbatus* in Romania. Apart from that, the area is characterized by the presence of 113 taxonomical elements (Iorgu *et al.* 2008, p.122, 128) belonging to *Orthoptera* order. Biogeographically the area where the taxon was found is of great importance, because it combines European, Mediterranean and Eurasian Orthoptera elements. It is also the interference point of continental, submediterranean, wet area, steppic and marine climate types as well.

This study offers data concerning the present distribution of *Asiotmethis limbatus* species in Romania and its habitat preferences.

Material and methods

The field investigations were carried out by employing the itinerary method, according to both vegetal associations and landscape forms. The species was identified by means of stridulation and body color, two major elements in identifying these insects. The captures were made using specific entomological mesh. In investigating the distribution of this taxa, the habitat characteristics were a significant reference point. The data was collected during the following research projects: the investigation of Dobrogea's Orthoptera fauna, the elaboration of the atlas of Orthoptera species from Danube Delta, the reconsideration of the limits for the functional area within the Danube Delta Biosphere Reserve, and investigations regarding the climate change in this area. The entomofaunistic studies were conducted during 8 years (2004 - 2011) and provided a series of of interesting elements. The taxa were found in NATURA 2000 habitats and they were described according to the Romanian Grasslands report (Sârbu *et al.* 2004).

The images were taken with a Canon S5IS photo camera.

Results and discussion

Genus *Asiotmethis* belongs to the class *Insecta*, order *Orthoptera*, suborder *Caelifera*, infraorder *Acrididea*, superfamily *Acridoidea*, family *Pamphagidae*, subfamily *Prionotropisinae* (Iorgu *et al.* 2008, p.128). Currently there are 24 species and subspecies known in to this genus, widely distributed in central and western Asia and southeastern Europe (Ünal 2007, p.191). It is recorded from eastern Siberia, southern Russia, northwest China, Kazakstan, Kyrgyzstan, Uzbekistan, northern Tajikistan, northern Iran, Azerbaijan, Armenia, northern Turkey, southern Ukraine, northern Greece, Macedonia and Romania (Ünal 2007, p.196). Most of the species are from central Asia.

Diagnosis. *Asiotmethis limbatus* has the metazona of pronotum short and narrow, less raised; hindwing yellow at base, its dark band short, never attaining basal part; inner side of hind legs orange or yellow; arolium small, reaching only to a half-length of claws or less; apical part of penis valves slender, their tips very narrow; posterior margin of epiphallus with a broadly rounded projection (Ünal 2007, p.192).

Ethology. Diurnal insect, *Asiotmethis limbatus* has xerophilous preferences and enjoys sunny places with low and sparse vegetation, generally in the vicinity of some wet areas. In spite of his medium to large corporal dimensions size, the calling song is somehow discreet.

Adults appear in late July and they can be seen until the end of September. It hibernates in egg stage and the larvae hatch in May.

Body length: male 25-28 mm (Fig. 1) (52 individuals measured); female 30-35 mm (Fig. 2) (74 individuals measured).

Habitats. *Asiotmethis limbatus* was observed in five collecting-sites in Dobrogea only (Fig. 5.):

- **Hagieni reserve (Constanța County)** (43° 47' 56" N, 28° 28' 39" E)

In Hagieni area the Natura 2000 habitat is 6240 - Subpannonic steppe grasslands - and is described as xerophilous feathergrass steppe grasslands. The plant species composition is mainly characterized by *Stipa lessingiana*, *Agropyron cristatum* ssp. *pectinatum*, *Festuca valesiaca*, *Centaurea marschalliana*, *Bombycilaena erecta*, *Taraxacum serotinum*, *Euphorbia nicaeensis* and *Dichanthium ischaemum*, forming a plant association

Bombycilaeno - Botriochloetum Dihoru et Doniță 1970.

The average altitude is 250 meters a.s.l., with a geological description of 65% - marl-clay deposits, 20% - grit-stone, 15% - sands. The climatic conditions for this flat area are 85% - warm and very warm, 15% - weak cold from temperature perspective, and from humidity point of view 85% - dry, 15% - few wet. The average cover of ligneous plants is 3 % (Sârbu *et al.* 2004, 31).

The second Natura 2000 habitat 4030 is in Hagieni area - Dry grasslands described as hill and plateau xero-mesophilous grasslands. The plant species composition is mainly characterized by *Achillea coarctata*, *Dichanthium ischaemum*, *Carex humilis*, *Convolvulus cantabrica*, *Dorycnium herbaceum*, *Festuca valesiaca*, *Tragopogon flocosus*, *Oxytropis pilosa*, *Stipa capillata*, *Iris pumila*, *Sedum sartorianum* ssp. *hillebrandtii*, *Centaurea trinervia*, *Salvia nutans*, *Dictamnus albus*, *Carpinus orientalis*, *Eryngium campestre* and *Goniolimon tataricum*, thus forming a plant association *Artemisio austriacae - Poetum bulbosae* Pop 1970.

The average altitude is 350 meters a.s.l., with a geological description of 45% - marl deposits, 15% - calcareous rocks and 10% - flysch. The climatic conditions for this flat area are 80% - warm and very warm, 20% - moderately cold from temperature perspective and from humidity point of view 80% - dry, 20% - moderately wet. The average cover of ligneous plants is 5 % (Sârbu *et al.* 2004, p. 30).

- **Istria village (Constanța County)** (44° 33' 50" N, 28° 42' 23" E)

In Histria area (Fig. 3.) the Natura 2000 habitat type is 2130* - Fixed coastal dunes with herbaceous vegetation (grey dunes) (Schneider, Drăgulescu 2005, p. 51), and is described as dry continental dune grassland. The plant species composition is mainly characterized by *Achillea ochroleuca*, *Alyssum desertorum*, *Bassia laniflora*, *Bromus tectorum*, *Centaurea diffusa*, *Carex stenophylla*, *Chondrilla juncea*, *Dianthus diutinus*, *Festuca vaginata*, *Euphorbia seguieriana*, *Polygonum arenarium*, *Plantago arenaria*, *Koeleria glauca* and *Helichrysum arenarium*, thus forming a plant association *Brometum tectorum* Bojko 1934.

The average altitude is 50 meters a.s.l., with a geological description of 90% - sands, 10% -

alluvial sands. The climatic conditions for this flat area are 84% - warm and very warm, 16% - cold from temperature perspective and from humidity point of view 84 - dry, 16% - moderately wet (Sârbu *et al.* 2004, 26).

- Southern part of **Babadag Forest (Tulcea County)** (44° 48' 29" N, 28° 42' 45" E)

In Babadag area the Natura 2000 habitat is 6110 - Limestone few fallow steppe grassland from Dobrogea and is described as steppe grassland on hill limestone. The plant species composition is mainly characterized by *Agropyron cristatum*, *Euphorbia nicaeensis*, *Dianthus nardiformis*, *Thymus zygioides*, *Pimpinella tragioides* ssp. *lithophila*, *Scorzonera mollis* and *Teucrium polium* ssp. *capitatum*, thus forming a plant association *Festucetum callierii*, Șerbănescu 1965.

The average altitude is 250 meters a.s.l., with a geological description of 55% - limestone rocks, 30% - conglomerate rocks, 15% - gritstone. The climatic conditions for this flat area are 85% - warm and very warm, 15% - moderately cold from temperature perspective and from humidity point of view 85% - dry, 15% - moderately wet. The average cover of ligneous plants is 2 % (Sârbu *et al.* 2004, p.33).

The second Natura 2000 habitat type found in Babadag area is 6240 - Subpannonic steppe grasslands and is described as xerophilous feathergrass steppe grasslands. The plant species composition is mainly characterized by *Stipa lessingiana*, *Festuca valesiaca*, *Bombycilaena erecta*, *Dianthus capitatus*, *Euphorbia nicaeensis*, *Dichanthium ischaemum*, thus forming a plant association *Bombycilaeno - Botriochloetum* Dihoru et Doniță 1970.

The average altitude is 250 meters a.s.l., with a geological description of 65% - marl-clay deposits, 20% - grit-stone, 15% - sands. The climatic conditions for this flat area are 85% - warm and very warm, 15% - weak cold from temperature perspective and from humidity point of view 85% - dry, 15% - few wet. The average cover of ligneous plants is 3 % (Sârbu *et al.* 2004, p.31).

- In the surroundings of **Agighiol village (Tulcea County)** (45° 01' 26" N, 28° 51' 44" E)

In Agighiol area (Fig. 4.), the Natura 2000 habitat is 4030 - Dry grasslands from Dobrogea and is described as steppe grassland on hill and plateau xero-mesophilous grasslands. The plant species composition is mainly characterized by *Artemisia*

austriaca, *Dichanthium ischaemum*, *Festuca valesiaca*, *Cleistogenes serotina*, *Stipa capillata*, *Goniolimon tataricum*, *Oxytropis pilosa*, *Poa bulbosa* and *Medicago lupulina*, thus forming a plant association *Medicagini – Festucetum valesiaca* Wagner 1941, *Artemisia austriacae – Poetum bulbosae* Pop 1970.

The average altitude is 350 meters a.s.l., with a geological description of 45% - marl deposits, 15% - calcareous rocks, 10% - flysh. The climatic conditions for this flat area are 80% - warm and very warm, 20% - moderately cold, from temperature perspective and from humidity point of view 80% - dry, 20% - moderately wet. The average cover of ligneous plants is 5 % (Sârbu *et al.* 2004, p.30).

- In central eastern part of **Tulcea County – Movila Cazacului Hill** (in the southern part of **Peceneaga village**) (44° 59' 39" N, 28° 08' 19" E)

In Movila Cazacului area, the Natura 2000 habitat is 6110 - Limestone few fallow steppe grasslands from Dobrogea and is described as steppe grassland on hill limestone. The species composition is mainly characterized by *Festuca callieri*, *Agropyron cristatum* ssp. *brandzae*, *Pimpinella tragi* ssp. *lithophila*, *Thymus zygoides*, *Teucrium polium* ssp. *capitatum* and *Allium saxatile*, thus forming a plant association *Festucetum callierii* Șerbănescu 1965.

The average altitude is 250 meters a.s.l., with a geological description of 55% - limestone rocks, 30% - conglomerate rocks, 15% - gritstone. The climatic conditions for this flat area are 85% - warm and very warm, 15% - moderately cold from temperature perspective and from humidity point of view 85% - dry, 15% - moderately wet. The average cover of ligneous plants is 2 % (Sârbu *et al.* 2004, p.33).

Conclusions

Species with relative large distribution, known from the South-eastern part of Europe and Asian part of Istanbul, *Asiotmethis limbatus* presents some special characteristics in Romania:

- *Asiotmethis limbatus* subspecies *motasi* lives only in Romania, Dobrogea being apparently the single area where its presence has been manifested, the presence in Dolj County being in doubt;
- Being diurnal and predominantly thermophile species, it prefers open spaces with powerful insolation, and low vegetation, near wetland areas;
- There are three major NATURA 2000 habitat where these taxa have been identified: 6240 - Subpannonic steppe grasslands described as xerophilous feathergrass steppe grasslands, 4030 - Dry grasslands described as hill and plateau xeromesophilous grasslands, and 6110 - Limestone few fallow steppe grasslands from Dobrogea described as steppe grassland on hill limestone, on average altitude oscillating between 100 and 150 meters a.s.l.

REFERENCES

- Iorgu 2008 Iorgu Ionuț, *Diversitatea ortopterelor (Insecta: Orthoptera) din estul României și semnificația lor ecologică* (2008) Iași, PhD Thesis (manuscript).
- Iorgu et al. 2008 Iorgu Ionuț, Pisică Elena, Păiș Laura, Lupu Gabriel, Iușan Claudiu, *Checklist of Romanian Orthoptera (Insecta: Orthoptera) and their distribution by ecoregions*. In: Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa LI (2008) Bucharest, p.119-135.
- Iorgu 2009 Iorgu Ionuț, *Bradyporus dasypus (Illiger, 1800) (Orthoptera: Tettigoniidae): some ethological aspects and distribution in Romania*. In: Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa LII (2009) Bucharest, p.143-149.
- Kis 1967 Kis Bela, *Studiul ortopterelor din R. S. România (sistematic, faunistic, zoogeografic și ecologic)*, (1967) Cluj-Napoca, PhD Thesis (manuscript).
- Ramme 1951 Rame Willy, *Zur Systematik Faunistik und Biologie der Orthopteren von Sudost-Europa und Vorderasien*. In: Mitteilungen Aus Dem Zoologischen Museum zu Berlin 27 (1951) Berlin, p. 383-385.
- Sârbu et al. 2004 Sârbu Anca, Coldea Gheorghe, Negrean Gavril, Cristea Vasile, Hanganu Jenică, Peter Veen, *Grasslands of Romania, Final report on National Grasslands Inventory 2000-2003*, Publisher University of Bucharest (2004) Bucharest, p. 26-33.
- Schneider, Drăgulescu 2005 Schneider Erika, Drăgulescu Constantin, *Habitate și situri de interes comunitar*. In: Editura Universității „Lucian Blaga”, (2005) Sibiu.
- Togănel, Chimișliu 2005 Togănel Florentina, Chimișliu Cornelia, *Orthopterans Species belonging to the entomological collection of the Natural Sciences Section Of The Oltenia Museum From Craiova*. In: Oltenia. Studii și comunicări. Științele Naturii XXI (2005) Craiova, p. 67-72.
- Ünal 2007 Ünal Mustafa, *Genus Asiotmethis Uvarov (Orthoptera, Pamphagidae) from Turkey, with a checklist of known taxa*. In: Journal of Orthoptera Research 16(2) (2007) Istanbul, p. 191-197.

LIST OF ILLUSTRATIONS

- Fig. 1.** *Asiotmethis limbatus motasi* ♀, 09.07.2009 - Istria (photo: Gabriel Lupu)
- Fig. 2.** *Asiotmethis limbatus motasi* ♂, 26.06.2008 - Agighiol (photo: Gabriel Lupu)
- Fig. 3.** Specific habitat – Istria area (photo: Gabriel Lupu)
- Fig. 4.** Specific habitat – Agighiol area (photo: Gabriel Lupu)
- Fig. 5.** Distribution of *Asiotmethis limbatus motasi* according to collecting sites (satellite image)

LISTA ILUSTRAȚIILOR

- Fig. 1.** *Asiotmethis limbatus motasi* ♀, 09.07.2009 - Istria (foto: Gabriel Lupu)
- Fig. 2.** *Asiotmethis limbatus motasi* ♂, 26.06.2008 - Agighiol (foto: Gabriel Lupu)
- Fig. 3.** Habitat specific – zona Istria (foto: Gabriel Lupu)
- Fig. 4.** Habitat specific – zona Agighiol (foto: Gabriel Lupu)
- Fig. 5.** Distribuția speciei *Asiotmethis limbatus motasi* conform punctelor de colectare (imagine satelitară)



Fig. 1. *Asiotmethis limbatus motasi* ♀, 09.07.2009 - Istria (photo: Gabriel Lupu)



Fig. 2. *Asiotmethis limbatus motasi* ♂, 26.06.2008 - Agighiol (photo: Gabriel Lupu)



Fig. 3. Specific habitat – Istria area (photo: Gabriel Lupu)



Fig. 4. Specific habitat – Agighiol area (photo: Gabriel Lupu)




Fig. 5. Distribution of *Asiotmethis limbatus motasi* according to collecting sites (satellite image)

FIRST RECORD OF *ISOPHYA PIENENSIS* IN THE WESTERN ROMANIAN CARPATHIANS (ORTHOPTERA: PHANEROPTERIDAE)

Ionuț Ștefan IORGU*

Abstract. The bush-cricket *Isophya pienensis* Mařan, 1952 is recorded for the first time from the Western Romanian Carpathians. Minor acoustic and morphological variations were noticed between the four studied populations from Romania: Breaza (Suceava County), Bistricioara (Neamț County), Baia Mare (Maramureș County) and the newly found one at Roșia Montană (Alba County).

Keywords: *Isophya pienensis*, bioacoustics, Carpathians, Romania

Rezumat. Cosașul *Isophya pienensis* Mařan, 1952 este menționat pentru prima dată din Carpații Occidentali, România. Mici variații acustice și morfologice au fost notate între cele 4 populații studiate din România: Breaza (județul Suceava), Bistricioara (județul Neamț), Baia Mare (județul Maramureș) și nou-găsită populație de la Roșia Montană (județul Alba).

Cuvinte cheie: *Isophya pienensis*, bioacustică, Carpați, România

Introduction

Currently there are 45 species of *Isophya* Brunner von Wattenwyl in Europe and 17 of these have been found in Romania, 4 being endemic: *Isophya harzi* Kis, *Isophya dobrogensis* Kis, *Isophya sicula* Orci, Szövényi & Nagy and *Isophya ciucasi* Iorgu & Iorgu (Eades *et al.* 2012).

Isophya pienensis Mařan, 1952 is known to occur in Austria, Czech Republic, Slovakia, Poland, Ukraine and Romania (Heller 2012). In Romania, the species populates the mountain meadows of the Eastern Carpathians in Maramureș, Transylvania (Kis 1964, Kis, Vasiliu 1970, Szövényi, Orci 2008), Bucovina and Moldavia (Iorgu 2011). During author's regular field research in Orthoptera, the bush-cricket *Isophya pienensis* was surprisingly found in the Metaliferi Mountains, Western Romanian Carpathians.

This bush-cricket is characterized by a broad left tegmen, angularly convex at the tip of the stridulatory vein and gradually curved inwards cerci, with a detached tooth at the upper edge of the tip (Heller *et al.* 2004). Females have a short ovipositor, 9-10 mm long.

Two subspecies have been recently described from Slovakia, *Isophya pienensis austromoravica* Chládek, 2010 and *Isophya pienensis sudetica* Chládek, 2011.

* Grigore Antipa National Museum of Natural History, nusi81@yahoo.com

Material and methods

All the specimens were collected as adults in a single trip at Roșia Montană, on the 9th of July 2011. Acoustic signals were recorded with the digital recorder EDIROL R-09HR, whose built-in microphones have the sound frequency response between 20-40.000 Hz (96.000 Hz sampling rate). Sound analysis was performed with the software Audacity 2.

Song terminology used: syllable - the sound produced by one complete up (opening) and down (closing) stroke of the forewings; impulse - the highly damped sound impulse resulting from the impact of one tooth of the stridulatory file (Heller *et al.* 2004).

Photos were taken with a Canon EOS DSLR camera and two lenses attached: a Canon 2.8 100 mm 1:1 macro lens was used for habitus photos and a Canon MP-E 65 mm 5:1 macro lens for photos of the morphological details. Male wing movements have been video recorded with the same camera.

Results

Order Orthoptera

Family Phaneropteridae

Isophya pienensis Mařan, 1954

Material. 6♂♂ 3♀♀, Roșia Montană, Metaliferi Mountains, Alba County, Romania, 2011.07.09, leg. I. Ș. Iorgu (Fig. 1).

Comparative material. 2♂♂ 1♀♀, Breaza, Suceava County, Romania, 47°38'31.30"N

25°18'45.81"E, 2011.06.27 (Fig. 1); 7♂♂ 5♀♀ Izvoarele Sucevei, Suceava County, Romania, 47°45'36.59"N 25°12'29.31"E, 2011.06.27; 3♂♂ 4♀♀, Borsec Pass, Harghita County, Romania, 46°57'35.92"N 25°29'24.72"E, 2011.06.30; 2♂♂, Bradu, Neamț County, Romania, 46°59'55.64"N 25°50'55.70"E, 2011.07.01; 3♂♂ 2♀♀, Bistricioara, Neamț County, Romania, 47°03'14.13"N 25°56'33.87"E, 2011.07.01; 2♂♂, Baia Mare, Maramureș County, Romania, 47°40'43.42"N 23°34'59.55"E, 2012.06.07; 1♂ 1♀, Dorna Candrenilor, Suceava County, Romania, 47°21'22.34"N 25°17'16.61"E, 2012.06.15; 3♂♂ 1♀, Hășmașul Mare Mts., Harghita County, Romania, 46°41'14.87"N 25°49'33.11"E, 2012.07.12; 2♂♂ 2♀♀, St. Ana Lake, Harghita County, Romania, 46°07'16.46"N 25°54'08.47"E, 2012.08.06, leg. I. Ș. Iorgu (Fig. 2).

Audio recordings. 2♂♂ recorded in the field at Breaza, temperature 20°C, 2011.06.27; 2♂♂ recorded in the laboratory, Bistricioara, temperature 22°C, 2011.07.01; 3♂♂ recorded in the field at Roșia Montană, temperature 24°C, 2011.07.09; 1♂ recorded in the laboratory, Baia Mare, temperature 23°C, 2012.06.07 (wave files, 24 bits/96 kHz).

Literature records: Borșa, Maramureș County, Romania (Kis 1964); Țibleș Mountains, Rodnei Mountains, Bistrița-Năsăud County, Romania; Borsec, Sovata, Mureș County, Romania; Harghita, Tușnad, Harghita County, Romania (Kis, Vasiliu 1970); Săpânța, Mara, Gutâi Pass, Poienile de Sub Munte, Maramureș County, Romania (Szövényi, Orci 2008) (Fig. 2).

Bioacoustics

The calling song consists of small groups of two or three syllables, repeated after several seconds or minutes, but isolated syllables have also been recorded. A syllable consists of a series of 75-115 impulses, lasting for about 250-500 ms at 20-24°C (Fig. 3). In the last 20-30 impulses, the amplitude of sound gradually decreases. All sounds are produced when the insect closes the tegmina.

The stridulatory file is 3.2-3.6 mm long and bears 205-220 teeth.

The spectrographic analysis reveals the most intense frequencies between 10-40 kHz, with the maximum amplitude at: 28 kHz (Baia Mare), 19 kHz (Bistricioara), 14 kHz (Breaza) and 23 kHz (Roșia Montană) (Fig. 3).

Habitat

At Roșia Montană, near Tăul Mare Lake, *Isophya pienensis* lives in mesophytic grasslands. The specimens were collected from broad leaves of *Urtica*, *Stachys*, *Rubus*, *Salvia*, *Centaurea* etc. The following bush-crickets and grasshoppers were found occurring simpatrically with *Isophya pienensis*: *Isophya stysi* Cejchan, 1957, *Tettigonia viridissima* (Linnaeus, 1758), *Decticus verrucivorus* (Linnaeus, 1758), *Metrioptera bicolor* (Philippi, 1830), *Pholidoptera transsylvanica* (Fischer, 1853), *Pseudopodisma transilvanica* Galvagni & Fontana, 1993, *Arcyptera fusca* (Pallas, 1773), *Euthystira brachyptera* (Ocskay, 1826), *Stauroderus scalaris* (Fischer de Waldheim, 1846), *Chorthippus parallelus* (Zetterstedt, 1821) etc.

Discussion

The bush-cricket *Isophya pienensis* was known in Romania only from the Central and Northern parts of the Eastern Carpathians and most probably the species has the Eastern distribution area limit near the lake Izvorul Muntelui (Bistricioara) (Fig. 2). The new collecting site from the Western Romanian Carpathians is relatively isolated from the other ones, located at more than 150 km from the nearest known locations. No major differences were revealed when we analyzed male acoustic signals, tegmina and cerci in 4 populations: Baia Mare, Breaza, Bistricioara and Roșia Montană (Figs. 3 and 4). However, the male from Baia Mare presents thinner cerci, while the ones from Breaza tend to have them less curved. The same males from Breaza have somewhat shorter wings, but the venation is constant in all populations. Syllables produced by males from Breaza are longer (340-500 ms) compared with the other studied populations (250-380 ms), but the same results were obtained by Heller *et al.* (2004) in individuals from Slovakia, Poland and Ukraine. The males from the newly discovered populations near Roșia Montană do not differ in any morphological or acoustical aspects from the ones in other populations.

Acknowledgements

Biologists Mr. Călin Hodor and Dr. Cosmin Ovidiu Mancu are acknowledged for making possible the trip to Roșia Montană and for excellent field work conditions. I am grateful to the anonymous reviewers that improved the paper.

REFERENCES

- Eades *et al.* 2012 Eades David, Otte Daniel, Cigliano María Marta, Braun Holger, *Orthoptera Species File Online* version 2.0/4.1, <http://Orthoptera.SpeciesFile.org>. (19.03.2012).
- Heller 2012 Heller Klaus - Gerhard, *Fauna Europaea: Arthropoda, Insecta, Orthoptera. Fauna Europaea* version 2.4, <http://www.faunaeur.org> (02.04.2012)
- Heller *et al.* 2004 Heller Klaus - Gerhard, Orci Kirill Márk, Grein Günther, Ingrisch Sigfrid, *The Isophya species of Central and Western Europe (Orthoptera: Tettigonioidea:Phaneropteridae)*. In: *Tijdschrift voor Entomologie* 147 (2004), p. 237-258.
- Iorgu 2011 Iorgu Ionuț Ștefan, *Bioacoustics of two newly recorded bush-crickets in the historical region of Moldavia: Isophya pienensis and Isophya sicalis (Insecta: Orthoptera)*. In: *Actual problems of protection and sustainable use of the animal world diversity*, Chișinău (2011), p. 117-119.
- Kis 1964 Kis Béla, *Contribuții la cunoașterea orthopterelor din R.P.R.* In: *Studii Universitatea Babeș-Bolyai* 2, Cluj-Napoca (1964), p. 69-73.
- Kis, Vasiliu 1970 Kis Béla, Vasiliu Maria, *Kritisches Verzeichnis des Orthopteren-Arten Rumäniens*. In: *Travaux du Muséum d'Histoire Naturelle "Grigore Antipa"* 10 (1970), p. 207-227.
- Szövényi, Orci 2008 Szövényi Gergely, Orci Kirill Márk, *Contributions to the Orthoptera fauna of Maramureș, with the first record of Isophya posthumoidalis Bazyluk, 1971 in Romania*. In: *Studia Universitatis Vasile Goldiș* 18 (2008), p. 235-241.

LIST OF ILLUSTRATIONS

- Fig. 1.** *Isophya pienensis* habitus: a. male, Roșia Montană, 2011.07.09; b. female, Roșia Montană, 2011.07.09; c. male, Breaza, 2011.06.27; d. female, Breaza, 2011.06.27.
- Fig. 2.** Distribution of species *Isophya pienensis* in Romania (own data - red points, literature data - blue points).
- Fig. 3.** Oscillographic (a-h) and spectrographic (i-l) sound analysis in *Isophya pienensis*: Baia Mare (a, b, i); Bistricioara (c, d, j); Breaza (e, f, k); Roșia Montană (g, h, l).
- Fig. 4.** *Isophya pienensis*: dorsal view of male wings (a, b, c, d) and cerci (e, f, g, h): Baia Mare (a, e); Bistricioara (b, f); Breaza (c, g); Roșia Montană (d, h).

LISTA ILUSTRAȚIILOR

- Fig. 1.** *Isophya pienensis* habitus: a. mascul, Roșia Montană, 2011.07.09; b. femelă, Roșia Montană, 2011.07.09; c. mascul, Breaza, 2011.06.27; d. femelă, Breaza, 2011.06.27.
- Fig. 2.** Distribuția speciei *Isophya pienensis* în România (date proprii - puncte roșii, date din literatura - puncte albastre).
- Fig. 3.** Analiza oscilografică (a-h) și spectrografică (i-l) a sunetului la *Isophya pienensis*: Baia Mare (a, b, i); Bistricioara (c, d, j); Breaza (e, f, k); Roșia Montană (g, h, l).
- Fig. 4.** *Isophya pienensis*: vedere dorsală asupra aripilor (a, b, c, d) și cercilor (e, f, g, h): Baia Mare (a, e); Bistricioara (b, f); Breaza (c, g); Roșia Montană (d, h).



Fig. 1. *Isophya pienensis* habitus




Fig. 2. Distribution of species *Isophya pienensis* in Romania (own data - red points, literature data - blue points).




Fig. 3. Oscillographic (a-h) and spectrographic (i-l) sound analysis in *Isophya pienensis*: Baia Mare (a, b, i); Bistricioara (c, d, j); Breaza (e, f, k); Roșia Montană (g, h, l).




Fig. 4. *Isophya pienensis*: dorsal view of male wings (a, b, c, d) and cerci (e, f, g, h): Baia Mare (a, e); Bistricioara (b, f); Breaza (c, g); Roșia Montană (d, h).

ARMOURED CRICKETS (ORTHOPTERA: TETTIGONIIDAE, BRADYPORINAE) IN THE NATURAL HISTORY MUSEUM COLLECTIONS OF SIBIU (ROMANIA)

Alexandru Ioan TATU*

Ioan TĂUȘAN**

Abstract. *The present paper contains data regarding the armored crickets (Bradyporinae) from the collections of the Natural History Museum of Sibiu preceded by brief details about the subfamily and the collections. The distribution maps of the collecting sites for the identified species are also provided. The preserved material is part of the following collections: “Dr. Arnold Müller”, “Rolf Weyrauch”, “Dr. Eugen Worell” and “Dr. Eckbert Schneider”.*

Keywords: *Bradyporinae, museum collections, systematical inventory, natural heritage.*

Rezumat. *În studiul de față, sunt oferite date asupra speciilor de Bradyporinae din colecțiile entomologice ale Muzeului de Istorie Naturală din Sibiu, împreună cu o serie de scurte informații legate de subfamilie respectiv de colecții. Hărțile de distribuție ale punctelor de colectare a speciilor identificate sunt de asemenea cuprinse în acest studiu. Materialul studiat face parte din următoarele colecții: “Dr. Arnold Müller”, “Rolf Weyrauch”, “Dr. Eugen Worell” și “Dr. Eckbert Schneider”.*

Cuvinte cheie: *Bradyporinae, colecții muzeale, inventar sistematic, patrimoniu natural.*

Introduction

The subfamily Bradyporinae comprises about 155 species (Eades *et al.* 2012). Its members are bulky, rather lazy bushcrickets with their forewings almost entirely covered by the pronotum, while the hindwings are absent. The antennae articulate below the eye level, closer to the clypeus rather than to the vertex (Harz 1969; Iorgu, Iorgu 2008). Their body colour varies from dark, metallic shining to light ochre or green with dark markings, “often with hieroglyph-like streaks or spots” (Harz 1969). For a more detailed description of their morphology see the aforementioned study. Concerning their distribution in Europe, they do not occur in the North, being limited to its central, southern and eastern parts (Fig. 1). Worldwide, according to Heller (2012), they can be found in the Near East and in the northern areas of Africa.

In Romania, several other papers exclusively concerning this subfamily have been undertaken by Vasiliu (1961), Kis (1962) and Iorgu (2009). Apart from these, the Orthoptera in the “Arnold Müller” collection have been researched by Vasiliu and Agapi who in 1958 published a catalogue of the collection.

* Babeș-Bolyai University of Cluj-Napoca,
alex_tatu@ymail.com, ioan.tausan@ubbcluj.ro

**Bruckenthal National Museum, Natural History Museum,
itausan@gmail.com

The entomological collections hosted at the Natural History Museum of Sibiu are some of the oldest and most valuable collections in Romania, dating back to 1827 (Pascu & Schneider, 1998).

This study aims to give an account of the armoured crickets present in the collections of the Sibiu museum.

Material

The preserved material is part of the four collections described below.

The “Arnold Müller” collection of Transylvanian and Palearctic Orthoptera is part of the wider Transylvanian Society for Nature Science of Sibiu collection and comprises 5.133 specimens. There are 12 species of armoured crickets in this collection. The “Rolf Weyrauch” entomological collection contains 16.436 specimens, of which 155 belong to the order Orthoptera and two species to the subfamily Bradyporinae. The specimens were collected mainly from 1950 to 1965 from Transylvania, especially from the surroundings of Sibiu. The “Eugen Worell” entomological collection contains 93.897 specimens (1041 Orthoptera). Its great value is given not only by this numbers but also by the fact that many of the species are rare, little known and some even new (Pascu & Schneider, 1998). There are two species of Bradyporinae in this collection. The “Eckbert Schneider” collection comprises 20.000 insect

specimens collected from all over the country, but mainly from the southern area of Transylvania. As in the previous collections, two species of armoured crickets have been found. The collection is still in state of processing (Pascu & Schneider, 1998). All four collections contribute to the knowledge of the species' distribution.

For the nomenclature and systematical order, we followed Eades *et al.* (2012) and Heller (2012).

The following abbreviations will be used in this paper: BG - Bulgaria; DZ - Algeria; ES - Spain; FR - France; GR - Greece; HR - Croatia; RO - Romania; RS - Serbia; Mts. - Mountains; spec./specs - specimen/specimens.

Listing of the species

Order Orthoptera
Family Tettigoniidae
Subfamily Bradyporinae

“Arnold Müller” Collection

Bradyporus Charpentier, 1825

Bradyporus (Bradyporus) dasypus (Illiger, 1800)

1 spec., Bălcic (Balcic, BG), VII. 1926, leg. Lepși;
1 spec., Balcic (BG), VIII.1927; 1 spec., Babadag,
Tulcea (Tulcea County, RO), 25.VII.1928; 1 spec.,
Caliakra, Bolatasteppe, Eichenm. (Bolata,
Kaliakra, BG), leg. Müller, 22.VII.1926; 2 specs.,
Kaliakra, Bolatasteppe, Eichenm. (Bolata, Kaliakra,
BG), leg. Müller, 22.VII.1926; 1 spec., Kaliakra,
Bolatasteppe, Eichensteppe (Bolata, Kaliakra, BG),
leg. Müller, 22.VII.1926; 1 spec., Kaliakra,
Bolatasteppe (Bolata, Kaliakra, BG), leg. Müller,
22.VII.1926; 1 spec., Dobrudscha, Maciner Berge
(Măcin Mts., Dobrogea, RO), leg. Müller,
12.VII.1927; 1 spec., Slava-Rusă, Tulcea (Slava
Rusă, Tulcea County, RO), 5.VII.1928; 1 spec.,
Slava-Rusă (Slava Rusă, Tulcea County, RO),
VII.1928.

Bradyporus (Callimenus) macrogaster longicollis
(Fieber, 1853)

1 spec., Serbia, Nisch (Niš, RS), leg. V. Örtzen; 1
spec., Caliacra, Bolatasteppe, Eichenm. (Bolata,
Kaliakra, BG), leg. Müller, 22.VII.1926.

Bradyporus (Callimenus) montandoni (Burr, 1898)

1 spec., Greci-Olt (Greci, Olt County, RO),
17.VI.1927; 2 specs., Plopșor, Dolju (Plopșor, Dolj
County, RO), VII.1925;

Comments: The material is labelled as *Callimenus oniscus* in the collections, however the species is not found in Romania (Iorgu *et al.* 2008).

Bradyporus (Callimenus) oniscus (Burmeister,
1838)

4 specs., Kaliakra, Bolatasteppe, Eichenm.
(Bolata, Kaliakra, BG), leg. Müller,
22.VII.1926; 1 spec., Balcic (Balcic, BG),
VII.1927; 1 spec., Serbia, Nisch (Niš, RS), leg.
Örtzen; 1 spec., Graecia, Atika (Attica, GR), leg.
Örtzen.

Ephippiger Berthold, 1827

Ephippiger ephippiger (Fiebig, 1784)

2 specs., Caracal, Rămănen (Caracal, Olt County,
RO), leg. Despaletz, 21.IX.1923; 1 spec.,
Herculesbad (Băile Herculane, Caraș-Severin
County, RO), leg. Müller, 26.VIII.1925; 1 spec.,
leg. Müller, 26.VIII.1925.

Comments: The material is labeled as *E. vitium* in
the collection, but this species is also not found in
Romania (Iorgu *et al.* 2008).

Ephippiger diurnus cunii Bolívar, 1877

1 spec., Bulgarien, Pirin, Elteppe (Pirin Mts., BG),
leg. Müller, 2.VIII.1931; 1 spec., Bulgarien, Pirin,
Elteppe (Pirin Mts., BG), leg. Müller, 7.VIII.1931;
2 specs., Bulgarien, Pirin, Elteppe (Pirin Mts., BG),
leg. Müller, 7.VIII.1931; 1 spec., Rivesaltes (FR),
VI.1891.

Comments: The first four specimens, collected
from Bulgaria cannot be *Ephippiger diurnus cunii*,
as this species' distribution is limited to Spain and
France (Eades *et al.* 2012).

Ephippiger provincialis Yersin, 1854

syn. *Ephippigera provincialis* (Yersin 1854)

1 spec., Collection Lucien Chopard 1919, Museum
Paris, Banyuls (FR), det. Bolívar, 9.IX.1909; 1
spec., Museum Paris, St. Tropez (FR), leg.
Bossavy, 1891.

Ephippiger discoidalis Fieber, 1853

syn. *Ephippigera sphacophila* Krauss, 1879

1 spec., Dalmația (HR), leg. Örtzen.

Uromenus Bolívar, 1878

Uromenus laticollis (Lucas, 1846-1849)

1 spec., ♀, Chabet el Aneur (DZ), 23.VI.1884.

Uromenus rugosicollis (Serville, 1838)

1 spec., Museum Paris, leg. Haury, 1907.

Steropleurus Bolívar, 1878

Steropleurus brunnerii (Bolívar, 1876-1878)

1 spec., Montarco Bei Madrid (Madrid, ES), leg.
Ebner, 16.VII.1924; det. Ebner, 1925; 1 spec.,

Montarco Bei Madrid (Madrid, ES), leg. Ebner, 16.VII.1924; det. Ebner, 1925.

Lluciapomaresius Barat, 2012

Lluciapomaresius stalii (Bolívar, 1878)

1 spec., Sierra de Guadarrama, Cercedilla, Kastilien (Madrid, ES), leg. Ebner, 18-24.VII.1924.

“Rolf Weyrauch” Collection

Bradyporus (Bradyporus) dasypus (Illiger, 1800)

2 specs., S.W. Dobr., V. Iortmac (Iortmac Lake, Lipnița, Constanța County, RO), leg. Weyrauch, 22/23.VI.1965; 1 spec., S.W. Dobr., V. Iortmac (Iortmac Lake, Lipnița, Constanța County, RO), leg. Weyrauch, 25.VI.1966; 1 spec., S.W. Dobr., V. Iortmac (Iortmac Lake, Lipnița, Constanța County, RO), leg. Weyrauch, 25.VI.1966.

Ephippiger ephippiger (Fiebig, 1784)

1 spec.; 1 spec., H.B. (Băile Herculane ?, Caraș-Severin County, RO), 15.VII.1957; 1 spec., Talmasch (Tălmaciu, Sibiu County, RO), 7.VII.1955; 1 spec., Rah. B. ?, 27.VII.1955.

“Eckbert Schneider” Collection

Bradyporus (Bradyporus) dasypus (Illiger, 1800)

3 specs., Dobrogea, Babadag (RO), leg. Schneider, 21.VI.1972.

Ephippiger ephippiger (Fiebig, 1784)

1 spec., Zakelsberg (Dealul Zakel), Sibiu County, RO), det. Worell, 1.IX.1954; 2 specs., Hammersdf. Bg b. Hermannstadt (Gușterița, Sibiu County, RO), leg. Schneider, 10.X.1969; 1 spec., Hammersdf. (Gușterița, Sibiu County, RO), leg. Schneider, 24.IX.1965; 1 spec., Tălmaciu, Deal. cu Pini (Tălmaciu, Sibiu County, RO), leg. Schneider, 21.IX.1975; 1 spec., Boița Wartberg (Boița, Sibiu County, RO), leg. Schneider, 26.IX.1967; 1 spec., Vânturarița (Vâlcea County, RO), leg. Schneider, 25. VIII.1964.

“Dr. Eugen Worell” Collection

Bradyporus (Bradyporus) dasypus (Illiger, 1800)

1 spec., Kaliakra, Bolatasteppe, Eichensteppe (Bolata, Kaliakra, BG), leg. Müller, 22.VII.1926; 1 spec., Caliacra, Bolatasteppe, Eichenm. (Bolata, Kaliakra, BG), leg. Müller; 1 spec., Donaudelta (Danube Delta, RO), leg. v. Spiess, det. Worell; 1 spec., Livezi, Doljiu, (Dolj County, RO), VIII.1925.

Comments: The last specimen is in fact a specimen of *Callimenus montandoni*, as the species is limited to Dobrogea (Iorgu 2009).

Ephippiger ephippiger (Fiebig, 1784)

2 specs., Michelsberg, Bei Hermannstadt (Cisnădioara, Sibiu County, RO), leg. Worell, 6.X.1941; 1 spec., Zakelsberg, Stolzburg (Dealul Zakel, Slimnic, Sibiu County, RO), leg. Worell, 26.VII.1947; 1 spec., Zakelsberg, Stolzenburg (Dealul Zakel, Slimnic, Sibiu County, RO), leg. Worell, 8.IX.1947; 1 spec., Götzenberg, Hermannstadt (Măgura Cisnădiei, Sibiu County, RO), leg. Worell, 18.VIII.1946; 1 spec., Hermannstadt, Hammersd., BG (Gușterița, Sibiu County, RO), leg. Worell, 16.VIII.1943; 1 spec., Hermannstadt, Hammersd., BG (Gușterița, Sibiu County, RO), leg. Worell, 17.VII.1945; 1 spec., Hermannstadt, Hammersd., BG (Gușterița, Sibiu County, RO), leg. Worell; 2 specs., Herkulesbad (Băile Herculane, Caraș-Severin County, RO), leg. Worell, 9.VI.1942; 2 specs., without other data.

Results and discussions

There are 12 species of armoured crickets in the entomological collections of the Natural History Museum of Sibiu, most of them preserved in the “Arnold Müller” collection: *Bradyporus (Bradyporus) dasypus* (Illiger, 1800), *Bradyporus (Callimenus) macrogaster longicollis* (Fieber, 1853), *Ephippiger ephippiger* (Fiebig, 1784) and *Bradyporus (Callimenus) montandoni* (Burr, 1898) which are present in the Romanian fauna (Iorgu *et al.* 2008) as well as eight additional foreign species: *Bradyporus (Callimenus) oniscus* (Burmeister, 1838), *Ephippiger diurnus cunii* Bolívar, 1877, *E. provincialis* Yersin, 1854, *E. discoidalis* Fieber, 1853, *Uromenus laticollis* (Lucas, 1846-1849), *U. rugosicollis* (Serville, 1839), *Steropleurus brunnerii* (Bolívar, 1876-1878) and *Lluciapomaresius stalii* (Bolívar, 1878).

Most of the specimens were collected from Romania, Bulgaria, the Czech Republic, Serbia, Spain, Greece and Algeria; however, others were obtained through exchanges with other collectors or museums (e.g. *Ephippiger provincialis* and *Uromenus rugosicollis* with the Museum of Paris or *Steropleurus brunnerii* and *Lluciapomaresius stalii* with Ebner, all of them being part of the “Arnold Müller” collection).

The subfamily *Bradyporinae* is systematically difficult and re-identification of all the specimens should be undertaken. However, of the ones listed

above, eight were definitely inaccurately identified. In the “Arnold Müller” collection, the three *Callimenus oniscus* specimens collected in Ploșor and Greci are actually *Callimenus montandoni*, the former taxon not being present in the Romanian fauna (Iorgu *et al.* 2008). The situation is identical with one of the *Bradyporus dasypus* specimens from the “Eugen Worell” collection, which was found in Livezi, in Romania this species being limited to Dobrogea (Iorgu 2009). The presence of one specimen of *Bradyporus dasypus* in the Danube Delta (also from the “Eugen Worell” collection) is another interesting finding, unknown so far. It is not specified on the label whether the specimen was

found strictly in the delta or outside of it, in other parts of the reserve such as the Razelm-Sinoe lakes where the species may be present. There is also the issue of the four specimens of *Ephippiger diurnus cunii* which are not present in the Bulgarian fauna. They could not be reidentified.

The collecting sites for all the specimens are displayed in Fig. 2 to 8.

Acknowledgements

The authors are grateful for the comments of Ionuț Ștefan Iorgu and Gabriel Lupu, which significantly improved the first version of the manuscript.

REFERENCES

- Harz 1969 Harz Kurt, *Die Orthopteren Europas/The Orthoptera of Europe I*. In: *Series Entomologica* V, (1969).
- Iorgu *et al.* 2008 Iorgu Ionuț, Pisciă Elena, Păiș Laura, Lupu Gabriel, Iușan Claudiu, *Checklist of Romanian Orthoptera (Insecta) and their distribution by eco-regions*. In: *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"* 51, Bucharest (2008), p. 119-135.
- Iorgu, Iorgu 2008 Iorgu Ionuț Ștefan, Iorgu Elena Iulia, *Bush-crickets, crickets and grasshoppers from Moldavia (Romania)*. In: *Editura Pim, Iași* (2008).
- Iorgu 2009 Iorgu Ionuț Ștefan, *Bradyporus dasypus (Illiger, 1800) (Orthoptera: Tettigoniidae): Some ethological aspects and distribution in Romania*. In: *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"* 52, Bucharest (2009), p. 143-149.
- Kis 1962 Kis Béla, *Revision der in Rumänien vorkommenden Bradyporinae - Arten (Orth.)*. In: *Sonderabdruck aus den Mitteilungen der Münchner Entomologischen Gesellschaft* 5 (1962), p. 115-122.
- Pascu, Schneider 1998 Pascu Mariana, Schneider Eckbert, *Colecțiile entomologice ale Muzeului de Istorie Naturală din Sibiu*. In: *Stud. și Com. Șt. Nat. Muz. Brukenthal* 27, Sibiu (1998), p. 201-218.
- Vasiliu 1961 Vasiliu Maria, *Contribuții la cunoașterea speciilor de Bradyporus Charp. (Orthoptera, Tettigoniidae) din R. P. R.* In: *Studii și Cercetări de Biologie, seria Biologie Animală (extras)* 13 (2), București (1961), p. 203-212.
- Vasiliu, Agapi 1958 Vasiliu Maria, Agapi Constantin, *Catalogul colecției de Orthoptere „Arnold Muller” de la Muzeul Brukenthal din Sibiu*. In: *Studii și Comunicări, Muzeul Brukenthal* 10, Sibiu (1958).
- Eades *et al.* 2012 Eades David, Otte Daniel, Cigliano Maria Marta, Braun Holger, *Orthoptera Species File Online*. Version 2.0/4.1. [21.05.2012]. <<http://Orthoptera.SpeciesFile.org>>
- Heller 2012 Heller Klaus-Gerhard, *Fauna Europaea: Orthoptera. Fauna Europaea version 2.4*. In: <http://www.faunaeur.org> (2012)

LIST OF ILLUSTRATIONS

- Fig. 1.** Distribution of armoured crickets in Europe, according to <http://faunaeur.org>
- Fig. 2.** Distribution of *Bradyporus (Bradyporus) dasypus* (Illiger, 1800) based on museum collections
- Fig. 3.** Distribution of *Bradyporus (Callimenus) macrogaster longicollis* (Fieber, 1853), based on museum collections
- Fig. 4.** Distribution of *Bradyporus (Callimenus) montandoni* (Burr, 1898), based on museum collections
- Fig. 5.** Distribution of *Bradyporus (Callimenus) oniscus* (Burmeister, 1838), based on museum collections
- Fig. 6.** Distribution of *Ephippiger diurnus cunii* Bolívar, 1877, *Ephippiger provincialis* Yersin, 1854 and *Ephippiger discoidalis* Fieber, 1853, based on museum collections
- Fig. 7.** Distribution of *Ephippiger ephippiger* (Fiebig, 1784), based on museum collections
- Fig. 8.** Distribution of *Uromenus laticollis* (Lucas, 1846-1849), *Steropleurus brunnerii* (Bolívar 1876-1878) and *Lluciapomaresius stalii* (Bolívar 1878)
- Fig. 9.** a) *Bradyporus (Bradyporus) dasypus* (Illiger, 1800); b) *Ephippiger ephippiger* (Fiebig, 1784); c) *Ephippiger diurnus cunii* Bolívar, 1877 ; d) *Ephippiger provincialis* Yersin, 1854; e) *Ephippiger discoidalis* Fieber, 1853; f) *Bradyporus (Callimenus) montandoni* (Burr, 1898), specimens from the “Arnold Müller” collection
- Fig. 10.** a) *Bradyporus (Callimenus) macrogaster longicollis* (Fieber, 1853); b) *Bradyporus (Callimenus) oniscus* (Burmeister, 1838); c) *Uromenus laticollis* (Lucas, 1846-1849); d) *Uromenus rugosicollis* (Serville, 1838); e) *Steropleurus brunnerii* (Bolívar, 1876-1878); f) *Lluciapomaresius stalii* (Bolívar 1878), specimens from the “Arnold Müller” collection

LISTA ILUSTRAȚIILOR

- Fig. 1.** Distribuția bradiporinelor în Europa potrivit <http://faunaeur.org>
- Fig. 2.** Distribuția lui *Bradyporus (Bradyporus) dasypus* (Illiger, 1800), bazată pe colecțiile muzeului
- Fig. 3.** Distribuția lui *Bradyporus (Callimenus) macrogaster longicollis* (Fieber, 1853), bazată pe colecțiile muzeului
- Fig. 4.** Distribuția lui *Bradyporus (Callimenus) montandoni* (Burr, 1898), bazată pe colecțiile muzeului
- Fig. 5.** Distribuția lui *Bradyporus (Callimenus) oniscus* (Burmeister, 1838), bazată pe colecțiile muzeului
- Fig. 6.** Distribuțiile lui *Ephippiger diurnus cunii* Bolívar, 1877, *Ephippiger provincialis* Yersin, 1854 și *Ephippiger discoidalis* Fieber, 1853, bazate pe colecțiile muzeului
- Fig. 7.** Distribuția lui *Ephippiger ephippiger* (Fiebig, 1784), bazată pe colecțiile muzeului
- Fig. 8.** Distribuțiile lui *Uromenus laticollis* (Lucas, 1846-1849), *Steropleurus brunnerii* (Bolívar 1876-1878) și *Lluciapomaresius stalii* (Bolívar 1878), bazate pe colecțiile muzeului
- Fig. 9.** a) *Bradyporus (Bradyporus) dasypus* (Illiger, 1800); b) *Ephippiger ephippiger* (Fiebig, 1784); c) *Ephippiger diurnus cunii* Bolívar, 1877 ; d) *Ephippiger provincialis* Yersin, 1854; e) *Ephippiger discoidalis* Fieber, 1853; f) *Bradyporus (Callimenus) montandoni* (Burr, 1898), specimene din colecția “Arnold Müller”
- Fig. 10.** a) *Bradyporus (Callimenus) macrogaster longicollis* (Fieber, 1853); b) *Bradyporus (Callimenus) oniscus* (Burmeister, 1838); c) *Uromenus laticollis* (Lucas, 1846-1849); d) *Uromenus rugosicollis* (Serville, 1838); e) *Steropleurus brunnerii* (Bolívar, 1876-1878); f) *Lluciapomaresius stalii* (Bolívar 1878), specimene din colecția “Arnold Müller”




Fig. 1. Distribution of armoured crickets in Europe, according to Heller (2012)




Fig. 2. Distribution of *Bradyporus (Bradyporus) dasypus* (Illiger, 1800) based on museum collections



Fig. 3. Distribution of *Bradyporus (Callimenus) macrogaster longicollis* (Fieber, 1853), based on museum collections




Fig. 4. Distribution of *Bradyporus (Callimenus) montandoni* (Burr, 1898), based on museum collections



Fig. 5. Distribution of *Bradyporus (Callimenus) oniscus* (Burmeister, 1838), based on museum collection




Fig. 6. Distribution of *Ehippiger diurnus cunii* Bolívar, 1877, *Ehippiger provincialis* Yersin, 1854 and *Ehippiger discoidalis* Fieber, 1853, based on museum collection




Fig. 7. Distribution of *Ehippiger ehippiger* (Fiebig, 1784), based on museum collections




Fig. 8. Distribution of *Uromenus laticollis* (Lucas, 1846-1849), *Steropleurus brunnerii* (Bolívar 1876-1878), *Lluiciapomaresius stalii* (Bolívar, 1877), based on museum collection




Fig. 9. a) *Bradyporus dasypus* (Illiger, 1800); b) *Ephippiger ephippiger* (Fiebig, 1784); c) *Ephippiger diurnus cunii* Bolívar, 1877 ; d) *Ephippiger provincialis* Yersin, 1854; e) *Ephippiger discoidalis* Fieber, 1853; f) *Bradyporus (Callimenus) montandoni* (Burr, 1898), specimens from the “Arnold Müller“ collection




Fig. 10. a) *Bradyporus (Callimenus) macrogaster longicollis* (Fieber, 1853); b) *Bradyporus (Callimenus) oniscus* (Burmeister, 1838); c) *Uromenus laticollis* (Lucas, 1846-1849); d) *Uromenus rugosicollis* (Serville, 1838); e) *Steropleurus brunnerii* (Bolívar, 1876-1878); f) *Lluciapomaresius stalii* (Bolívar, 1878), specimens from the “Arnold Müller” collection

**ANT FAUNA (HYMENOPTERA: FORMICIDAE) OF SIBIU COUNTY
(TRANSYLVANIA, ROMANIA)**

Ioan TĂUȘAN*
Mădălina Maria JERPEL**
Ioana Roxana PUȘCAȘU**
Cătălin SĂDEANU**
Roberta Elena BRUTARU**
Lavinia Ana RĂDUȚIU**
Valeria GIURESCU**

Abstract. *In the present paper, the authors give a first checklist of the ant species from Sibiu County. The data was compiled from published literature, unpublished data from museum collections and personal collections. Altogether 70 species were recorded. Tapinoma subboreale Seifert, 2012 is new for the ant fauna of Sibiu. Distribution maps are given for all species.*

Keywords: *ants, checklist, new records, distribution, Sibiu.*

Rezumat. *În prezenta lucrare, autorii oferă prima listă a furnicilor din județul Sibiu. Datele au fost compilate, din literatura existentă, date nepublicate din colecții muzeale și din colectări personale. În total au fost identificate 70 de specii. Tapinoma subboreale Seifert, 2012 este nouă pentru fauna județului. Adicional, sunt prezentate hărți ale distribuției speciilor.*

Cuvinte cheie: *furnici, listă faunistică, noi semnalări, distribuție, Sibiu*

Introduction

The ant fauna of Romania has been studied in the last decade (Markó *et al.* 2006) and 109 ant species are now known from Romania (Ionescu-Hirsch *et al.* 2009, Markó *et al.* 2006, Markó 2008a, b, Moscaliuc 2009, Czekes *et al.* 2012). Comparatively with other neighbouring countries, the number appears low: Hungary – 125 species (Csősz *et al.* 2011), Bulgaria – 163 species (Lapeva-Gjonova *et al.* 2010), Montenegro – 135 (Karaman 2004, 2008), Serbia – 141 (Petrov 2006). Due to intensive myrmecological investigations the number is expected to grow.

The entomofauna of Sibiu was studied in particular, due to the scientific activity of the Transylvanian Natural History Society of Sibiu (Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt – 1850 – 1946). The bases of the Transylvanian Society Collection of Natural Sciences were established by Karl Fuss, considered to be the father of entomology in Transylvania (Pascu, Schneider 1998).

The entomological collections, of great scientific value, preserved in the Natural History Museum of Sibiu proves that important steps were made, regarding the research in the Romanian entomology field (Bucșa, Tăușan 2011).

Despite that the ant fauna of Sibiu County was rather well studied (Paraschivescu 1975, Markó, Csősz 2002, Csősz, Markó 2005, Tăușan, Markó 2009, Tăușan *et al.* 2011, Tăușan *et al.* 2012) no ant check list is available.

This study renders the first ant checklist with sampling data and the distribution map of the species recorded in Sibiu County.

Study area, material and methods

Having a surface of 5575 km², Sibiu County is situated in the central part of Romania. The geographic position of the County at the limit between the Southern Carpathians (which occupies the southern part of the county) and the Transylvanian Depression (which occupies the central and northern part) and its large extension in the two tectonic-structural units give the aspect of an amphitheatre opened northwards (Fig. 1). The landform's evolution is subdued to the Carpathian chain represented on the county's territory by

* Babeș-Bolyai University of Cluj-Napoca and Brukenthal National Museum, Natural History Museum, itausan@gmail.com, ioan.tausan@brukenthalmuseum.ro

** Lucian Blaga University of Sibiu, Faculty of Sciences,

Făgăraș, Cindrel and Lotru Mountains (Costea 2011).

Using published and unpublished records, we have compiled the ant fauna of Sibiu County. In addition we used unpublished data from the Natural History Museum of Sibiu collections.

The ant material of the Natural History Museum of Sibiu, consisting of 1577 specimens (Markó, Csősz 2002; Csősz, Markó 2005), is made up of two separate collections: that of the former Transylvanian Natural History Society of Sibiu (Siebenbürgisches Verein für Naturwissenschaften zu Hermannstadt) mostly arranged by Arnold Müller and containing Paul Rösler's types, and Dr. Eugen Worrell's personal collection, which was acquired by the Museum in 1957 (Paraschivescu 1975). Additional, the museum hosts also Eckbert Schneider's collection, 330 specimens, almost all of the specimens being unidentified.

The identification of ant species was carried out on the basis of several available identification keys (Czechowski *et al.* 2002, Seifert 2007, Markó *et al.* 2009). The following codes are used in the paper: w – worker; m – male; q – queen; coll. Schneider – Dr. Eckbert Schneider's collection, coll. Tăușan – Ioan Tăușan's collection; in square brackets the Romanian name of the sampling place.

List of species

Family Formicidae

Subfamily Dolichoderinae

Tribe Dolichoderini

Dolichoderus quadripunctatus (Linnaeus, 1767) (Fig. 2)

Published records: Cisnădie, Sibiu, (Markó, Csősz 2002).

New records: 1 w, Transsilv. merid. Hamba, Sibiu, 4-16.08.1973, leg. E. Schneider, coll. E. Schneider.

Tapinoma erraticum (Latreille, 1798) (Fig. 2)

Published records: Dealul Cetății, Tălmăciu, Gușterița, Cisnădioara, Petiș, Cisnădie (Markó, Csősz 2002).

New records: 2 w, Transsilv. merid. Șura Mare, 23.05.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012.

Tapinoma subboreale Seifert, 2012 (Fig. 2)

New records: 3 w, Ocna Sibiului, 07.2009, leg. Tăușan *et al.* coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan *et al.*, coll. Tăușan.

Subfamily Formicinae

Tribe Camponotini

Camponotus aethiops (Latreille, 1798) (Fig. 3)

Published records: Sibiu, Gușterița (Csősz, Markó 2005).

New records: 2 w, Șura Mare.7.8.1971 leg. E. Schnieder, as [*Camponotus silvaticus* Ol.], det. Tăușan *et al.* 2012; 1w, Transsilv. merid., Șura Mare, 23.05.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012.

Camponotus fallax (Nylander 1856) (Fig. 3)

Published records: Sibiu (Csősz, Markó 2005).

New records: 1 w, the garden of NHMS, 06.06.2012, leg. Brutaru & Tăușan, coll. Tăușan.

Camponotus herculeanus (Linnaeus, 1758) (Fig. 4)

Published records: Păltiniș, Cindrel Mountains, Prislop-Rășinari (Csősz, Markó 2005).

New records: 1 w, Mts. Cibin, 1300-1400 m a.s.l., 11.08.1957, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 1 w, V. Sebeșului, 27.08.1966, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012.

Camponotus ligniperdus (Latreille, 1802) (Fig. 4)

Published records: Păltiniș, Cisnădioara, Măgura Cisnădioarei, Cisnădioara, Sibiu, Gușterița, Rîu Sadu (Csősz, Markó 2005); Cisnădioara (Tăușan *et al.* 2012).

New records: 1 q, Poplaca b. Hermannstadt, 11.09.1969, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 4 w, Hammersdorf [Gușterița], 25.06.1970, leg. E. Schneider; 1 w Transsilv. merid. Șura Mare, 20.04.1973, leg. E. Schneider coll. E. Schneider, det. Tăușan *et al.* 2012; 1 w Transsilv. merid. Șura Mare, 13.06.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 3 w, Hammersdorf Bg. [Dealul Gușteriței], 22.09.1974, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 4 w, Transsilv. merid. Cisnădioara, 3.08.1975, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 1 w Transsilv. merid. Podu Olt, 7.05.1977, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 1 w, Mt. Cibin, Valea Sadului, 8.08.1980, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 2 w, Podu Olt, 18.05.1984, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 1 w, Racovița, 06.06.2009, leg. G. Mărginean, det. Tăușan *et al.* 2012, coll. Tăușan.

Camponotus piceus (Leach, 1825) (Fig. 5)

Published records: Șura Mare, Gușterița (Csősz, Markó 2005).

New records: 1 w, Sibiu, Veștem, 3.10.1969, b. Hermannstadt [Sibiu], leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1w, Hammersdorf Tal. [Gușterița], 19.06.1970, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Transsilv. merid. Șura Mare, 23.05.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1w, Transsilv. merid., Ocna Sibiului, 13.07.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 2 w, Transsilv. merid. Șura Mare, 20.07.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 2 w, Transsilv. merid., Ocna Sibiului, 19.05.1974, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Dealul Gușteriței, 4.10.1974, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 q, 1 w, Transsilv. merid. Ocna Sibiului, 30.05.1976, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 2 w, Racovița, 26.09.2010, leg. G. Mărginean, det. Tăușan et al. 2012, coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan.

Camponotus truncatus (Spinola, 1808) (Fig. 3)

Published records: Sibiu, Cîsnădie (Csősz, Markó 2005).

Camponotus vagus (Scopoli, 1763) (Fig. 5)

Published records: Sibiu, Cîsnădie, Dealul Cetății, Tălmăciu, Cîsnădioara, Măgura Cîsnădioarei, Cîsnădioara (Csősz, Markó 2005).

New records: 1 w, Transsilv. merid. Șura Mare, 28.04.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1w, Transsilv. merid. Șura Mare, 23.05.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 2 w, Porumbacu de Sus, Carpat. Merid. 8.07.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 2 w, Transilv. Sibiu, D. Gușterița, 18.05.1975, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012.

Tribe Formicini

Formica cinerea (Mayr, 1853) (Fig. 6)

Published records: Gușterița/Szenterzsebet, Sibiu, 28.10.1918, leg. Müller, coll. NHMS; 6 w, Hammersdorf [Gușterița], 05.1910, leg. Anonymous, coll. NHMS (Csősz, Markó 2005); Dumbrava (Tăușan, Markó 2009); Cîsnădioara (Tăușan et al. 2012).

New records: 1 w, V. Sadu, Cîinăia, 1700 m a.s.l., 3.10.1971, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 3 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan; 11 w Sadu valley, 06. 2010 leg. Tăușan et al. coll. Tăușan, 6 w Racovița, 31.10.2010 leg. G. Mărginean, det.

Tăușan et al. coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan.

Formica clara Forel, 1886 syn. *Formica lusatica* Seifert, 1997 (Seifert, Schultz 2009) (Fig. 6)

Published records: as *Formica lusatica* Seifert, 1997, Gușterița (Tăușan, Markó 2009).

New records: 2 w, Sibiu, 4.04.1970, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 2 w, Transsilv. merid. Șura Mare, 23.05.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan.

Formica cunicularia (Latreille, 1798) (Fig. 6)

Published records: Sibiu, Gușterița, Bungard (Csősz, Markó 2005); Cîsnădioara (Tăușan, Markó 2009; Tăușan et al. 2012).

New records: 1 w, Dealul Gușteriței, Sibiu, 04.04.1971, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Sadu valley, 06. 2010 leg. Tăușan et al. coll. Tăușan, 2 w, 14.10.2010, Racovița, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al., coll. Tăușan.

Formica fusca (Linnaeus, 1758) (Fig. 6)

Published records: Păltiniș, Măgura Cîsnădioarei, (Csősz, Markó 2005).

New records: 3 w, V. Sadu, Cîinăia, 1700 m a.s.l., 3.10.1971, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Șura Mare, 30.04.1972, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Transsilv. merid. Șura Mare, 28.04.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Transsilv. merid. Șura Mare, 23.05.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Munții Cibin, R. Sadului Val. Vîrjoghii (Cibin Mountains), 28.09.1975, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Sadu valley, 06. 2010 leg. Tăușan et al., coll. Tăușan, 1 w, 2010 leg. G. Mărginean, det. Tăușan et al. coll. Tăușan ; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al., coll. Tăușan.

Formica gagates (Latreille, 1798) (Fig. 7)

Published records: Gușterița, Agârbiciu (Csősz, Markó 2005).

Formica lemani (Bondroit, 1917) (Fig. 6)

Published records: Sibiu, Gușterița, Păltiniș, Cîndrel (Csősz, Markó 2005).

New records: 5 w, V. Sadu, Cîinăia, 1700 m a.s.l., 3.10.1971, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w Sadu valley, 06. 2010

leg. Tăușan et al. coll. Tăușan; 1 w Racovița, 21.08.2010, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al., coll. Tăușan.

Formica polyctena (Forster, 1850) (Fig. 8)

Published records: Sibiu, Rășinari, Măgura Cisnădioarei (Csősz, Markó 2005).

New records: 2 w Racovița, 31.10.2010 leg. G. Mărginean, det. Tăușan et al., coll. Tăușan.

Formica pratensis (Retzius, 1783) (Fig. 8)

Published records: Dumbrava Sibiului, Sibiu, Dealul Cetății-Tălmăciu, Turnu Roșu, Măgura Cisnădioarei, Cristian (Csősz, Markó 2005).

As *Formica nigricans* Emery, 1909, Măgura Cisnădioarei, Cisnădioara, Cindrel, Gușterița in (Csősz, Markó 2005).

New records: 5 w, Șura Mare, 10.10.1971, leg. E. Schneider, coll. E. Schneider; 1 q, Șura Mare, 24.8.1972, leg. E. Schneider, coll. Schneider; 1 q, Transsilv. Merid. Șura Mare, 20.07.1973, leg. E. Schneider, coll. E. Schneider; 1 w Sadu valley, 06. 2010 leg. Tăușan et al. coll. Tăușan.

Formica rufa (Linnaeus, 1758) (Fig. 7)

Published records: Cristian (Csősz, Markó 2005).

Formica rufibarbis (Fabricius, 1793)

Published records: Cisnădie (Csősz, Markó 2005); Cisnădioara, Gușterița (Tăușan, Markó, 2009).

New records: 2 w, Hermmannstadt Reussbachwiesen, 26.04.1969, leg. E. Schneider, Coll. E. Schneider; 1 w, Slimnic, D. Zakel, 24.08.1972, leg. E. Schneider, coll. E.Schneider, det. Tăușan et al. 2012; 1 w, Trassilv. merid., Ocna Sibiului, 19.05.1974, leg. E. Schneider, coll. E.Schneider, det. Tăușan et al. 2012; 2 w, Transilvania merid. Șura Mare, 23.5.1975, leg. E. Schneider, Coll. E. Schneider, det. Tăușan et al. 2012; 3 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan; 1 w, Sadu valley, 06. 2010 leg. Tăușan et al. coll. Tăușan; 3 w, Racovița 26.09.2010 leg. G. Mărginean, det. Tăușan et al. coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan;

Formica sanguinea (Latreille, 1798) (Fig. 7)

Published records: Gușterița, Dealul Cetății-Tălmăciu (Csősz, Markó 2005).

New records: 4 w, V. Sadu (Cânaia), 1700 m a.s.l., 3.10.1971, leg. E. Schneider, coll. E. Schneider; 3 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan; 1 w, Sadu valley, 06. 2010, leg. Tăușan et al. coll. Tăușan.

Polyergus rufescens (Latreille, 1798) (Fig. 7)

Published records: Bungard, Sibiu (Csősz, Markó 2005).

New records: 2 w, Ocna Sibiului, 25.07.1971, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 3 w, Dumbrăveni, 07.2011, leg. Tăușan et al. coll. Tăușan; 3 w, Pădurea Dumbrava, 9.07.2012, leg. Tăușan, coll. Tăușan.

Tribe Lasiini

Lasius alienus (Förster, 1850) (Fig. 9)

Published records: Cisnădie (Csősz, Markó 2005).

New records: 1 w Racovița, 21.08.2010, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan.

Lasius bicornis (Förster, 1850) (Fig. 9)

Published records: Viile Sibiului Sibiu (Csősz, Markó 2005).

Lasius brunneus (Latreille, 1798) (Fig. 10)

Published records: Ocna Sibiului, Sibiu, Rîu Sadu Măgura Cisnădioarei, Dumbrava Sibiului, Turnu Roșu, Cisnădie, (Csősz, Markó 2005); Cisnădioara (Tăușan, Markó 2009).

New records: 1 w, Dealul Gușteriței (Gușterița Hill), 28.07.1971, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 4 w Sadu valley, 06. 2010 leg. Tăușan et al. coll. Tăușan, 6 w, Racovița, 2010, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan.

Lasius citrinus Emery, 1922 (Fig. 9)

Published records: as *Lasius afinis* Sibiu (Csősz, Markó 2005).

Lasius distinguendus (Emery, 1916) (Fig. 9)

Published records: Cindrel (Csősz, Markó 2005).

Lasius emarginatus (Olivier, 1922) (Fig. 9)

Published records: Sibiu (Csősz, Markó 2005).

New records: 5 w Racovița, 06.10.2010, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan ; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan.

Lasius flavus (Fabricius, 1781) (Fig. 10)

Published records: Păliniș, Cindrel, Cisnădie, [Dealul Cetății-Tălmăciu, Cisnădie (Csősz, Markó 2005); Cisnădioara (Tăușan et al. 2012).

New records: 4 w, Sibiu, 11. 04.1971, Dumbrava, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Munții Cibin, Rășinari Buru, 3.05.1975, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 6 w, Cornățel, Cașolț,

24.04.1983, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 3 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan.

Lasius fuliginosus (Latreille, 1798) (Fig. 10)

Published records: Sibiu, Gușterița, Slimnic, Turnu Roșu, Cisnădioara (Csősz, Markó 2005); Cisnădioara, Gușterița (Tăușan, Markó, 2009).

New records: 4 w, Sibiu, 28.03.1971, Dealul Gușteriței, leg. E. Schneider, coll. E. Schneider; 3 w, Racovița, 31.10.2010, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan.

Lasius niger (Linnaeus, 1758) (Fig. 11)

Published records: Gușterița, Sibiu, Cisnădie, Tălmăciu, Dumbrava Sibiului (Csősz, Markó 2005); Cisnădioara (Tăușan, Markó 2009; Tăușan et al. 2012); Gușterița, Seviș (Tăușan, Markó 2009).

New records: 3 w, Hermannstadt, Reussbachwiesen, 26.04.1969, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 2 w, Transsilv. Merid. 23.05.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 2 w, Pasul Turnu Roșu Lotrioara, 24.03.1974, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 4 w, Cornățel, Cașolt, 24.04.1983, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 3 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan; 15 w, Racovița 06.2009 leg. Tăușan et al. coll. Tăușan; 10 w, Sadu valley, 06.2010 leg. Tăușan et al. coll. Tăușan, 3 w Racovița, 06.10.2010, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan.

Lasius paralienus (Seifert, 1992) (Fig. 11)

Published records: Sibiu, Gușterița, Turnișor (Csősz, Markó 2005); Cisnădioara (Tăușan et al. 2012).

New records: 1 w, Hammersdorf, Sibiu [Gușterița, Sibiu], 19.06.1970, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan; 3 w, 21.08.2010, Racovița, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan.

Lasius platythorax (Seifert, 1922) (Fig. 11)

Published records: Cisnădie, (Csősz, Markó 2005).

New records: 1 w, Racovița, leg. G. Mărginean, det. Tăușan et al., coll. Tăușan; 2 w, Sibiu, Hammersdorf Tal. [Dealul Gușterița], 19.06.1970, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 3 w, Ocna Sibiului, 07.2009, leg. Tăușan

et al. coll. Tăușan; 06.2009; 4 w Sadu valley, 06.2010 leg. Tăușan et al. coll. Tăușan.

Lasius umbratus (Nylander, 1846)

Published records: Păltiniș (Csősz, Markó 2005).

Prenolepis nitens (Mayr, 1853) (Fig. 12)

Published records: Viile Sibiului, Cristian (Csősz, Markó 2005).

Tribe Plagiolepidini

Plagiolepis pygmaea (Latreille, 1798) (Fig. 12)

Published records: Sibiu (Csősz, Markó 2005).

New records: 1 w, Sibiu, Hammersdorf Tal. [Dealul Gușteriței], 19.06.1970 leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Șura Mare, 10.10.1971, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 2 w, Transsilv. merid., Șura Mare, 23.05.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Trassilv. merid., Ocna Sibiului, 13.07.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012.

Plagiolepis taurica Santschi, 1920 – Senior synonym of *P. vindobonnensis* Lomnicki, 1925 (Radchenko, 1989) (Fig. 12)

Published records: Gușterița, Cisnădie (Csősz, Markó 2005).

Subfamily Myrmicinae Lepeletier, 1836

Tribe Crematogastrini

Crematogaster schmidtii (Mayr, 1852) (Fig. 13)

Published records: Păltiniș (Markó, Csősz 2002).

Tribe Formicoxenini

Formicoxenus nitidulus (Nylander, 1846) (Fig. 14)

Published records: Dealul Cetății-Tălmăciu, Sibiu (Markó, Csősz 2002).

Leptothorax acervorum (Fabricius, 1793) (Fig. 14)

Published records: Măgura Cisnădiei (Markó, Csősz 2002).

New records: 1 w, Șura Mare, Sibiu, 10.10.1971, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012.

Temnothorax affinis (Mayr, 1855) (Fig. 14)

Published records: as *Leptothorax affinis*, Gușterița, Cisnădie (Markó, Csősz 2002).

Temnothorax corticalis (Schenck, 1852) (Fig. 14)

Published records: as *Leptothorax corticalis* Sibiu (Markó, Csősz 2002).

Temnothorax clypeatus (Mayr, 1853) (Fig. 14)

Published records: as *Leptothorax clypeatus*, Cîsnădie (Markó, Csősz 2002).

Temnothorax crassispinus (Karavajev, 1926) (Fig. 14)

Published records: as *Leptothorax crassispinus*, Cîsnădie (Markó, Csősz 2002); as *Temnothorax crassispinus*, Cîsnădioara (Tăușan *et al.* 2012).

New records: 1 w, Transsilv. merid. Șura Mare, 23.05.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 3 w, Dumbrăveni 07.2011, leg. Tăușan *et al.* coll. Tăușan

Temnothorax unifasciatus (Latreille, 1798) (Fig. 14)

Published records: as *Leptothorax unifasciatus*, Cîsnădie, as *Leptothorax tuberosum* var. *interrupta* det. Müller, Cîsnădie (Markó, Csősz 2002).

Tribe Myrmicini

Manica rubida (Latreille, 1802) (Fig. 15)

Published records: Păltiniș, Munții Cîbinului (Markó, Csősz 2002).

New records: 3 q, 4 m and 9 w, V. Sadu, Cînaia, 1700 m a.s.l., 3.10.1971, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012.

Myrmica constricta Karawajew, 1934 (Fig. 18)

Published records: Gușterița (Markó, Csősz 2002); Media (Seifert *et al.* 2009).

Myrmica gallienii Bondroit, 1920 (Fig. 18)

Published records: Ocna Sibiului (Czekes *et al.* 2012).

Myrmica lobicornis Nylander, 1846 (Fig. 15)

Published records: Cîbin Mts. (Paraschivescu 1975).

New records: 1 w, Sadu Cînaia, 11.09.1969; 2 w, Sadu valley, 06. 2010 leg. Tăușan *et al.* coll. Tăușan; 1 w Racovița, 31.10.2010, leg. G. Mărginean, det. Tăușan, coll. Tăușan.

Myrmica lonae Finzi, 1926 (Fig. 18)

Published records: Erbota Valley (Făgăraș Mts.) (Markó, Csősz 2001).

Myrmica rubra (Linnaeus, 1758) (Fig. 16)

Published records: Sibiu, Păltiniș, Sibiu (Markó, Csősz 2002); Dumbrava Sibiului (Tăușan, Markó 2009).

New records: 1 q and 1 m, Dl., 28.07.1971, leg. E. Schneider, coll. E. Schneider; 2 w, Mts. Cîbin, Michelsberg (Cîsnădioara), 1.05.1973, leg. E.

Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 1 w, Transsilv. Merid. Șura Mare, 23.05.1973, leg. E. Schneider, det. as [*M. rugulosa*]; 4 w, Pasul Turnu Roșu Lotrioara, 24.03.1974, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 1w, 2 q, Transsilv. merid. Cîsnădioara, 3.08.1975, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 9 w, Sadu valley, 06. 2010 leg. Tăușan *et al.* coll. Tăușan; 9 w Racovița 2010, leg. G. Mărginean, det. Tăușan *et al.* coll. Tăușan.

Myrmica ruginodis Nylander, 1846 (Fig. 16)

Published records: Măgura Cîsnădiei, Cîsnădie, Dealul Cetății-Tălmăciu, Sibiu (Markó, Csősz 2002); Dumbrava Sibiului (Tăușan, Markó, 2009), Cîsnădioara (Tăușan *et al.* 2012).

New records: 1 w, Poplaca, 11.09.1969, leg. E. Schneider, coll. E. Schneider; 1 w Sibiu, Hammersdorf [Sibiu, Gușterița], 23.06.1970, leg. E. Schneider, coll. E. Schneider; 3 w, Ocna Sibiului, 07.2009, leg. Tăușan *et al.* coll. Tăușan; 3 w, Sadu valley, 06. 2010 leg. Tăușan *et al.* coll. Tăușan, 1 w 2010 Racovița, leg. G. Mărginean, det. Tăușan *et al.* coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan *et al.* coll. Tăușan.

Myrmica rugulosa Nylander, 1849 (Fig. 15)

Published records: Sibiu, Gușterița (Markó, Csősz 2002).

New records: 1 w, Poplaca b. Hermannstadt [Poplaca, Sibiu], 11.09.1969, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012.

Myrmica sabuleti Meinert, 1860 (Fig. 17)

Published records: Dealul Cetății-Tălmăciu, Sibiu, Gușterița (Markó, Csősz 2002).

New records: 3 w, Ocna Sibiului, 07.2009, leg. Tăușan *et al.* coll. Tăușan; 1 w, Sadu valley, 06. 2010 leg. Tăușan *et al.* coll. Tăușan, 2 w Racovița, 26.09.2010 leg. G. Mărginean, det. Tăușan *et al.* coll. Tăușan; 1 w, Transsilv. Merid. Șura Mare, 23.05.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012.

Myrmica scabrinodis Nylander, 1846 (Fig. 17)

Published records: Sibiu, Gușterița, Păltiniș, (Markó, Csősz 2002); Cîsnădioara (Tăușan *et al.* 2012)

New records: 1 m, Hermannstadt [Sibiu], 27.09.1969, leg. E. Schneider, coll. E. Schneider; 1 q, Hermannstadt, Salzburger Bg., 27.09.1969, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.* 2012; 4 w, Sibiu, 11. 04.1971, Dumbrava, leg. E. Schneider, coll. E. Schneider, det. Tăușan *et al.*

2012; 3 w, V. Sadu, Cînaia, 1700 m a.s.l., 3.10.1971, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 2 w, Trassilv. merid., Ocna Sibiului, 19.05.1974, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 3 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan; 1 w, Sadu valley, 06. 2010 leg. Tăușan et al. coll. Tăușan, 5 w Racovița, 21.08.2010, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan;

Myrmica schenki Emery, 1894 (Fig. 15)

Published records: Gușterița, Sibiu, Dumbrava Sibiului (Markó, Csősz 2002).

New records: 3 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan; 1 w, Racovița, 31.10.2009, leg. G. Mărginean, det. Tăușan et al. 2012, coll. Tăușan; 2 w, Sadu valley, 06. 2010 leg. Tăușan et al. coll. Tăușan;

Myrmica slovac Sadi, 1952 (Fig. 15)

Published records: as *Myrmica salina*, Mediaș, Rîu Sadu (Markó, Csősz 2002).

Myrmica specioidea Bondroit, 1918 (Fig. 15)

Published records: Sibiu, (Markó, Csősz 2002); Cîsnădioara (Tăușan et al. 2012)

New records: 2 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan.

Myrmica sulcinodis Nylander, 1846 (Fig. 18)

Published records: Cibin Mts. (Paraschivescu, 1975).

Tribe Myrmecini

Myrmecina graminicola (Latreille, 1802) (Fig. 13)

Published records: Sibiu, Dealul Zakei-Slimnic, (Markó, Csősz 2002); Cîsnădioara (Tăușan et al. 2012).

New records: 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan.

Tribe Pheidolini

Aphaenogaster subterranea (Latreille, 1798) (Fig. 19)

Published records: Dumbrăveni (Tăușan et al. 2011)

Messor cf. structor sensu Schlick-Steiner et al. (2006) (Fig. 19)

Published records: Agârbiciu, Gușterița, Slimnic, Ocna Sibiului, (Markó, Csősz 2002).

New records: 3 w, Sibiu, Hammersdorf Tal. [Dealul Gușteriței], 27.09.1970, as *Messor barbarous*, leg. E. Schneider, coll. E. Schneider,

det. Tăușan et al. 2012; 1 w, Racovița, 21.08.2010, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan.

Tribe Solenopsidini

Solenopsis fugax (Latreille, 1798) (Fig. 13)

Published records: Dealul Gușteriței, Gușterița, Sibiu (Markó, Csősz 2002); Cîsnădioara (Tăușan et al. 2012).

New records: 5 w, Hermannstadt, Reussbachwiesen, 26.04.1969, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 q, Slimnic, D. Zackel, 24.08.1972, leg. E. Schneider, coll. E. Schneider; 3 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan.

Monomorium pharaonis (Linnaeus, 1758) (Fig. 13)

Published records: Sibiu (Tăușan, Markó 2011)

Tribe Stenammini

Stenamma debile (Förster, 1850) (Fig. 13)

Published records: Dumbrava Sibiului (Tăușan, Markó 2009)

New records: 1 w, Racovița, 06.2009, leg. G. Mărginean, det. Tăușan et al. coll. Tăușan; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan.

Tribe Tetramoriini

Tetramorium cf. caespitum (Linnaeus, 1758) (Fig. 19)

Published records: Agârbiciu, Gușterița, Păltiniș, Rășinari, Rîu Sadului (Markó, Csősz 2002).

New records: 3 w, Hermannstadt, Reussbachwiesen, 26.04.1969, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 4 q, Poplaca, b. Hermannstadt, 11.09.1969, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 1 w, Trassilv. merid., Ocna Sibiului, 13.07.1973, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 4 w, Trassilv. merid., Ocna Sibiului, 19.05.1974, leg. E. Schneider, coll. E. Schneider, det. Tăușan et al. 2012; 9 w, Ocna Sibiului, 07.2009, leg. Tăușan et al. coll. Tăușan; 9 w, Racovița, 21.08.2010 leg. G. Mărginean, det. Tăușan et al. coll. Tăușan ; 3 w, Dumbrăveni 07.2011, leg. Tăușan et al. coll. Tăușan.

Tetramorium ferox Ruzsky, 1903 (Fig. 19)

Published records: Sibiu (Tăușan, Markó 2009)

Tetramorium cf. impurum (Förster, 1850) (Fig. 19)

Published records: Sibiu (Markó, Csősz 2002).

Subfamily Ponerinae Lepeletier, 1836

Tribe Ponerini

Ponera coarctata (Latreille, 1802) (Fig. 20)

Published records: Sibiu (Markó, Csősz 2002).

Ponera testacea Emery, 1895 as *Ponera coarctata*
var. *testacea* Emery, 1985 (Fig. 20)

Published records: Sibiu (Markó, Csősz 2002).

Comments

Altogether 70 species were identified, one being exclusively indoor species. This number is quite high, considering that it represents 76.3 % of the Romanian myrmecofauna. A possible explanation could be the high diversity of habitats (mountains, hilly areas, lowlands, rivers meadows etc.) and the temperate climate. Nevertheless several myrmecological studies were undertaken in this area. Thus other species can occur, but the number cannot increase spectacularly. Though the presence of *Crematogaster schmidtii* in mountain areas such as Păltiniș is questionable, the proof material is available. One species is new for the ant fauna of Sibiu County, namely *Tapinoma subboreale*. It

was described by Seifert (2012), replacing the formerly known *Tapinoma ambiguum* Emery, 1925. *Tapinoma subboreale* is widespread in north and central Europe (Seifert, 2012).

Acknowledgements

The authors are grateful for the comments of Klara Benedek and Dr. Corneliu Bucșa, who improved the first version of the manuscript. We are in debt to Georgiana Mărginean, who provided the ant samples from Racovița. We owe thanks to Aurelia Ștefu, Oana Bota and Adriana Cravă for the help on the ant material from Dumbrăveni and also Aurelian Bordei for the help on the ant material from Sadu. Ana-Maria Mesaroș is acknowledged for constructive comments.

This work was possible with the financial support of the Sectoral Operational Programme for Human Resources Development 2007-2013, cofinanced by the European Social Fund, under the project number POSDRU/107/1.5/S/76841 with the title „Modern Doctoral Studies: Internationalization and Interdisciplinarity”.

REFERENCES

- Bucşa, Tăușan 2011 Bucşa Corneliu, Tăușan Ioan, *Istoricul cercetărilor entomologice din împrejurimile Sibiului. Partea I* In: Brudea et al. (eds.) *Volum de lucrări al Simpozionului "Biodiversitatea și Managementul Insectelor din Romania" in memoria entomologului bucovinean Ioan Nemeș*, Suceava (2011), p. 179-199.
- Costea 2011 Costea Marioara, *Regionalization of present modelling processes and their risk character. Case study: Sibiu County*. In: Brukenthal. Acta Musei VI.3, Sibiu (2011), p. 559-572.
- Csösz, Markó 2005 Csösz Sándor, Markó Bálint, *European ant species (Hymenoptera: Formicidae) in the ant collection of the Natural History Museum of Sibiu (Hermannstadt/Nagyszeben), Romania II. Subfamily Formicidae*. In: *Annales Historico-Naturales Musei Nationalis Hungarici* 97, (2005), p. 225-240.
- Csösz et al. 2011 Csösz Sándor, Markó Bálint, Gallé László, *The myrmecofauna (Hymenoptera: Formicidae) of Hungary: an updated checklist*. In: *North-Western Journal of Zoology* 7(1), Oradea (2011), p. 55-62.
- Czekes et al. 2012 Czekes Zsolt, Radchenko Alexander, Csösz Sándor, Szász-Len Annamária, Tăușan Ioan, Kiss Klára, Markó Bálint, *The genus Myrmica Latreille, 1804 (Hymenoptera: Formicidae) in Romania: new data, distribution of species and key for their identification*. In: *Entomologica Romanica* 17, Cluj-Napoca (2012) (in press).
- Czechowski et al. 2002 Czechowski Wojciech, Radchenko, Alexander, Czechowska, Wiesława, *The ants (Hymenoptera, Formicidae) of Poland*. In: *Museum and Institute of Zoology PAS, Studio 1*, Warsaw (2002).
- Karaman 2004 Karaman Marko, *Checklist of known species of ants (Hymenoptera, Formicidae) in the fauna of Montenegro*. In: *Natura Montenegrina* 3, Podgorica (2004), p. 83-92.
- Karaman 2008 Karaman Marko, *Two new species of the Crematogaster scutellaris group, Crematogaster gordani, sp. nov. and C. montenegrinus sp. nov. (Insecta: Hymenoptera: Formicidae) from Crna Gora (Montenegro) with a key to this group from Southern Europe*. In: *Natura Montenegrina* 7(1), Podgorica (2008), p. 5-24.
- Lapeva-Gjonova et al. 2010 Lapeva-Gjonova Albena, Antonova Vera, Radchenko Alexander, Atanasova Maria, *Catalogue of ants (Hymenoptera: Formicidae) of Bulgaria*. In: *ZooKeys* 62, (2010), p. 1-124.
- Markó 2008a Markó Bálint, *Ants (Hymenoptera, Formicidae) of the „Sand Dunes of Foieni” protected area and its surroundings (Satu Mare County, Romania) and a new species for the Romanian fauna*. In: *Acta Scientiarum Transylvanica – Múzeumi Füzetek* 16(3), Cluj-Napoca (2008), p. 87-99.
- Markó 2008b Markó Bálint, *Pyramica baudueri (Emery, 1875) – a new ant species (Hymenoptera: Formicidae) for the Romanian fauna*. In: *Fragmenta Faunistica* 51(2), Warsaw (2008), p. 101-106.
- Markó, Csösz 2001 Markó Bálint, Csösz Sándor, *Nine new ant species in the Romanian fauna (Hymenoptera: Formicidae): morphology, biology, and distribution*. In: *Entomologica Romanica* 6, Cluj-Napoca (2001), p.127-132
- Markó, Csösz 2002 Markó Bálint, Csösz Sándor, *Die europäischen Ameisenarten (Hymenoptera: Formicidae) des Hermannstädter (Sibiu, Rumänien) Naturkundemuseums I.: Unterfamilien Ponerinae, Myrmicinae und Dolichoderinae*. In: *Annales Historico-Naturales Musei Nationalis Hungarici* 94, (2002), p. 109-121.
- Markó et al. 2006 Markó Bálint, Sipos Botond, Csösz Sándor, Kiss Klára, Boros István, Gallé László, *A comprehensive list of the ants of Romania (Hymenoptera: Formicidae)*. In: *Myrmecological News* 9, (2006), p. 65-76.
- Markó et al. 2009 Markó, B., Ionescu-Hirsch, A., Szász-Len, A., *Genus Camponotus Mayr, 1861 (Hymenoptera: Formicidae) in Romania: distribution and identification key to the worker caste*. In: *Entomologica Romanica* 14, Cluj-Napoca, (2009), p. 29-41.
- Moscaliuc 2009 Moskaliuc Liviu, *Notes on the distribution of the genus Plagiolepis Mayr, 1861*

- (Hymenoptera: Formicidae) in Dobrogea, and the first record of *Plagiolepis obscuriscapa* Santschi, 1923 in Romania. In: *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"* 52, Bucharest (2009), p. 297-301.
- Paraschivescu 1975 Paraschivescu Dinu, *Investigations upon the Formicidae belonging to the collections of the "Gr. Antipa" museum – Bucharest*. In: *Travaux du Muséum d'Histoire Naturelle "Grigore Antipa"* 16, Bucharest, (1975), p. 187-192.
- Pascu, Schneider 1998 Pascu Mariana, Schneider Eckbert, *Colecțiile entomologice ale Muzeului de Istorie Naturală din Sibiu*. In: *Studii și Comunicări de Științele Naturii, Muzeul Brukenenthal* 27, Sibiu (1997) p. 201-218.
- Petrov 2006 Petrov Ivan, *Mravi Srbije i Crne Gore*, In: SANU, Belgrad (2006).
- Radchenko 1989 Radchenko Alexander, *The ants of the genus Plagiolepis MAYR of the European part of USSR*. In: *Vestnik Zoologii* 9, (1989), p. 153-156.
- Seifert 2007 Seifert Bernhard, *Die Ameisen Mittel-und Nordeuropas*. In: Lutra-Verlag, (2007) Tauer
- Seifert 2012 Seifert Bernhard, *Clarifying naming and identification of the outdoor species of the ant genus Tapinoma Förster, 1850 (Hymenoptera: Formicidae) in Europe north of the Mediterranean region with description of a new species*. In: *Myrmecological News* 16, (2012), p. 139-147.
- Seifert, Schultz 2009 Seifert Bernhard, Schultz Roland, *A taxonomic revision of the Formica rufibarbis Fabricius, 1793 group (Hymenoptera: Formicidae)*. In: *Myrmecological News* 12, (2009), p. 255-272.
- Seifert et al. 2009 Seifert Bernhard. Schlick-Steiner, Birgit, Steiner Florian, *Myrmica constricta Karavajev, 1934 - a cryptic sister species of Myrmica hellenica Finzi, 1926 (Hymenoptera: Formicidae)*. In: *Soil Organisms* 81(1), (2009), p. 53-76.
- Schlick-Steiner et al. 2006 Schlick-Steiner Birgit. Steiner Florian, Konrad Heino, Markó Bálint, Csősz Sándor, Heller Gerhard, Ferencz Beatrix, Sipos Botond, Christian Erhard, Stauffer Christian, *More than one species of Messor harvester ants (Hymenoptera: Formicidae) in Central Europe*. In: *European Journal of Entomology* 103, (2006), p. 469-476.
- Tăușan, Markó 2009 Tăușan Ioan, Markó Bálint, *Comparative analysis of ant communities (Hymenoptera: Formicidae) in the surroundings of Sibiu (Romania)*. In: *Brukenenthal Acta Musei* IV.3, Sibiu (2009), p. 635-644.
- Tăușan, Markó 2011 Tăușan Ioan, Markó Bálint, *New records of the pharaoh ant Monomorium pharaonis (Linnaeus, 1758) (Hymenoptera: Formicidae) in Romania*. In: *Entomologica Romanica* 16, Cluj-Napoca (2011), p. 23-26.
- Tăușan et al. 2011 Tăușan Ioan, Bota Oana, Ștefu Aurelia, Cravă Adriana, *Aphaenogaster subterranea (Latreille, 1798) (Hymenoptera: Formicidae) in Romania: new records, distribution and habitat preferences*. In: *Brukenenthal Acta Musei* VI.3, Sibiu (2011), p. 459-464.
- Tăușan et al. 2012 Tăușan Ioan, Tatu Alexandru, Cravă Adriana, *Comparative analysis of pitfall traps with different liquids for studying ground-dwelling insects with special reference to ant communities (Hymenoptera: Formicidae)*. In: *Acta Oecologica Carpatica* V, Sibiu (2012), p. 145-152.

LIST OF ILLUSTRATIONS

- Fig. 1.** Sibiu County limits (modified after Google Earth)
- Fig. 2.** Distribution of *Dolichoderus quadripunctatus* (*Dolq*), *Tapinoma erraticum* (*Tera*) and *T. suboreale* (*Tsub*) (modified after Google Earth)
- Fig. 3.** Distribution of *Camponotus aethiops* (*aet*), *C. fallax* (*fal*) and *C. truncatus* (*trun*) (modified after Google Earth)
- Fig. 4.** Distribution of *Camponotus ligniperdus* (*lig*) and *C. herculeanus* (*her*) (modified after Google Earth)
- Fig. 5.** Distribution of *Camponotus piceus* (*pic*) and *C. vagus* (*vag*) (modified after Google Earth)
- Fig. 6.** Distribution of *Formica cinera* (*cin*), *F. cunicularia* (*cun*), *F. fusca* (*fus*), *F. lemani* (*lem*) and *F. clara* (*clara*) (modified after Google Earth)
- Fig. 7.** Distribution of *Polyergus rufescens* (*pol*), *Formica sanguinea* (*sang*), *F. rufa* (*rufa*) and *F. gagates* (*gag*) (modified after Google Earth)
- Fig. 8.** Distribution of *Formica polystena* (*pol*), *F. pratensis* (*pra*) and *F. rufibarbis* (*rufi*) (modified after Google Earth)
- Fig. 9.** Distribution of *Lasius alienus* (*ali*), *L. bicornis* (*bic*), *L. citrinus* (*cit*), *L. distinguendus* (*dist*), *L. emarginatus* (*emar*) and *L. umbratus* (*umbr*) (modified after Google Earth)
- Fig. 10.** Distribution of *Lasius fuliginosus* (*ful*), *L. flavus* (*fla*) and *L. brunneus* (*brun*) (modified after Google Earth)
- Fig. 11.** Distribution of *Lasius platythorax* (*plat*), *L. paralienus* (*par*) and *L. niger* (*nig*) (modified after Google Earth)
- Fig. 12.** Distribution of *Plagiolepis pygmaea* (*plpyg*), *P. taurica* (*pltau*) and *Prenolepis nitens* (*pnit*) (modified after Google Earth)
- Fig. 13.** Distribution of *Stenamma debile* (*deb*), *Crematogaster schmidti* (*csch*), *Solenopsis fugax* (*fug*), *Monomorium pharaonis* (*mon*) and *Myrmecina graminicola* (*mgram*) (modified after Google Earth)
- Fig. 14.** Distribution of *Formicoxenus nitidulus* (*fnit*), *Leptothorax acervorum* (*lacer*), *Temnothorax affinis* (*tafis*), *T. corticalis* (*tcor*), *T. crassispinus* (*tcrass*), *T. unifascitatus* (*tunif*) and *T. clypeatus* (*tcly*) (modified after Google Earth)
- Fig. 15.** Distribution of *Manica rubida* (*rubida*), *Myrmica slovacica* (*slo*), *M. rugulosa* (*rug*), *M. specioidea* (*spe*), *M. schenki* (*sck*) and *M. lobicornis* (*lob*) (modified after Google Earth)
- Fig. 16.** Distribution of *Myrmica ruginodis* (*rug*) and *Myrmica rubra* (*rub*) (modified after Google Earth)
- Fig. 17.** Distribution of *Myrmica scabrinodis* (*sca*) and *M. sabuleti* (*sab*) (modified after Google Earth)
- Fig. 18.** Distribution of *Myrmica constricta*, *M. lonae*, *M. gallienii* and *M. sulcinodis* (modified after Google Earth)
- Fig. 19.** Distribution of *Aphaenogaster subterranea* (*asub*), *Messor* cf. *structor* (*mes*) and *Tetramorium* cf. *caespitum* (*tcae*), *T.* cf. *impurum* (*timp*) and *T. ferox* (*tfer*) (modified after Google Earth)
- Fig. 20.** Distribution of *Ponera coarctata* and *P. testacea* (modified after Google Earth)

LISTA ILUSTRAȚIILOR

- Fig. 1.** Limitele județului Sibiu (modificat după Google Earth)
- Fig. 2.** Distribuția speciilor *Dolichoderus quadripunctatus* (Dolq), *Tapinoma erraticum* (Tera) și *T. suboreale* (Tsub) (modificat după Google Earth)
- Fig. 3.** Distribuția speciilor *Camponotus aethiops* (aet), *C. fallax* (fal) și *C. truncatus* (trun) (modificat după Google Earth)
- Fig. 4.** Distribuția speciilor *Camponotus ligniperdus* (lig) și *C. herculeanus* (her) (modificat după Google Earth)
- Fig. 5.** Distribuția speciilor *Camponotus piceus* (pic) și *C. vagus* (vag) (modificat după Google Earth)
- Fig. 6.** Distribuția speciilor *Formica cinera* (cin), *F. cunicularia* (cun), *F. fusca* (fus), *F. lemani* (lem) și *F. clara* (clara) (modificat după Google Earth)
- Fig. 7.** Distribuția speciilor *Polyergus rufescens* (pol), *Formica sanguinea* (sang), *F. rufa* (rufa) și *F. gagates* (gag) (modificat după Google Earth)
- Fig. 8.** Distribuția speciilor *Formica polyctena* (pol), *F. pratensis* (pra) și *F. rufibarbis* (rufi) (modificat după Google Earth)
- Fig. 9.** Distribuția speciilor *Lasius alienus* (ali), *L. bicornis* (bic), *L. citrinus* (cit), *L. distinguendus* (dist), *L. emarginatus* (emar) și *L. umbratus* (umbr) (modificat după Google Earth)
- Fig. 10.** Distribuția speciilor *Lasius fuliginosus* (ful), *L. flavus* (fla) și *L. brunneus* (brun) (modificat după Google Earth)
- Fig. 11.** Distribuția speciilor *Lasius platythorax* (plat), *L. paralienus* (par) și *L. niger* (nig) (modificat după Google Earth)
- Fig. 12.** Distribuția speciilor *Plagiolepis pygmaea* (plpyg), *P. taurica* (pltau) și *Prenolepis nitens* (pnit) (modificat după Google Earth)
- Fig. 13.** Distribuția speciilor *Stenamma debille* (deb), *Crematogaster schmidti* (csch), *Solenopsis fugax* (fug), *Monomorium pharaonis* (mon) și *Myrmecina graminicola* (mgram) (modificat după Google Earth)
- Fig. 14.** Distribuția speciilor *Formicoxenus nitidulus* (fnit), *Leptothorax acervorum* (lacer), *Temnothorax affinis* (tafis), *T. corticalis* (tcor), *T. crassispinus* (tcrass), *T. unifasciatus* (tunif) și *T. clypeatus* (tcly) (modificat după Google Earth)
- Fig. 15.** Distribuția speciilor *Manica rubida* (rubida), *Myrmica slovacica* (slo), *M. rugulosa* (rugu), *M. specioidea* (spe), *M. schenki* (sck) și *M. lobicornis* (lob) (modificat după Google Earth)
- Fig. 16.** Distribuția speciilor *Myrmica ruginodis* (rug) și *Myrmica rubra* (rub) (modificat după Google Earth)
- Fig. 17.** Distribuția speciilor *Myrmica scabrinodis* (sca) și *M. sabuleti* (sab) (modificat după Google Earth)
- Fig. 18.** Distribuția speciilor de *Myrmica constricta*, *M. lonae*, *M. gallienii* și *M. sulcinodis* (modificat după Google Earth)
- Fig. 19.** Distribuția speciilor *Aphaenogaster subterranea* (asub), *Messor* cf. *structor* (mes) și *Tetramorium* cf. *caespitum* (tcae), *T.* cf. *impurum* (timp) și *T. ferox* (tfer) (modificat după Google Earth)
- Fig. 20.** Distribuția speciilor *Ponera coarctata* și *P. testacea* (modificat după Google Earth)




Fig. 1. Map of Sibiu County limits




Fig. 2. Distribution of *Dolichoderus quadripunctatus* (Dolq), *Tapinoma erraticum* (Tera) and *T. suboreale* (Tsub)




Fig. 3. Distribution of *Camponotus aethiops* (aet), *C. fallax* (fal) and *C. truncatus* (trun)




Fig. 4. Distribution of *Camponotus ligniperdus* (lig) and *C. herculeanus* (her)




Fig. 5. Distribution of *Camponotus piceus* (pic) and *C. vagus* (vag)




Fig. 6. Distribution of *Formica cinera* (cin), *F. cunicularia* (cun), *F. fusca* (fus), *F. lemani* (lem) and *F. clara* (clara)




Fig. 7. Distribution of *Polyergus rufescens* (pol), *Formica sanguinea* (sang), *F. rufa* (rufa) and *F. gagates* (gag)




Fig. 8. Distribution of *Formica polycetena* (pol), *F. pratensis* (pra) and *F. rufibarbis* (rufi)




Fig. 9. Distribution of *Lasius alienus* (*ali*), *L. bicornis* (*bic*), *L. citrinus* (*cit*), *L. distinguendus* (*dist*), *L. emarginatus* (*emar*) and *L. umbratus* (*umbr*)




Fig. 10. Distribution of *Lasius fuliginosus* (*ful*), *L. flavus* (*fla*) and *L. brunneus* (*brun*)




Fig. 11. Distribution of *Lasius platythorax* (plat), *L. paralienus* (par) and *L. niger* (nig)



Fig. 12. Distribution of *Plagiolepis pygmaea* (plpyg), *P. taurica* (pltau) and *Prenolepis nitens* (pnit)




Fig. 13. Distribution of *Stennama debille* (deb), *Crematogaster schmidti* (csch), *Solenopsis fugax* (fug), *Monomorium pharaonis* (mon) and *Myrmecina graminicola* (mgram)




Fig. 14. Distribution of *Formicoxenus nitidulus* (fnit), *Leptothorax acervorum* (lacer), *Temnothorax affinis* (tafis), *T. corticalis* (tcor), *T. crassispinus* (tcrass), *T. unifascitatus* (tunif) and *T. clypeatus* (tcly)




Fig. 15. Distribution of *Manica rubida* (rubida), *Myrmica slovacica* (slo), *M. rugulosa* (rugu), *M. specioides* (spe), *M. schenki* (sck) and *M. lobicornis* (lob)




Fig. 16. Distribution of *Myrmica ruginodis* (rug) and *Myrmica rubra* (rub)




Fig. 17. Distribution of *Myrmica scabrinodis* (sca) and *M. sabuleti* (sab)




Fig. 18. Distribution of *Myrmica constricta*, *M. lonae*, *M. gallieni* and *M. sulcinodis*




Fig. 19. Distribuion of *Aphaenogaster subterranea* (asub), *Messor* cf. *structor* (mes) and *Tetramorium* cf. *caespitum* (tcae), *T.* cf. *impurum* (timp) and *T. ferox* (tfer)




Fig. 20. Distribution of *Ponera coarctata* and *P. testacea*

THE BIELZ MALACOLOGICAL SLUG COLLECTION FROM THE NATURAL HISTORY MUSEUM IN SIBIU

Ana-Maria MESAROS*
Raluca STOICA**

Abstract. 75 slug specimens belonging to the Bielz Malacological collection from the Natural History Museum in Sibiu were listed and studied. Because of the general storage conditions and the lack of staff, during time, the majority of the slug specimens found in the Bielz collection dried up. The authors registered the specimens and their conservation aspects in order to identify the proper curatorial methods that should be appointed to these particular museum pieces. Active research is necessary to support passive conservation measures in order to stop further degradation to the wet Malacological collection.

Keywords: slug collection, museum specimen conservation, Natural History Museum from Sibiu

Rezumat. Au fost inventariate 75 specimene de limacomorfe aparținând colecției malacologice Bielz din cadrul Muzeului de Istorie Naturală (Sibiu). Datorită condițiilor generale de depozitare și a lipsei personalului calificat, de-a lungul timpului, majoritatea specimenelor de limacomorfe din colecția Bielz conservate în lichid s-au uscat. Autorii au inventariat speciemenele și au studiat aspectele de conservare cu privire la aceste piese muzeale cu scopul de a identifica care sunt metodele optime ce ar trebui aplicate acestora. Cercetarea activă a stării actuale de deteriorare este necesară pentru a stabili măsurile de restaurare viitoare ce pot fi aplicate speciemenelor și pentru a opri degradarea altor preparate umede din colecția Malacologică a muzeului.

Cuvinte cheie: colecția de limacși, conservarea speciemenelor muzeale, Muzeul de Istorie Naturală Sibiu

Introduction

The Natural History Museum from Sibiu was founded in 1895 by the Transylvanian Society for Natural Sciences from Sibiu (*Siebenbürgischer Verein für Naturwissenschaften zu Hermannstadt*). The collections were initiated in 1849 long before the museum was open to the public.

The Malacological collection from the Natural History Museum in Sibiu gathers today 515.147 specimens, and the value of the collection consists not only in its numbers but also in its historical background, offering many opportunities in approaching systematic, ecological, biogeographic and biodiversity assessment studies of different time scale (Mesaros 2010, 649).

The Malacological collection of the museum includes the Kimakowicz collection (305.431 specimens) and the Transylvanian Society for Natural Sciences of Sibiu Malacological collection (209.716 specimens).

The Transylvanian Society for Natural Sciences of Sibiu Malacological collection originated in 1852 when Johann L. Neugeboren, founding member of the Society, donated 210 mollusc shells.

The Bielz Malacological collection, counting over 200.000 pieces, was initiated by Michael Bielz (1787-1866), one of the founding members of the museum and the first director of the Transylvanian Society. Edward Albert Bielz (1827-1898), his son, president of the Society for many years also, shared this passion for natural sciences and Malacology.

The collection was donated in 1927 to the Society by Julius Bielz (1884 – 1958) the grandson of Eduard Albert Bielz.

The material found in the Bielz Malacological collection was mainly collected and identified by Eduard Albert Bielz. The results of his Malacological studies were published in the Society journal *Verhandlungen und Mittheilungen des siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* from 1850 until 1865 as short papers and later all the data from these communications was gathered in a volume entitled *Fauna der Land- und Süßwasser*

*Brukenthal National Museum, Natural History Museum,

*ana.mesaros@yahoo.com,

ana.mesaros@brukenthalmuseum.ro;

**ralucastoicas@yahoo.com,

raluca.stoica@brukenthalmuseum.ro

Mollusken Siebenbürgens published in 1863. Eduard Albert Bielz described and listed, in the journal and in the volume published in 1863, the specimens collected and included in the collection but not with inventory numbers.

The Bielz Collection has two main categories: wet and dried specimens. The slug collection is part of the wet collection.

Unfortunately, because of the general storage conditions and the lack of staff, during time, the majority of the slug specimens dried out. After listing the museum specimens one of the objectives was to stop further degradation

Material and methods

Curatorial aspects

The entire Transylvanian Society Malacological collection holds 103 slug specimens, from which 75 belong, according to their labels, to the Bielz Collection, the rest were collected by museum curators between 1952 and 1959.

As the Bielz Malacological collection was donated to the Society, it is since 1927 included, by the general accounting inventory books, in the Transylvanian Society for Natural Sciences of Sibiu Malacological collection. All donations that entered the museum, in that period, were included in the Society collection. These collections were not kept separately, as they were received, but each specimen was given an inventory number and stored together with the other specimens of the same genera and species. The inventory numbers changed and the specimens were moved by museum curators during time.

Regarding the Bielz Malacological collection the unit-level data is included on old original labels which in many cases are incomplete or hard to read, or even more disturbing some specimen's labels are missing. The recording of the collecting data associated to the specimens was not a priority to collectors, because the number of specialists and museum staff was reduced and the Society members were collecting natural specimens belonging to different scientific areas. For example, the museum includes both a mineralogical and molluscan Bielz collection. Also, the Society did not benefit from a true headquarter until the museum was opened in 1895, and the collections were moved from one place to another and thus, we presume that a lot of written information was lost. That is why the authors, consider, that this might be the reason why there are no written inventory records by Michael or Eduard Albert Bielz regarding their Malacological collection.

There were identified in the museum archive some donation letters for specimens but none of them are related to slug specimens.

In order to gather more information for the slug specimens, to complete the data missing from the original labels the Society scientific journal *Verhandlungen und Mittheilungen des siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* and E. A. Bielz's *Fauna der Land- und Süßwasser Mollusken Siebenbürgens* were used.

Slug specimens are mentioned for the first time in the paper written by Michael Bielz in 1851 and published in the museum journal, in which the author mentioned the species, the collecting sites but not the total number of specimens or an inventory number. The species collected were: *Arion empiricorum* var *ater* Férussac, 1819 and *A. empiricorum* var *rufus* (Linnaeus, 1758) synonym *Arion ater* (Linnaeus, 1758) collected from Hermannstadt meaning Sibiu (Sibiu County), *Arion albus* Linnaeus synonym *Arion ater* (Linnaeus, 1758) collected from Gurariu am Valye plaje mening Gura Răului (Sibiu County) on the river bank, *Arion subfuscus* (Draparnaud, 1805) no collecting sites received from A. Szombath, *Arion hortensis* Férussac, 1819 collected from Hermannstadt meaning Sibiu (Sibiu County), *Limax cinereus* O. F. Müller, 1774 today *Limax maximus* Linnaeus, 1758 in many varieties as the author mentioned collected from Hermannstadt, Sibiu, *Limax agreste* Linnaeus, 1758 today *Deroceras agreste* (Linnaeus, 1758) collected from Hermannstadt, Sibiu and *Limax coeruleus* M. Bielz, 1851 synonym *Bielzia coeruleus* (M. Bielz, 1851) described for the first time by the author in this paper, collected on May 24, 1847 from Draguscher und Arpaser Gebirge, Rotunde bey Kapnikbánya meaning Drăguş (Braşov County), mountain area near Arpaşu (Sibiu County), Pasul Rotunda between Căvnic and Budeşti (Maramureş County) (Bielz, 1851, 14-15).

Eduard Albert Bielz in the first chapter of *Beitrag zur kenntniss der siebenbürgischen Land- und Süßwasser mollusken* mentioned the same species but without the collecting dates or sites and in the second subchapter entitled *Bemerkungen zu dem Verzeichnisse der Siebenbürger Land- und Süßwassermollusken nebst der Beschreibung der neuen Arten* he included *Daudebardia brevipes* Draparnaud, 1805, no collecting sites or date (Bielz, 1853, 113). In the 1856 December number of the museum journal the same author listed the following species without mentioning the

collecting area and dates: *Arion empiricorum* Férussac, 1819 [*Arion ater* (Linnaeus, 1758)], *Arion albus* Linnaeus, 1758 [*Arion ater* (Linnaeus, 1758)], *Arion subfuscus* (Draparnaud, 1805), *Arion hortensis* Férussac, 1819, *Limax cinereus* var. *coerulans*, *Limax agrestis* Linnaeus, 1758 [*Deroceras agreste* (Linnaeus, 1758)] and *Daudebardia brevipes* Draparnaud, 1805 (Bielz, 1856, 225).

The numbers 9 and 10 of the museum journal from 1859 included general observations regarding the genera *Arion* Férussac, 1819, *Limax* Linnaeus, 1758 and *Daudebardia* Hartmann, 1821 and the first description of the species *Daudebardia transsilvanica* E. A. Bielz, 1859 synonym *Cibinia transsilvanica* (E.A. Bielz, 1859), collected from Transylvania, several localities, also near Braşov (Bielz, 1859, 214).

In *Fauna der Land- und Süßwasser Mollusken Siebenbürgens*, E. A. Bielz listed the following slug species: *Daudebardia transsilvanica* E. A. Bielz, 1859 synonym *Cibinia transsilvanica* (E.A. Bielz, 1859), *Arion olivaceus* Kotula, 1884 synonym *Arion subfuscus* (Draparnaud, 1805), *Arion hortensis* Férussac, 1819, *Limax cinereus* O.F. Müller, 1774 synonym *Limax maximus* Linnaeus, 1758, *Limax marginatus* Draparnaud 1805 synonym *Milax rusticus* Millet, 1843, *Limax variegatus* Draparnaud, 1801 synonym *Limax flavus* Linnaeus, 1758, *Limax agreste* Linnaeus, 1758 synonym *Deroceras agreste* (Linnaeus, 1758), *Limax silvaticus* Goldfuss, 1856 synonym *Lehmannia marginata* (Müller, 1774) (Bielz, 1863, 25-32).

The last paper published in the museum journal in which E.A. Bielz mentioned slug species is the *Catalogus molluscorum terrestrium et fluviatilium Imperii Austriaci* in which the author enumerated the slug species found in the Austrian Empire including Romania (Bielz, 1865, 139).

For the inventory list the authors compared the specimen scientific names and the collecting sites from the labels to those included in the written work in order to narrow down the periods when the specimens were collected and who was the collector.

As a general observation the specimens found in the collection were gathered mainly from Romania (Fig. 1, Tab. 1).

The Bielz collection slug specimens are held in glass jars (Fig. 2 a, b, c). The majority of the jars are closed with cork and covered with paper. The collector wrote on the paper covering the jar's cork

the name of the species and the collecting sites (Fig. 3 a, b).

In general the slug specimens belonging to the Bielz Collection have personalized labels with the name of Eduard Albert Bielz printed on them. The labels are glued to the jars making it easy to recognize and separate them from other donations (Fig. 4 a, b).

From a curatorial point of view it is important that the original data included on the labels should be mentioned for each specimen followed by the appropriate translations, where it is necessary. The information written on the labels from each jar was computerized, the specimens were taken out of the jars photographed and their general conservation aspects evaluated.

Conservation aspects

The Natural History Museum building was constructed in 1894 and its main design and purpose was to store the Society's collections and function as a museum.

The entire Bielz collection is held in a basement room of the museum together with other Malacological donations, bought dried and wet pieces are stored here.

The room is part of the museum storage area, which is effectively a series of rooms that open in a hallway. The storage area insulation is the same for the whole building and it is limited to brick walls, there are no modern modifications.

The Malacological collection storage room communicates through a door with the Mineralogy collection room.

The Malacological collection room lacks special climate controls. Thick orange drapes are fitted to the one window present here, in order to minimize solar gain. The room is exposed during the day to sun light especially at noon.

Heating in the storage area is provided by one radiator with thermostatic valve. The radiator is turned off from April until November. There is no controlled ventilation and the air enters underneath the door and through the window when it is opened. Also air exchange or associated dust movement through the floorboards is not prevented.

Being of smaller size and packed with wood cabinets, the room is prone to heat up quickly from radiant heating on the outside wall of the building, especially during summer.

The Malacological specimens are deposited in wooden boxes held in wood cabinets. The wood

cabinets, very old, are used to deposit dried and wet collection pieces together. The Bielz slug collection is stored in such conditions (Fig. 5 a, b).

The slug specimens are held in glass jars sealed with cork and covered with ordinary paper. The specimens are whole animals that were preserved in an unknown conservation fluid.

The Bielz wet Malacological collection was not checked periodically to top off the conservation liquid if necessary and now the liquid completely evaporated and the specimens dried out.

Results and discussion

Curatorial aspects

The slug specimens found in the collection were collected by Eduard Albert Bielz. The years when the specimens were collected are not mentioned on the label for each specimen.

The list with the slug specimens found in the Bielz collection, including the inventory number (Inv. No.), number of pieces found at that inventory number, the original identification done by E.A. Bielz, today's synonym, the collecting sites as written on the label and the collecting period if mentioned:

- Inv. No. 9274, 1 specimen, *Limax arboretum* probably *Limax arborum* Boughard, 1838, synonym to *Lehmannia marginata* O.F. Müller 1774 (Grossu 1983, 280), Hermannstadt Spital meaning the hospital in Sibiu, 9. XI. 1890;
- Inv. No. 66024, 66025, 66026, 66027, 66028, 66029, 6 specimens, *Arion subfuscus transilvanicus* Simroth synonym *Arion subfuscus quadrifasciata* Jensen synonym *Arion subfuscus* (Draparnaud, 1805) (Grossu, 1983, 47), Békéscsaba (Hungary);
- Inv. No. 66030, 66031, 66032, 66033, 66034, 66035, 66036, 66037, 66038, 66039, 66040, 66041, 12 specimens, *Limax marginatus* Draparnaud 1805 synonym *Milax rusticus* Millet, 1843 (Grossu, 1983, 220), Freckes Gebirge Transilvania meaning Avrig (Sibiu County), mountain area, Transylvania;
- Inv. No. 66042, 1 specimen, *Limax cinereus* O.F. Müller, 1774 synonym *Limax maximus maximus* Linnaeus, 1758 (Grossu, 1983, 242), Michelsberg meaning Cîsnădioara, (Sibiu County);
- Inv. No. 66043, 66044, 66045, 66046, 4 specimens, *Limax agrestis* synonym *Limax agreste* Linnaeus, 1758 synonym *Deroceras agreste* (Linnaeus, 1758) (Grossu, 1983, 334), Hermannstadt, Gutenwaisenhaus, meaning Sibiu, orphanage;

- Inv. No. 66047, 66048, 66049, 66050, 66051, 66052, 66053, 7 specimens, *Arion subfuscus transilvanicus* Simroth synonym *Arion subfuscus quadrifasciata* Jensen synonym *Arion subfuscus* (Draparnaud 1805) (Grossu, 1983, 47), Klausenburg Bükk. meaning Cluj-Napoca, Bükk. with no correspondence;

- Inv. No. 66054, 66055, 2 specimens, *Limax agrestis* synonym *Limax agreste* Linnaeus, 1758 synonym *Deroceras agreste* (Linnaeus, 1758) (Grossu, 1983, 334), Bihargebirge Branisora bis Stenisora meaning Bihor Mountains, from Bănişoara to Stănişoara;

- Inv. No. 66056, 1 specimen, *Limax maximus var Bielzii* Seibert, 1873 synonym *Limax bielzii* Seibert, 1873, St. Gotthard bei Graz, Steiermark meaning the Castle St. Gotthard, north Graz, Steiermark, Austria;

- Inv. No. 66057, 66058, 2 specimens, *Arion olivaceus* Kotula, 1884 synonym *Arion subfuscus* (Draparnaud, 1805), Freckes Gebirge, Transsilvan. meaning Avrig (Sibiu County), mountain area, Transylvania;

- Inv. No. 66059, 1 specimen, *Limax cinereoniger* Wolf, 1803, Hermannstadt, Transilv. meaning Sibiu, Transylvania;

- Inv. No. 66060, 1 specimen, *Limax unicolor* Heynemann synonym *Limax maximus var. unicolor* Heynemann (Grossu, 1983, 246), Transylvania;

- Inv. No. 66061, 66062, 66063, 66064, 4 specimens, *Arion subfuscus transilvanicus* Simroth synonym *Arion subfuscus quadrifasciata* Jensen synonym *Arion subfuscus* (Draparnaud 1805) (Grossu, 1983, 47), Hohe Rinne, meaning Păltiniş, Cindrel Mountains;

- Inv. No. 66065, 66066, 2 specimens, *Amalia marginata* Draparnaud 1805 synonym *Milax rusticus* Millet, 1843 (Grossu, 1983, 220), Gödesberg in Schloss, meaning Gödesberg castle, in Bad Gödesberg, Bonn (Germany);

- Inv. No. 66067, 1 specimen, *Limax transilvanicus* Hazay, 1883 synonym *Bielzia coerulans* (M. Bielz, 1851) (Grossu, 1983, 304), Karzisoara Transsilvanien meaning Cârţişoara (Sibiu County), Transylvania;

- Inv. No. 66068, 66069, 66070, 3 specimens, *Limax marginatus* Draparnaud, 1805 synonym *Milax rusticus* Millet, 1843 (Grossu, 1983, 220), Hohe Rinne, meaning Păltiniş, Cindrel Mountains;

- Inv. No. 66071, 66072, 66073, 66074, 66075, 66076, 6 specimens, *Limax cinereoniger* Wolf, 1803, Mühlbachthal, Colone Bisztra, meaning Sebeş, Bistra (Alba County);
- Inv. No. 66077, 66078, 2 specimens, *Arion subfuscus transilvanicus* Simroth synonym *Arion subfuscus quadrifasciata* Jensen synonym *Arion subfuscus* (Draparnaud 1805) (Grossu, 1983, 47), Klausenburg meaning Cluj-Napoca;
- Inv. No. 66079, 1 specimen, *Limax cinereoniger* Wolf, 1803, Graz (Austria);
- Inv. No. 66080, 1 specimen, *Limax cinereoniger* Wolf, 1803, Klausenburg meaning Cluj-Napoca;
- Inv. No. 66081, 66082, 2 specimens, *Arion subfuscus transilvanicus* Simroth synonym *Arion subfuscus quadrifasciata* Jensen synonym *Arion subfuscus* (Draparnaud 1805) (Grossu, 1983, 47), Hermannstadt, Hallerweise, meaning Sibiu, meadow area;
- Inv. No. 66083, 66084, 2 specimens, *Arion hortensis* Férussac, 1819, Transylvania;
- Inv. No. 66085, 66086, 2 specimens, *Limax agrestis* synonym *Limax agreste* Linnaeus, 1758 synonym *Deroceras agreste* (Linnaeus 1758) (Grossu, 1983, 334), Hidegşamos, meaning Someşul Rece River;
- Inv. No. 66087, 1 specimen, *Limax cinereoniger* Wolf, 1803, Hercules Bad meaning Herculan Baths,;
- Inv. No. 66088, 1 specimen, *Limax cinereoniger* Wolf, 1803, Bullea – See, meaning Bălea Lake, Făgăraş Mountains;
- Inv. No. 66089, 1 specimen *Arion subfuscus* (Draparnaud 1805), Schnee grube Riascus, no equivalent;
- Inv. No. 66090, 66091, 66092, 66093, 66094, 5 specimens, *Limax marginatus* Draparnaud 1805 synonym *Milax rusticus* Millet, 1843 (Grossu, 1983, 220), collected by E. A. Biele from Bullea-Sea, meaning Bălea Lake, Făgăraş Mountains;
- Inv. No. 66095, 1 specimen, *Limax cinereus* var. *coerulans*, *Bielzia coerulans* (M. Biele, 1851) Bihargebirge, Banisor bei Stănişoara, meaning Bihor Mountains, Stănişoara, Bănişor (Sălaj County);
- Inv. No. 66096, 66097, 2 specimens, *Arion subfuscus transilvanicus* Simroth synonym *Arion subfuscus quadrifasciata* Jensen synonym *Arion subfuscus* (Draparnaud 1805) (Grossu, 1983, 47),

Klausenburg, Sigmundgarten, meaning Cluj-Napoca, Sigmundgarten.

As the specimens are dried out (Fig. 6 a, b) it is impossible, with the current museum means, to confirm that the names of the species written on the original labels are correct.

Conservation aspects

Fundamentally, the temperature and humidity regimes of collections spaces are architectural issues, and it is the nature of the building in which collections are housed that will determine the levels of control able to be applied to the collections space (Padfield, Larsen 2004, 131).

The chamber where the collection is held is not a proper museum storage unit. There is no ventilation, there are no devices to measure the external factors that influence the specimen conservation status not even the temperature. Temperature affects the state of the preserving fluid. Over time, even small fluctuations in temperature will cause most closure to fail, and evaporation will occur. As the temperature rises, the vapor pressure within the storage container increases putting stresses on the storage containers and their seals. Temperature increases also increase the rates of reaction (Carter 2003, 101-102).

Viscri *et al.* (2006, 94) considered that manual control of temperature in an effort to reduce humidity fluctuations in a poorly scaled building, with inadequate levels of staffing, is an inefficient and ineffective use of human resources. At the Museum of Natural History in Sibiu climate control in the storage area has to be done using limited resources in an antiquated building. Unfortunately the storage conditions of these specimens can not be changed because of the lack of space and funding.

The current museum conservation technician has sheltered the listed slug specimens from light exposure during storage. Light caused permanent, irreversible damage to specimens (the effect is cumulative).

The container used to store a specimen, and the fluid preservative, surrounding the specimen constitute a micro-environment. If placed in a good, stable storage environment and maintained well, fluid-preserved specimen should last for hundreds of years. This micro-environment is affected by the macro-environment (light, temperature, and handling). Wet collections require constant maintenance and monitoring (Carter 2000, 19).

The wood cabinet in which the Bielz collection slug specimens are stored are opened often as some of the material held in them are often used for temporary exhibits and specimens are extracted to be studied for scientific papers. The cork sealing could have been damaged by the vibration produced as the cabinet was often opened. As the jars cork seals are broken and highly fluctuating temperatures caused the liquid to evaporate. Evaporation happens especially with older and dried-out sealing material.

The original jars in which the specimens were held have to be completely changed or at least the cork should be avoided when closing the jars, our specimens being the perfect example of what happens when the jar is closed with cork. It tends to be acidic and decomposes over time and also cork, a compressible material, reacts with preservative fluid and loses their compressibility.

In a perfectly sealed storage container the relative humidity should have little effect, but in our case the jars were not perfectly sealed. Most storage containers found in the Malacological wet collection have imperfect seals leading to fluid loss through evaporation and hence open to being affected by relative humidity levels. Even in the Society collection slug specimens collected during 1952 and 1959, even if the liquid did not evaporate and the jars are sealed better there can be observed the reduction of the liquid levels. However the seals on storage containers are rarely completely airtight, and many of the materials used for the construction of jar lids, are themselves very permeable to oxygen. The provision of oxygen into the fluid preservative can have numerous effects (Carter 2003, 103). As an example many of our museum wet specimens are preserved in formaldehyde. Oxidation can promote the pH of formaldehyde solutions to fall, and can degrade lipids, amino acids and pigments.

The paper used to cover the cork seals is not acid free and so it became yellow and is decomposing. Even so it is a positive aspect, as in many other wet collection specimens found in the museum collections the original labels inserted in the jars disintegrated, making it impossible for the curator to extract the data associated to the specimens.

In the Bielz slug collection there are specimens that are severely deteriorated, for example the inventory numbers: 66042 (Fig. 7, a) and 66059 (Fig. 7 b,c).

However, if the tissue dried out it may not be useless. Strum and Mayhew (2006, 24) mention the name of scientists that have written a couple of

papers based on anatomical data drawn from dried soft parts found inside old shells from museum collections, some of them more than 40 years old. For example, Houbrick (1992, 2) extracted radulae from museum *Cerithium* Bruguière specimens dried in their shells.

Strum (2006, 55) considers that if the preservative has evaporated and left behind desiccated tissue, rehydrating solutions are not always the best. If the specimen was stored in alcohol and it is now dried out then it should be left like that.

Rehydration is a possibility in the case of our collection dried out slug specimens. In order to select a less invasive method to rehydrate the specimens the authors consulted the scientific studies conducted in this area until today.

Studies show that the specimen can be soaked in 0.5% aqueous solution of trisodium phosphate (Van Cleave, Ross 1947, 318).

Thompson *et al.* (1966, 109) used a method that involved soaking the dried specimen in a mixture of 50% ethylene glycol and 50% water.

Carter (2000, 23) offers the following method for rehydrating specimens:

- first a solution of Decon 90 detergent and water is prepared in approximate concentration 1: 20, as the phosphates in the detergent penetrate the cells allowing water to osmose into them;
- the solution is added to the dried up material making sure that no labels are affected as detergent causes serious damage to paper and script;
- the specimen is left for an average of 3 days, checking daily to ascertain if the solution needs to be changed or renewed. The material is then rinsed thoroughly in running cold water before re-immersing.
- after rehydration is complete, the specimen is rinsed in running cold water for at least 10 minutes and introduced in alcohol (usually 80%) with information that it has been rehydrated.

The final material has to be checked for a further week. If the appearance of the alcohol changes it means that the material needs a further rinsing in water.

Another step towards a better conservation of the slug collection is a good record keeping

The procedures used to collect each specimen should have been noted, along with the methods used in preparing, preserving and using the specimen. If a specimen deteriorates, these records

can tell us why, may allow us to repair damage, and prevent it from happening to other museum specimens. Our museum did not benefit from such records from the past curators. The museum conservation technician has started conservation records for the museum specimens including the Bielz slug collection.

The way forward in caring for the Bielz slug collection, in the current conditions, is by controlling the specimen microenvironment and by constant monitorization.

Acknowledgements

The authors are grateful to Dr. Heike Riese from the Senckenberg Museum für Naturkunde Görlitz (Germany) and to Dorina Golban, conservation specialist from the Țării Crișurilor Museum, Oradea (Romania) for the scientific and constructive remarks.

REFERENCES

- Bielz 1851 Bielz Michael, *Verzeichniss der Land und Süsswasser-Mollusken Siebenbürgens*. In: *Verhandlungen und Mittheilungen des siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* II Jahrgang Januar no 1, Hermannstadt (1851), p. 14 – 15.
- Bielz 1853 Bielz Eduard, Albert, *Beitrag zur kenntniss der siebenbürgischen Land- und Süsswasser mollusken*. In: *Verhandlungen und Mittheilungen des siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* IV Jahrgang Juli no 7, Hermannstadt (1853), p. 113.
- Bielz 1856 Bielz Eduard Albert, *Malacologische Notizen aus Siebenbürgen*. In: *Verhandlungen und Mittheilungen des siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt*, VII Jahrgang December no 12, Hermannstadt (1856), p. 225.
- Bielz 1859 Bielz Eduard Albert, *Ueber einige neue Arten und Formen der siebenbürgischen Molluskenfauna*. In: *Verhandlungen und Mittheilungen des siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* X Jahrgang September v. October no. 9, 10, Hermannstadt (1859), p. 214.
- Bielz 1863 Bielz Eduard Albert, *Fauna der Land- und Süsswasser-Mollusken Siebenbürgens*. In: *Commission von Th. Steinhaussen's Buchhandlung, Sibiu* (1863), p. 25-32.
- Bielz 1869 Bielz Eduard Albert, *Catalogus molluscorum terrestrium et fluviatilium Imperii Austriaci*. In: *Verhandlungen und Mittheilungen des siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt* XVI Jahrgang August no 8, Hermannstadt (1869), p. 139.
- Carter 2000 Carter Julian, *The Conservation of Molluscan Collections*. In: *The Biology Curator. The Publication of the Biology Curator's Group*, issue 18 (2000), p. 19, 23.
- Carter 2003 Carter Julian, *The effects of preservation and conservation treatments of the DNA of museum invertebrate fluid preserved collections*, Wales (2003). Thesis presented for the degree of master of philosophy (manuscript), p. 101-103.
- Grossu 1983 Grossu Alexandru, *Gastropoda Romaniae vol. 4 Ordo. Stylommatophora, Suprafam. Arionacea, Zonitacea, Ariohantacea și Helicacea*. (1983) In: *Editura Litera, București* (1983), p. 47, 220, 242, 246, 280, 304, 334.
- Houbrick 1992 Houbrick Richard, *Monograph of the genus Cerithium Bruguière in the Indo-Pacific (Cerithiidae: Prosobranchia)*. In: *Smithsonian Contributions to Zoology, number 510* Smithsonian Institution Press, Washington (1992), p.2.
- Kozłowski 2012 Kozłowski Jan, *The significance of alien and invasive slug species for plant communities in agrocenoses*. In: *Journal of plant protection research* 52(1), (2012) p. 67.
- Mesaroş 2010 Mesaroş Ana-Maria, *The family Hygromiidae Tryon (Gastropoda: Stylomatophora) from the Malacological Collections of the Natural History Museum from Sibiu*. In: *Brukenthal Acta Musei* V.3, Sibiu (2010), p. 649.
- Padfield, Larsen 2004 Padfield Tim, Larsen Poul Klens, *How to design museums with a naturally stable climate*. In: *Studies in Conservation* 49, (2004), p. 131.
- Strum, Mayhew 2006 Strum Charles, Mayhew Ross, *Field and laboratory methods in Malacology*. In:

- Strum, Charles, Pearce, Timothy, Valdés, Ángel, *Mollusks: A guide to their study, collection and preservation. A publication of the American Malacological Society Los Angeles and Pittsburgh, Universal Publishers, Boca Raton, Florida* (2006), p. 24.
- Strum 2006 Strum Charles, *Archival and curatorial methods*, Chapter 5. In: Strum, Charles, Pearce, Timothy, Valdés, Ángel, *Mollusks: A guide to their study, collection and preservation. A publication of the American Malacological Society Los Angeles and Pittsburgh, Universal Publishers, Boca Raton, Florida* (2006), p. 55.
- Thomson *et al.* 1966 Thomson John, Thomson Mary, Drummond Shelby, *A method for restoring dried crustacean specimens to taxonomically usable conditions*. In: *Crustaceana* 10(1), (1966), p. 109.
- Van Cleave, Ross, 1947 Van Cleave Harley, Ross Jean, *A method of reclaiming dried zoological specimens*. In: *Science* 105, (1947), p. 318.
- Viscardi *et al.* 2006 Viscardi Paolo, Sigwart Julia, Monaghan Nigel, *Climate control in an Uncontrollable Building*. In: *Collection Forum* 21, (2006), p. 94.

LIST OF ILLUSTRATIONS

- Fig. 1.** The areas from which the slug specimens were collected
- Fig. 2 a, b, c.** General aspect of the glass jars in which the specimens are held
- Fig. 3 a, b.** The original hand writing from the jars paper tops
- Fig. 4 a, b.** Personalized original labels from the Bielz Collection
- Fig. 5 a, b.** The general storage area (a) and the wooden boxes in which the jars with slug specimens were held (b – circled area)
- Fig. 6 a, b.** Close up aspects of a dried up specimen (inventory number 66088)
- Fig. 7 a, b, c.** Deteriorated dried up specimens
- Tab. 1.** The slug species found in the Bielz collection, the collecting sites and number of specimens

LISTA ILUSTRAȚIILOR

- Fig. 1.** Zonele din care speciemenele au fost colectate
- Fig. 2 a, b, c.** Aspectul general al borcanelor de sticlă în care sunt păstrate speciemenele
- Fig. 3 a, b.** Inscripții originale de pe hârtia ce îmbracă dopul de plută
- Fig. 4 a, b.** Etichete originale personalizate din colecția Bielz
- Fig. 5 a, b.** Spațiul de depozitare general (a) și cutia de lemn în care borcanele cu specimene limacomorfe erau ținute (b – zona încercuită)
- Fig. 6 a, b.** Aspecte ale unui speciemen desicat (numarul de inventar 66088)
- Fig. 7 a, b, c.** Specimene deteriorate sever
- Tab. 1.** Specile de limacomorfe din colecția Bielz, zonele de colectare și numărul de specimene pentru fiecare specie




Fig. 1. The areas from which the slug specimens were collected (edited using <http://www.simplemappr.net/> [accessed May 2012])




Fig. 2 a, b, c. General aspect of the glass jars in which the specimens are held inventory number 66042 (a), 66041(b), 66054-66055 (c).



Fig. 3 a, b. The original hand writing from the jars paper tops



Fig. 4 a, b. Personalized original labels from the Bielz Collection



Fig. 5 a, b. The general storage area (a) and the wooden boxes in which the jars with slug specimens were held (b – circled area)




Fig. 6 a, b. Close up aspects of a dried up specimen (inventory number 66088)




Fig. 7 a, b, c. Deteriorated dried up specimens

Tab. 1. The slug species found in the Bielz collection, the collecting sites and number of specimens

Species	Collecting sites	Number of specimens
<i>Lehmannia marginata</i> O.F. Müller, 1774	Sibiu (Sibiu County, Romania)	1
<i>Arion subfuscus</i> (Draparnaud, 1805)	Békéscsaba (Hungary)	6
	Cluj-Napoca (Cluj County, Romania)	7
	Avrig (Sibiu County, Romania) mountain area	2
	Păltiniş (Sibiu County, Romania) Cindrel Mountains	4
	Cluj-Napoca (Cluj County, Romania)	4
	Sibiu (Sibiu County, Romania)	2
<i>Milax rusticus</i> Millet, 1843	Avrig (Sibiu County, Romania) mountain area	12
	Gödesberg Castle, Bad Gödesberg, Bonn (Germany)	2
	Păltiniş (Sibiu County, Romania) Cindrel Mountains	3
	Bălea Lake, Făgăraş Mountains (Romania)	5
<i>Limax maximus maximus</i> Linnaeus, 1758	Cisnădioara (Sibiu County, Romania)	1
<i>Deroceras agreste</i> (Linnaeus, 1758)	Sibiu (Sibiu County, Romania)	4
	Bihor Mountains from Bănişoara to Stănişoara (Sălaj County, Romania)	2
	Someşul Rece River (Romania)	2
<i>Limax bielzii</i> Seibert, 1873	Castle St. Gotthard, north Graz (Austria)	1
<i>Limax cinereoniger</i> Wolf, 1803	Sibiu (Sibiu County, Romania)	1
	Sebeş, Bistra (Alba County, Romania)	6
	Graz (Austria)	1
	Cluj-Napoca (Cluj County, Romania)	1
	Herculane Baths (Caraş-Severin County, Romania)	1
	Bălea Lake, Făgăraş Mountains (Romania)	1
<i>Limax maximus</i> var. <i>unicolor</i> Heynemann	Transylvania	1
<i>Bielzia coerulans</i> (M. Bielz, 1851)	Cârţişoara (Sibiu, County, Romania)	1
	Bihor Mountains from Bănişoara to Stănişoara (Sălaj County, Romania)	1
<i>Arion hortensis</i> Férussac, 1819	Transylvania	2

HELICIDS IN LAND SNAIL COMMUNITIES FROM SECONDARY GRASSLANDS OF SOUTHERN TRANSYLVANIA

Voichița GHEOCA*

Abstract. Terrestrial land snail communities from secondary grasslands and the importance of helicid snails in these communities was analyzed from 120 quantitative samples taken in four grasslands. The main goal of the study was to analyze the qualitative structure of the community as well as the degree of association between land snail species and their affinity for the habitat. A number of 14 species were identified. The representative species for these communities are *V. excentrica*, *C. tridens* and *C. vindobonensis*. Helicids are an important component regarding number of individuals and the major component from the perspective of biomass.

Keywords: helicids, land snails communities, secondary grassland fauna, Transylvania.

Rezumat. Lucrarea prezintă comunitățile de gastropode terestre caracteristice pajiștilor secundare și rolul helicidelor în cadrul acestora, prin analiza unui număr de 120 de probe cantitative prelevate în patru pajiști. A fost testată structura calitativă a comunităților, gradul de asociere dintre specii precum și gradul de afinitate pentru habitat. Au fost identificate 14 specii de gastropode. Speciile reprezentative pentru acest tip de pajiști sunt *V. excentrica*, *C. tridens*, *C. vindobonensis*, helicidele fiind o componentă importantă din punct de vedere numeric și componenta dominantă din punctul de vedere al biomasei.

Cuvinte cheie: helicide, comunități de gastropode terestre, fauna din pajiști secundare, Transilvania.

Introduction

In Transylvania most of the Subcarpathian hills (400-700 m altitude) were originally covered with forest. The consequence of deforestation was a replacement of primary vegetation with secondary grasslands, i.e. xeromezophyllous herbaceous vegetation. Depending on the slope, their xerophyllous character is more or less striking. These grasslands were originally vegetal associations like *Festucetum rupicolae* and *Agrostio tenuis* – *Festucetum rupicolae*, with a Central – European toward continental, character. Some of the characteristic species are: *Festuca rupicola*, *Asperula cianachica*, *Gallium mollugo*, *Scabiosa ochroleuca*, *Salvia pratensis*, but also mezophyllous species, weeds, and in micro depressions there can be found even hygrophyllous coenoses, with *Phragmites australis* (Fig.1).

Generally, these are more or less fragmented grasslands, limited by clumps of trees, shrubs or cultivated lands. Unlike the calcareous grasslands, which are known to have an extraordinary high diversity of plants and invertebrates, including land snails (Baur *et al.* 2007, Kirby 2001, Wallis De Vries *et al.* 2002), the land snail fauna of these

number of species as in number of individuals. These kinds of grasslands have restrictive conditions for most land snail species, among them are the aridity and the vegetation. Most of the land snail species cannot live in dry conditions (Cameron 1970), as for the vegetation the dominance of Poacea in grasslands limits the food resource for herbivorous snails, which are known to not prefer grass species due to the high silica and fiber content of the leaves (Iglesias, Castillejo 1999), and to their toughness (Dirzo 1980, Speiser, Rowell-Rahier 1991, Boschi, Baur 2007, Rathcke 1985). Although it was proved that poacea are part of their food, even if they are not preferred (Chevalier 2001, Silva, Teresa 1992).

The present paper will set forth an analysis of land snail community of grasslands from Sibiu and Braşov counties in southern Transylvania.

Methods

Qualitative samples of 0.625 cm² (25x25 cm) and 1 m² for the large species, were taken from four grasslands located in southern Transylvania in the interval 1995-2011. The location of sampling points is presented in figure 2. In each of four stations an equal number of 30 samples were taken, all the snails were collected from the grass

* Lucian Blaga University of Sibiu, Faculty of Sciences, vgheoca@yahoo.com

and soil, and a soil sample was also taken on 10 cm dept. The soil was sieved in laboratory and the biological material conserved. The identification was made according to Grossu (1981, 1983 and 1987), Kerney, Cameron (1979) and Bank (2011).

In order to describe the qualitative structure of land snail communities the abundance (A), the frequency (F) and the ecological significance index (Dzuba, $W = (A \times F) / 100$) were used. The quantification of the degree of association between land snail species, was made using CCM contingency coefficient, Cole coefficient (C) for interspecific association and Fager affinity index (I_{AB}). The χ^2 and t tests were applied in order to analyze if the species are statistically significantly related to each other at 5% level of probability. The snail species' affinity for the habitat was tested with Fager affinity index.

Results and discussion

A number of 14 land snail species, from 6 families were identified (table 1). The number of species in each station ranges between 6 and 10. The registered density was ranged between 8 ind./m² and 24 ind./m², while the biomass was 5.18 g/m² – 14.25 g/m²

Analysing the relative abundance values for the 14 species the most abundant species are *Vallonia excentrica*, *Chondrula tridens*, *Cochlicopa lubricella* and *Cepaea vindobonensis*.

Regarding the frequencies, *C. vindobonensis* seem to be the most frequent, but elevated frequency values have also *C. tridens*, *C. lubricella*, *V. excentrica*. The ecologic significance index, register high values in all the sampling stations for *Vallonia excentrica*, *Chondrula tridens*, *Cepaea vindobonensis*.

Considering the values of relative abundance, frequency and ecologic significance, we can conclude that in this grasslands there is a community of xerophylous land snails with tree outstanding species *Vallonia excentrica*, *Chondrula tridens* and *Cepaea vindobonensis*.

The species' distribution on families considering the abundance of each species (Fig. 2) is dominated by Valoniidae (38.28%). High values have also Helicidae (18.04%), Enidae (16.8%), Cochlicopodae (12.91%), Hygromiidae (9.85), Ferussaciidae (2.11%) and Pupiliidae (1.94%).

The Hygromiids are represented by two species *Monacha cartusiana* and *Xerolenta obvia*, and the

helicids by three species *Cepaea vindobonensis*, *Helix pomatia* and *Helix lutescens*

The biomass does not observe the numeric model due to the difference in size between the representatives of different families. So in biomass the helicids (66.79 %), followed by Enidae (22.11%), and Hygromiidae (9.17%) (Fig. 3) are dominant.

For helicids, the three dominant species are the same, but the size of the two species of the *Helix* genus (*H. pomatia* and *H. lutescens*), is reflected in an increased biomass, and as a result a more equilibrated distribution between the three species (Fig. 4)

Let us mention that the presence of *H. pomatia* and *H. lutescens* is dependent on the existence of trees in these grasslands' vicinity.

For the association degree between the terrestrial gastropod species analyze, only eight species, the most common, were selected: *Cochlicopa lubricella*, *Vallonia excentrica*, *Chondrula tridens*, *Ceciloides acicula*, *Monacha cartusiana*, *Xerolenta obvia*, *Cepaea vindobonensis*, *Helix lutescens*. The Cole coefficient indicate significant association ($p=0.05$) between *Cochlicopa lubricella* and *Vallonia excentrica* ($\chi^2 = 11.409$; CCM = 0.374; C= 0.87 ± 0.187).

We have also compiled a matrix of co-occurrences, and Fager's index of affinity was calculated (table 2). The highest level of association was found between *Vallonia excentrica* and *Chondrula tridens*, and these two species and *Cepaea vindobonensis* (Fig. 5).

Conclusions

The land snail community from secondary grasslands is relatively poor and made up of species characteristic of dry habitats, which are also present in calcareous grasslands as *V. excentrica*, *C. tridens*, *Granaria frumentum*, species that feed on fungi and debris, but do not form large populations here as in the conditions of an increased amount of calcium carbonate. Specific for these nutrient rich grasslands is the presence of other species that depend on the neighbouring trees and shrubs. The phytophagous species become essential elements in these communities, as is the case of *C. vindobonensis*, or occur more or less frequently (*H. lutescens* and *H. pomatia*), but enough, due to their dimension, to represent essential items in the nutrient cycle and food chains of these habitats.

Acknowledgements

We would like to thank prof. dr. Constantin Drăgulescu, for his help in the description of the vegetation of the sampling areas.

This work was co-financed from the European Social Found through Sectoral Operational

Programme Human Resources Development 2007-2013, project number POSDRU/89/1.5/S/63258 "Postdoctoral school for zootechnical biodiversity and food biotechnology based on the eco-economy and the bioeconomy required by ecosanogenesis"

REFERENCES

- Bank 2011 Bank Ruud, *Fauna Europaea: Mollusca, Gastropoda*. In: Fauna Europaea version 2.4 (2011) <http://www.faunaeur.org> (Accessed, March 2012).
- Baur *et al.* 2007 Baur Baur, Cremene Cristina, Groza Gheorghe, Schileyko Anatoli, Baur Anette, Erhardt Andreas, *Intensified grazing affects endemic plant and gastropod diversity in alpine grasslands of the Southern Carpathian mountains (Romania)*. In: *Biologia* 62, (2007), p. 438-445.
- Boschi, Baur 2007 Boschi Cristina, Baur Bruno, *The effect of horse, cattle and sheep grazing on the diversity and abundance of landsnails in nutrient-poor calcareous grasslands*. In: *Basic and Applied Ecology* 8(1), (2007), p. 55-65.
- Chevalier *et al.* 2001 Chevalier Laure, Desbuquois Christophe, Le Lannic Joseph, Charrier Maryvonne, *Les Poacées dans le régime alimentaire de l'escargot Helix aspersa Müller (Gastropoda, Pulmonata)*. In: *Comptes Rendus de l'Académie des Sciences - Series III - Sciences de la Vie* 324(11), (2001), p. 979-987.
- Dirzo 1980 Dirzo Rodolfo, *Experimental studies on slug-plant interactions. I. The acceptability of thirty plant species to the slug Agriolimax caeruleus*. In: *J. Ecol.* 68, (1980), p. 981-998.
- Grossu 1981 Grossu Alexandru Victor, *Gastropoda Romaniae. Vol. 3, Ordo Stylommatophora: Suprafamilii Clausiliacea și Achatinacea*. In: *Editura Litera, București* (1981).
- Grossu 1983 Grossu Alexandru Victor, *Gastropoda Romaniae, Vol. 4, Ordo Stylommatophora, Suprafamilii: Arionacea, Zonitacea, Ariophantacea și Helicacea*. In: *Editura Litera, București* (1983).
- Grossu 1987 Grossu Alexandru Victor, *Gastropoda Romaniae. Vol. 2. Ordo Basommatophora, Ordo Stylommatophora, Suprafam. Succinacea, Cochlicopacea, Pupillacea*. In: *Editura Litera, București* (1987).
- Iglesias, Castillejo 1999 Iglesias Javier, Castillejo Jose, 1999, *Field observations on the land snail Helix aspersa Muller*. In: *J. Mol. Stud.* 65, (1999), p. 411-423.
- Kerney, Cameron 1979 Kerney Michael, Cameron Robert, *A field guide to the Land Snails of Britain and North-West Europe*. Collins, London (1979).
- Kirby 2001 Kirby Peter, *Habitat Management for Invertebrates: a Practical Handbook*. In: *Royal Society for the Protection of Birds, Bedfordshire* (2001).
- Rathcke 1985 Rathcke, Beverly, *Slugs as Generalist Herbivores: Tests of Three Hypotheses on Plant Choices*. In: *Ecology* 66(3), (1985), p. 828-836.
- Silva, Teresa 1992 Silva Oliveira, Teresa Maria, *Effects of mollusc grazing on the development of grassland species*. In: *Journal of Vegetation Science* 3(2), (1992), p. 267-270.
- Speiser, Rowell-Rahier 1991 Speiser Bernhard, Rowell-Rahier Martine, *Effect of food availability, nutritional value and alkaloids on food choice in the generalist herbivore Arianta arbustorum (Gastropoda: Helicidae)*. In: *Oikos* 62, (1991), p. 306-318.
- Wallis DeVries *et al.* 2002 Wallis DeVries Michiel, Poschlod Peter, Willems Jo, *Challenges for the conservation of calcareous grasslands in Northwestern Europe: integrating the requirements of flora and fauna*. In: *Biological Conservation* 104, (2002), p. 265-273.

LIST OF ILLUSTRATIONS

- Fig. 1.** Weeded grassland – Nou Român
Fig. 2. Location of sampling stations
Fig. 3. Relative abundance of terrestrial gastropod families from grasslands; the percentage of different helicid species within the family
Fig. 4. Biomass of terrestrial gastropod families from grasslands; the percentage of different helicid species within the family
Fig. 5. Tree diagram of degree of cenotic affinity for the land snail species from grasslands (using the values of Fager affinity index)
The species' codification :
1- *Cochlicopa lubricella*, 2 – *Vallonia excentrica*, 3 – *Chondrula tridens*, 4 – *Cecilioides acicula*, 5 – *Monacha cartusiana*, 6 – *Xerolenta obvia*, 7 – *Cepaea vindobonensis*, 8 – *Helix lutescens*.
- Tab. 1.** List of land snails from grasslands: D – density (ind/m²), A – relative abundance(%), F – frequency (%), W – Dzuba ecologic significance index; S* species with only shells.
Tab. 2. Matrix of the Fager affinity index values for the land snail species from grasslands (with hyphen are marked pairs of species for which the value of index is not significant).
The species' codification :
1- *Cochlicopa lubricella*, 2 – *Vallonia excentrica*, 3 – *Chondrula tridens*, 4 – *Cecilioides acicula*, 5 – *Monacha cartusiana*, 6 – *Xerolenta obvia*, 7 – *Cepaea vindobonensis*, 8 – *Helix lutescens*

LISTA ILUSTRAȚIILOR

- Fig. 1.** Pajiște îmburuientă – Nou Român
Fig. 2. Localizarea stațiilor de colectare
Fig. 3. Abundența relativă a familiilor de gastropode terestre din pajiști; ponderea diferitelor specii de helicide în cadrul familiei
Fig. 4. Ponderea biomasei familiilor de gastropode terestre din pajiști; ponderea diferitelor specii de helicide în cadrul familiei
Fig. 5. Matricea valorilor indicelui de afinitate Fager (cu liniuță sunt marcate perechile de specii pentru care valorile indicelui sunt nesemnificative)
Codificarea speciilor este următoarea :
1- *Cochlicopa lubricella*, 2 – *Vallonia excentrica*, 3 – *Chondrula tridens*, 4 – *Cecilioides acicula*, 5 – *Monacha cartusiana*, 6 – *Xerolenta obvia*, 7 – *Cepaea vindobonensis*, 8 – *Helix lutescens*.
- Tab. 1.** Lista speciilor de gastropode terestre din pajiști: D – densitatea (ind/m²), A – abundența relativă (%), F – frecvența (%), W – indicele de semnificație ecologică Dzuba; S* specii prezente doar sub formă de cochilii.
Tab. 2. Valorile indicelui de afinitate Fager pentru speciile de gastropode terestre din pajiști (cu liniuță sunt marcate perechile de specii pentru care valoarea indicelui Fager este nesemnificativă)
Codificarea speciilor este următoarea :
1- *Cochlicopa lubricella*, 2 – *Vallonia excentrica*, 3 – *Chondrula tridens*, 4 – *Cecilioides acicula*, 5 – *Monacha cartusiana*, 6 – *Xerolenta obvia*, 7 – *Cepaea vindobonensis*, 8 – *Helix lutescens*.



Fig. 1. Grassland with weeds in Southern Transylvania – Nou Român




Fig. 2. Location of the four sampling sites




Fig. 3. Relative abundance of terrestrial gastropod families from grasslands; the percentage of different helicid species within the family




Fig. 4. Biomass of terrestrial gastropod families from grasslands; the percentage of different helicid species within the family




Fig. 5. Tree diagram of degree of cenotic affinity for the land snail species from grasslands (using the values of Fager affinity index).

The species' codification: 1- *Cochlicopa lubricella*, 2 – *Vallonia excentrica*, 3 – *Chondrula tridens*, 4 – *Cecilioides acicula*, 5 – *Monacha cartusiana*, 6 – *Xerolenta obvia*, 7– *Cepaea vindobonensis*, 8 – *Helix lutescens*.

Tab. 1. List of land snails from grasslands

(D – density (ind/m²), A – relative abundance(%), F – frequency (%), W – Dzuba ecologic significance index, S* species with only shells.

Species	Șura Mare				Tâlmăciu			
	D	A	F	W	D	A	F	W
Cochlicopidae								
<i>Cochlicopa lubricella</i> (Porro, 1838)	3.25	13.9	36.36	5.05	-	-	-	-
<i>Cochlicopa lubrica</i> (Müller, 1774)	-	-	-	-	-	-	-	-
Valloniidae								
<i>Vallonia pulchella</i> (Müller, 1774)	-	-	-	-	-	-	-	-
<i>Vallonia costata</i> (Müller, 1774)	0.5	2.15	18.18	0.39				
<i>Vallonia excentrica</i> (Sterki, 1892)	10.75	46.23	81.8	13.27	2	25.8	36.36	9.38
Pupillidae								
<i>Pupilla muscorum</i> (Linnaeus, 1758)	0.25	1.07	9.05	0.09	-	-	-	-
Chondrinidae								
<i>Granaria frumentum</i> (Draparnaud, 1801)	-	-	-	-	0.5	6.45	18.18	1.17
Enidae								
<i>Chondrula tridens</i> (Müller, 1774)	4.5	19.35	45.45	8.79	,	12.9	27.27	3.51
Ferussaciidae								
<i>Cecilioides acicula</i> (Müller, 1774)	1.75	7.52	54.54	4.10	-	-	-	-

Hygromiidae								
<i>Xerolenta obvia</i> (Menke 1828)	-	-	-	-	-	-	-	-
<i>Monacha cartusiana</i> (Müller, 1774)	0.5	2.15	18.18	0.39	0.75	9.67	18.18	1.75
Helicidae								
<i>Cepaea vindobonensis</i> (Pfeiffer, 1828)	1.75	7.52	63.63	4.78	3	38.7	81.81	38.66
<i>Helix pomatia</i> (Linnaeus, 1758)	-	-	-	-	0.5	6.45	18.18	1.17
<i>Helix lutescens</i> (Rossmässler, 1837)	-	-	-	-	-	-	-	-
	Nou Român				Făgăraș			
Species	D	A	F	W	D	A	F	W
Cochlicopidae								
<i>Cochlicopa lubricella</i> (Porro, 1838)	6.88	29	77.77	22.55	0.5	2.39	22.22	0.53
<i>Cochlicopa lubrica</i> (Müller, 1774)	1.55	6.37	22.22	1.46	-	-	-	-
Valloniidae								
<i>Vallonia pulchella</i> (Müller, 1774)	1.33	5.63	11.15	0.62	-	-	-	-
<i>Vallonia costata</i> (Müller, 1774)	S*				0.24	1.15	11.11	0.12
<i>Vallonia excentrica</i> (Sterki, 1892)	6.88	29	88.88	25.77	9	43.18	88.88	38.38
Pupillidae								
<i>Pupilla muscorum</i> (Linnaeus, 1758)	-	-	-	-	1.4	6.71	44.45	2.98
Chondrinidae								
<i>Granaria frumentum</i> (Draparnaud, 1801)	-	-	-	-	-	-	-	-
Enidae								
<i>Chondrula tridens</i> (Müller, 1774)	5.22	22.56	77.77	17.16	1.25	5.99	33.33	1.99
Ferussaciidae								
<i>Cecilioides acicula</i> (Müller, 1774)	0.22	0.93	11.15	0.1	-	-	-	-
Hygromiidae								
<i>Xerolenta obvia</i> (Menke 1828)	-	-	-	-	3	14.39	77.77	11.19
<i>Monacha cartusiana</i> (Müller, 1774)	-	-	-	-	2.75	13.19	77.77	10.26
Helicidae								
<i>Cepaea vindobonensis</i> (Pfeiffer, 1828)	1.1	4.69	66.66	3.12	1.5	7.19	55.6	4.00
<i>Helix pomatia</i> (Linnaeus, 1758)	-	-	-	-	0.2	0.95	11.11	0.106
<i>Helix lutescens</i> (Rossmässler, 1837)	0.44	1.87	33.33	0.62	1	4.79	22.22	1.06

Tab. 2. Matrix of the Fager affinity index values for the land snail species from grasslands
(with hyphen are marked pairs of species for which the value of index is not significant)

	1	2	3	4	5	6	7	8
1	1							
2	0.603	1						
3	-	0.686	1					
4	0.2	0.226	-	1				
5	0.162	0.3	0.182	-	1			
6	0.25	0.182	0.154	0.091	0.138	1		
7	-	0.682	-	0.077	0.068	0.259	1	
8	-	0.157	-	-	0.08	0.2	0.16	1

CONSIDERATIONS REGARDING LAND SNAILS DIVERSITY IN URBAN AREAS. CASE STUDY - THE CITY OF SIBIU

Voichița GHEOCA*

Abstract. During the past two decades the city of Sibiu developed in two directions, by extension of the inhabited surface in the suburban area and by construction of new buildings on nearly every available land patch. The consequence was the reduction of the green zones and the supplementary habitat fragmentation, process which is characteristic in urban areas. The present study aims to assess the terrestrial gastropod diversity in urban areas with different destinations by qualitative sampling in 35 points located in seven different areas of the city. A number of 39 species of terrestrial gastropod species were identified. Except Dumbrava Forest and Sub Arini Park, which still preserve conditions for these animals' development, the city presents very low diversity of terrestrial gastropods (between 2 and 9 species per station), underlying the importance of the conservation of areas with suitable conditions. We mention the presence in Dumbrava Forest of *Chilostoma banaticum*, species of community interest from Annex II of the EU Habitats Directive.

Keywords: land snail communities, biodiversity, urban areas, Sibiu, Transylvania

Rezumat. În ultimele două decenii, orașul Sibiu s-a dezvoltat pe două direcții, prin extinderea suprafeței construite în zona periurbană și prin construirea de clădiri pe practic orice suprafață disponibilă din perimetrul intravilan. Rezultatul a fost o diminuare drastică a zonelor verzi, accentuând și mai mult fragmentarea habitatelor, caracteristică zonelor urbane. Lucrarea urmărește diversitatea gastropodelor terestre din perimetrul urban și periurban al orașului Sibiu, prin evaluarea cantitativă a faunei de gastropode din 35 de puncte localizate în șapte zone ale orașului. Au fost identificate un număr de 39 de specii de gastropode terestre, și exceptând Pădurea Dumbrava și Parcul Sub Arini, care mai conservă condiții pentru dezvoltarea acestor animale, în restul zonelor investigate diversitatea este extrem de scăzută, fiind cuprinsă între 2 și 9 specii, subliniind importanța conservării spațiilor în care se mai păstrează condiții favorabile prezenței acestor animale. Menționăm prezența speciei *Chilostoma banaticum* în Pădurea Dumbrava, specie de interes comunitar din Anexa II a Directivei Habitats.

Cuvinte cheie: gastropode terestre, comunități, biodiversitate, zone urbane, Sibiu, Transilvania

Introduction

Often underestimated the analysis of terrestrial ecosystems biodiversity, gastropods represent a group whose importance is given by their contribution as a food source for a wide variety of animals, from small mammals to birds, reptiles and amphibians, and also as an important source of calcium for some animals, especially birds. The shells content in calcium carbonate is a critical supplement for the female birds or both parents during the raise of young (Kerney, Cameron 1979).

Recently, as new data on these animals' efficiency as bioindicators accumulate, the interest regarding terrestrial gastropods increases. There are a series of studies using land snail communities as models for different disturbances as anthropic activity and habitat fragmentation (Frest 2002, Dedov, Penev

2004, Gotmark *et al.* 2008, Kappes *et al.* 2009), while other studies use land snail species as accumulation indicators (Rabitsch 1996, Regoli *et al.* 2006).

The interest towards land snails from human settlements was mainly related to their effect as pests for many cultivates species (Baker 2002; Sanderson, Sirgel 2002). Only in the last years came into attention the loss of biodiversity in urban areas. Urbanization was shown to have a strong influence on biodiversity, several studies dealing with the effect of urbanization on the diversity of different groups (Gilbert 1989, Dedov, Penev 2000, Fernández-Juricic 2002, Smith *et al.* 2006a, 2006b, Knapp *et al.* 2008) some of them considering land snails (Tappert 1995, Horsák *et al.* 2009).

In the past two decades the city of Sibiu developed in two directions, by extension of the inhabited surface in the suburban area, and by construction of new buildings on nearly every available land

* Lucian Blaga University of Sibiu, Faculty of Sciences, vgheoca@yahoo.com

patch, with a consequent reduction of the green zones (especially private gardens). Sometimes the buildings and parking places represents in this newly build spaces over 80-90% of the entire surface, and the loss of natural and semi natural habitats is alarming. The present study aims to assess the diversity of terrestrial gastropod fauna from Sibiu, in urban areas with different destinations.

Material and methods

The study was conducted over six years (2007-2012), qualitative samples being taken from 35 points in seven locations which differ by the degree of human intervention: Dumbrava forest, at the western limit of the city, the main park, which is actually an extension of the forest in the city, the city center, one industrial area and three different residential areas. Fig.1 presents the location of the investigated areas.

Results and discussions

Habitat quality and terrestrial gastropod fauna

39 species of terrestrial gastropods belonging to 16 families were found in the investigated areas from Sibiu. The highest number of species was identified in the largest park (Sub Arini) of the city (25 species) and in Dumbrava Forest (24 species). 13 species are common for the two areas and some were found only in these two areas. Among them are *Clausilia dubia*, *Vestia elata* and *Aegopinella epipedostoma*. Another area that preserves some diversity of snail fauna is represented by the banks of Cibin River, where some patches of willows were left. In the rest of the city the diversity of land snail fauna is much lower, only 2 to 7 species being found. Tab. 1 presents the list of land snail species and the number of species in each area.

Analyzing the preferences for humidity of the identified gastropod species, 64% are mesophylous and 13% are mesoxerophylous, due to the absence of humid habitats from the city (Fig. 2).

These results emphasize that public parks and gardens are refuges for biodiversity in urban areas, where the disappearance of private gardens limit the distribution of most animal species. The natural origin of the central park is emphasized by the presence of tree species from the former forest, as *Alnus glutinosa*, *Quercus robur*, *Carpinus betulus*, *Fraxinus excelsior*. Along these native species there are ornamental trees, planted beginning with the 19th Century. Created along the rivulet,

consisting of an alder belt flanked by oak forest, the park is a continuation of Dumbrava Forest.

The faunistic composition of the park reflects partially the forests' fauna, especially in mobile groups, for which the interruption of the landscapes' continuity between the two elements, the forest and the park, by the residential area and associated facilities, is not an impediment for dispersal. Other groups, as it is the case of terrestrial gastropods, cannot surpass the barrier and local populations are developing in these fragmented habitats. Thus, their survival depends primarily on the preservation of suitable local conditions.

Land snail species of special interest

Among the land snail species *Drobacia banatica*, a Quaternary Carpathian relict, whose eastern distribution limit is Romania and Ukraine, requires special interest. For some invertebrate species, like land snails, the Carpathian mountain chain contains a significant number of endemic species, and can be considered the center of surviving and autochthonous postglacial evolution of some species with low mobility (Varga 2001). This is the case of *D. banatica*, species from Annex II of EU Habitats Directive. Its presence in Dumbrava Forest is a supplementary argument for the conservation of this area, and against the extension of residential buildings into the forest.

Another remarkable element is the presence of *Cepaea hortensis* (Fig 3) almost everywhere in Sibiu, in public parks and gardens, as well as in private gardens. *C. hortensis*, the smallest of the four species of the *Cepaea* genus, was mentioned for the first time in Romania by Csiki (1918), near Oradea, but its presence there was not confirmed later (Grossu 1983). In 1999 the species was identified in Sibiu (Gheoca 2004), its presence is considered the consequence of a deliberate introduction of this species, based on the shells from the Kimakowicz malacological collection from Sibiu Natural History Museum. This collection contains shells of *C. hortensis* originating from the same garden in Sibiu, with the mention Manheim on the labels. The population was apparently surveyed during the period 1911 – 1934.

C. hortensis is present in moister habitats than *C. vindobonensis*, and generally the two species are not found together. However, in one of the sampling areas they were found together. The populations from Sibiu are equally homomorphic and polymorphic, but the polymorphism was lower than known for this species.

Is possible that all the populations from the city area originated from a single founder population but their polymorphism is lower than that of the population studied by Kimakowicz at the beginning of the 20th Century. A later accidental introduction of this species in Sibiu is also not excluded (Gheoca 2004).

Whichever the species' provenience, its presence in Sibiu in the past Century is certain. In the condition of urban development and reduction of green spaces, its future in this isolated enclave is uncertain and dependent on the quality of suitable habitats.

Another mentionable species is *Lozekia transsilvanica*, one of two species of the genus *Lozekia*, genus with a relatively restricted distribution in the Carpathian-Pannonian region (Fehér *et al.* 2009). The type locality for *L. transsilvanica* is Sibiu, Dumbrava Forest, and its conservation in this habitat is important in the context of disputes over the intent of residential development in the area.

Conclusions

The diversity of land snail fauna from Sibiu is reduced and almost entirely dependent on the presence of the forest neighbouring the city and of the emergent park.

The conservation of these two habitats is essential for the local biodiversity, including terrestrial gastropods, but as demonstrated by Götmark *et al.*

(2008), the microhabitat factors are not the only responsible for species richness and composition in land mollusks. These sedentary organisms seem to be substantially influenced by the surrounding landscape, which should be considered in conservation work and in future urbanistic plans. Habitat fragmentation is also a major cause for species loss and its effect on invertebrates with low active capacity of dispersal, as it is the case of land snails, is even more accentuated. The response to fragmentation seems to be specific, as demonstrated by Kappes *et al.* (2009), and to depend on habitat specialization and macroclimatic conditions. This is especially important in case of protected species like *Drobacia banatica*, whose management has to be a priority of conservation works.

For some species the exploitation can also be a factor of numerical reduction, as it is the case of edible snail *Helix pomatia*, a species collected from urban and rural areas, but also from natural habitats.

Acknowledgements

This work was co-financed from the European Social Found through Sectoral Operational Programme Human Resources Development 2007-2013, project number POSDRU/89/1.5/S/63258 "Postdoctoral school for zootechnical biodiversity and food biotechnology based on the eco-economy and the bio-economy required by eco-san-genesys"

REFERENCES

- Barker 2002 Barker Glen, (eds) *Molluscs as crop pests*. In: CABI Publishing, Wallingford (2002).
- Dedov, Penev 2000 Dedov Ivailo, Penev Lyubomir, *Species composition and origins of the terrestrial gastropod fauna of Sofia City, Bulgaria*. In: *Ruthenica* 10, Moscow (2000), p. 121–131.
- Dedov, Penev 2004 Dedov Ivailo, Penev Lyubomir, *Spatial variation in terrestrial gastropod communities (Gastropoda, Pulmonata) along urban-rural gradients in Sofia City, Bulgaria*. In: Penev L, Niemelä J, Kotze J, Chipev N (eds) *Ecology of the city of Sofia. Species and communities in an urban environment*. Pensoft Publishers, Moscow (2004), p. 307–318.
- Fehér et al. 2009 Fehér Zoltán, Varga András, Deli Tamás and Domokos Tamás, *Geographic distribution and genital morphology of the genera Lozekia Hudec, 1970 and Kovacsia Nordsieck, 1993 (Mollusca, Gastropoda, Hygromiidae)*. In: *Zoosyst. Evol.* 85 (1), (2009), p. 151–160.
- Gheoca 2004 Gheoca Voichița, *Data concerning the presence of Cepaea hortensis Müller, 1774 (Gastropoda, Helicidae) in Romania*. In: *Studii și comunicări, Revista muzeului de Istorie Naturala Sibiu* 29, Sibiu (2004), p. 101–106.
- Grossu 1983 Grossu Alexandru Victor *Gastropoda Romaniae. Ordo Stylommatophora, vol.4, Suprafam. Arionacea, Zonitacea, Ariophantacea și Helicacea*, Ed Litera (1983), București.
- Götmark et al. 2008 Götmark Frank, Von Proschwitz Ted, Franc, Niklas, *Are small sedentary species affected by habitat fragmentation? Local vs. landscape factors predicting species richness and composition of land molluscs in Swedish conservation forests*. In: *Journal of Biogeography* 35(6), (2008), p. 1062–1076.
- Fernández-Juricic 2002 Fernández-Juricic Esetban, *Can human disturbances promote nestedness? A case study with breeding birds in urban habitat fragments*. In: *Oecologia* 131, (2002), p. 269–278.
- Frest 2002 Frest Terrence, *Native snails: indicators of ecosystem health*. In: Wuerthner G, Matteson M (eds) *Welfare ranching, the subsidized destruction of the American West*. Island Press, Washington DC (2002), p. 211–215.
- Gilbert 1989 Gilbert Oliver, *The ecology of urban habitats*. In: Chapman and Hall, London (1989).
- Horsák et al. 2009 Horsák Michal, Juříčková Lucie., Kintrová Katerina, Hájek Ondřej, *Patterns of land snail diversity over a gradient of habitat degradation: a comparison of three Czech cities*, In: *Biodiversity and Conservation* 18(13), (2009), p. 3453–3466.
- Kappes et al. 2009 Kappes Heike, Jordaens Kurt, Hendrickx Frederik, Maelfait Jean-Pierre, Lens Luc Backeljau Thierry, *Response of snails and slugs to fragmentation of lowland forests in NW Germany*. In: *Landscape Ecology* 24, (2009), p. 685–697.
- Kerney, Cameron 1979 Kerney Michael, Cameron Robert, *A field guide to the Land Snails of Britain and North-West Europe*, London (1979).
- Knapp et al. 2008 Knapp Sonja, Kühn Ingolf, Mosbrugger Volker, Klotz Stefan, *Do protected areas in urban and rural landscapes differ in species diversity?* In: *Biodivers Conserv* 17, (2008), p. 1595–1612.
- Rabitsch 1996 Rabitsch Werner, *Metal Accumulation in Terrestrial Pulmonates at a Lead/Zinc Smelter Site in Arnoldstein, Austria*. In: *Bulletin of environmental contamination and toxicology* 56(5), (1996), 734–741.
- Regoli et al. 2006 Regoli Francesco, Gorbi Stefania, Fattorini Daniele, Tedesco Sara, Notti Alessandra, *Use of the Land Snail Helix aspersa as Sentinel Organism for Monitoring Ecotoxicologic Effects of Urban Pollution: An Integrated Approach*. In: *Environ Health Perspect.* 114, (2006), p. 63–69.
- Sanderson, Sirgel 2002 Sanderson Graeme, Sirgel Willem, *Helicidae as pests in Australian and South African grapevines*, In G.M. Barker (eds.), *Molluscs as crop*, CABI Publishing

- New York, (2002), p. 255-270.
- Smith *et al.* 2006a Smith Richard, Warren Philip, Thompson Ken, Gaston Kevin, *Urban domestic gardens (VI): environmental correlates of invertebrate species richness*. In: *Biodivers. Conserv.* 15, (2006), p. 2415–2438.
- Smith *et al.* 2006b Smith Richard. Gaston Kevin. Warren Philip, Thompson Ken, *Urban domestic gardens (VIII): environmental correlates of invertebrate abundance*. In: *Biodivers. Conserv.* 15, (2006), p. 2515–2545.
- Tappert 1995 Tappert Andrea, *Die Molluskenfauna von Köln*. In: *Dacheniana, Beihefte* 35, (1995), p. 579–643.
- Varga 2001 Varga Zoltan, *Post-glacial dispersal strategies of Orthoptera and Lepidoptera in Europe and in the Carpathian basin*. In: *Proc. 13th Int. Coll. EIS*, (2001), p. 93-105.

LIST OF ILLUSTRATIONS

- Fig. 1.** Location of sampling areas
1. Sub Arini - public parc
 2. Dumbrava Forest
 3. Cibin riverside
 4. Industrial area – periphery of the town
 5. Guterita Hill
 6. Garden in Terezian residential area
 7. Turnișor residential area
 8. Tineretului Park
 9. Ceferiștilor residential area
- Fig. 2.** The number of land snail species for each city area
- Fig. 3.** Distribution of land snail species considering their demand towards humidity
- H – hygrophylous
MH – mesohygrophylous
M – mesophylous
MX – mesoxerophylous
- Fig. 4.** Some individuals of *Cepaea hortensis* from Sibiu (foto V. Gheoca).
- Tab. 1.** List of the land snail species from the nine analyzed areas of Sibiu city

LISTA ILUSTRAȚIILOR

- Fig. 1.** Localizarea punctelor de colectare
1. Parcul Sub Arini
 2. Pădurea Dumbrava
 3. Malul Râului Cibin
 4. Zonă industrială – periferia oraului
 5. Dealul Guterita
 6. Grădină în cartierul Terezian
 7. Zonă verde în cartierul Turnișor
 8. Parcul Tineretului
 9. Grădină în cartierul Ceferiștilor
- Fig. 2.** Numărul speciilor de gastropode terestre identificate în fiecare zonă
- Fig. 3.** Distribuția speciilor de gastropode terestre în funcție de cerințele lor față de umiditate
- H – higrofile
MH – mezohigrofile
M – mezofile
MX – mezoxerofile
- Fig. 4.** Indivizi de *Cepaea hortensis* din Sibiu (foto V. Gheoca).
- Tab.1.** Lista speciilor de gastropode terestre identificate în cele nouă zone analizate ale orașului Sibiu.



Fig. 1. Location of sampling areas




Fig. 2. Distribution of land snail species considering their demand towards humidity



Fig. 3. Some individuals of *Cepaea hortensis* from Sibiu (foto V. Gheoca).

Tab. 1. List of the land snail species from the nine analyzed areas of Sibiu city.

<i>Species</i>	1. Sub Arini public parc	2. Dumbrava forest	3. Cibin river	4. Periphery of the town	5. Gusterita Hill	6. Garden Terezia	7. Turnisor resid. area	8. Tineretului park	9. Ceferistilor residential area
<i>Carichium tridentatum</i> (Risso, 1826)			X						
<i>Vallonia costata</i> (Muller, 1774)							X		
<i>Vallonia pulchella</i> (Muller, 1774)						X			
<i>Succinea putris</i> (Linnaeus, 1758)	X	X	X						
<i>Cochlicopa lubrica</i> (Muller, 1774)	X	X							
<i>Chondrula tridens</i> (Küster, 1852)				X					
<i>Columella edentula</i> (Draparnaud, 1805)		X							
<i>Cochlodina laminata</i> (Montagu, 1803)	X								
<i>Ruthenica filograna</i> (Rossmässler, 1836)	X								
<i>Clausilia dubia</i> (Draparnaud, 1805)	X	X							
<i>Vestia elata</i> (Rossmässler, 1836)	X	X							
<i>Alinda biplicata</i> (Montagu, 1803)		X							
<i>Cecilioides acicula</i> (Müller, 1774)						X			
<i>Punctum pygmaeum</i> (Draparnaud, 1801)		X							
<i>Vitraea transsylvanica</i> (Clessin, 1877)		X							
<i>Vitrea cristallina</i> (Müller, 1774)		X							
<i>Euconulus fulvus</i> (Muller, 1774)		X							
<i>Zonitoides nitidus</i> (Muller, 1774)	X	X	X						
<i>Oxychillus draparnaudi</i> (Beck 1837)	X	X		X				X	X
<i>Oxychillus depressus</i> (Sterki, 1880)		X							X
<i>Aegopinella epipedostoma</i> (Fagot, 1879)	X	X							
<i>Aegopinella minor</i> (Stabile, 1864)	X				X				
<i>Vitrina pellucida</i> (Müller 1774)	X								
<i>Limax maximus</i> (Linnaeus, 1758)	X		X						X
<i>Limax cinereoniger</i> (Wolf, 1803)		X							
<i>Deroceras reticulatum</i> (Müller, 1774)			X						X
<i>Arion subfuscus</i> (Draparnaud, 1805)		X	X						
<i>Arion circumscriptus</i> (Johnston, 1828)	X	X	X						X
<i>Arion hortensis</i> (Férussac, 1819)	X						X		
<i>Fruticicola fruticum</i> (Müller 1774)	X	X	X	X			X		X
<i>Euomphalia strigella</i> (Draparnaud, 1801)	X	X	X		X				
<i>Monacha cartusiana</i> (Müller, 1774)	X		X	X	X				
<i>Lozekia transsylvanica</i> (Westerlund, 1876)	X	X							
<i>Pseudotrachia rubiginosa</i> (A. Schmidt 1853)	X	X							
<i>Perforatella bidentata</i> (Gmelin 1788)	X								
<i>Drobacia banatica</i> (Rossmässler, 1838)		X							
<i>Cepaea hortensis</i> (Muller, 1774)		X	X	X		X			X
<i>Cepaea vindobonensis</i> (Férussac, 1821)	X		X	X	X		X	X	
<i>Helix pomatia</i> (Linnaeus, 1758)	X	X	X	X	X	X	X		X
Number of species	25	24	9	7	4	4	5	2	7

THE IMPORTANCE OF HOMOGENEOUS VS. HETEROGENOUS WETLANDS IN RALLID (RALLIDAE) PHENOLOGICAL SEASONS

Alexandru Nicolae STERMIN*
Liviu Răzvan PRIPON*
Alin DAVID*

Abstract. Seven rallid species inhabit the European homogenous and heterogeneous wetlands. Analyzing the presence and absence of rail species in one homogenous (Sic Reed beds) and one heterogeneous (Sucutard Pond) wetland from Romania in Sic Reed beds we found six rail species during the migration season and two during the breeding season while at Sucutard Pond four species were located in both seasons. We conclude that homogeneous reed beds are the typical habitats used by rails during migration as important travel corridors and buffer zones.

Keywords: lifetime seasons, travel corridors, conservation, habitat preference, spatial distribution.

Rezumat. Șapte specii de ralide ocupă zonele umede, omogene sau heterogene din Europa. Analizând prezența sau absența acestor specii în zonele umede omogene (Stufărișurile de la Sic) și în cele heterogene (Heleșteele de la Sucutard) din România am constatat că la Stufărișurile de la Sic au fost prezente șase specii de ralide în timpul migrației de primăvară și două în perioada de reproducere iar pe Heleșteele de la Sucutard patru specii în ambele perioade. Am concluzionat că stufărișurile omogene sunt habitate tipice pentru ralide, fiind folosite de acestea ca importante coridoare de migrație.

Cuvinte cheie: sezoane fenologice, coridoare de migrație, preferința de habitat, distribuție spațială.

Introduction

The optimization theory assumes that adaptive traits of species have evolved to maximize evolutionary success (Brooks, McLennan, 1991; Price *et al.* 1997, Adamik *et al.* 2003). According to this, one may expect species-specific preferences for a particular habitat type as a consequence of a historical relationship between species and their environment (Adamik *et al.* 2003) but, if wetlands in a region are spatially diverse and temporally variable, wildlife may require the use of several different wetlands during a phenological season (Haig *et al.* 1997, Joyal *et al.* 2001, Neugle *et al.* 2001, Roe, Georges 2007).

Originally inhabiting forests, rails are nowadays widely distributed, occupying many habitat types, from forests to wetlands and grasslands, both natural and artificial (Taylor 1998). However, this remains one of the most poorly represented groups in ornithological research (Brambilla, Jenkins 2009). This situation is unfortunate because the

number of extinct species is high in this group: since the 1600s 16 species became extinct from a total of more than 150 species (Robinson, Horsfall 2004). The main threats for rail species are habitat loss (e.g. as a consequence of wetland draining), agricultural intensification and deforestation (Collar *et al.* 1994, Taylor, 1998, Jenkins, Ormerod, 2002).

Seven rallid species (Eurasian Coot *Fulica atra*, Moorhen *Gallinula chloropus*, Water Rail *Rallus aquaticus*, Spotted Crake *Porzana porzana*, Little Crake *Porzana parva*, Baillon's Crake *Porzana pusilla* and Corncrake *Crex crex*) are inhabiting the European wetlands (Taylor 1998). Depending on the plant communities, water level and management, wetland habitats are heterogeneous or homogeneous having the possibility to evolve from one to another. Heterogeneous habitats are conservation priorities, being important not only for generalist but also for specialized species.

In this context, the aims of our study are: (i) to illustrate the difference in rallid species composition regarding the use of both heterogeneous and homogeneous habitats during two phenological seasons (spring migration and breeding season), (ii) to reveal the importance of

* Babeș-Bolyai University, Department of Taxonomy and Ecology, Cluj-Napoca, Romania,
sandu.stermin@yahoo.com.
liviu.pripon@gmail.com
adavid.ubb@gmail.com

each of the two wetland types in the species ecological necessities.

Study areas

The research was carried out in Sucutard Pond (SP) and Sic Reed beds (SR), two wetlands located 13 km apart from each other, in the central part of the Transylvanian Plane, Romania (24°10' E; 46°50' N) (David 2008). SP is an anthropogenic wetland with an area of 46 ha, covered 50% by open water, 15% by Reed (*Phragmites australis*) and 35% by Cattail (15% dominated by *Typha latifolia* and 20% by *Typha angustifolia*) (Fig. 1). The spatial distribution and the structure of different plant communities make this pond a heterogeneous wetland.

With an area of 252.68 ha which is in 98% covered by Reed, SR is the largest reed bed from the Transylvanian Plane being a homogeneous habitat (Stermin *et al.* 2011). It has a length of around 4 km with a maximum compact reed width of 550 m. For our studies we selected a 40 ha area part of SR (Fig. 1), here the density of the vegetation is around 400 plant stems per square meter and the water level does not exceed 1.5 m, generally ranging between 20 and 50 cm (Stermin *et al.* 2011).

Methods

We selected 10 observation points inside each study area (Fig. 1). In SR the points were located 100 m apart from each other, in a transect which crosses the reed bed from one edge to another (Fig. 1). In addition, we selected 10 points inside SP, situated 50 m apart from each other, distributed to cover all habitat types (Fig. 1). Six points were located at different distances to the edge in a habitat dominated by *T. angustifolia*, two points in habitats dominated by *T. latifolia*, one in Reed and one in open water (Fig. 1). Because the habitat dominated by *T. angustifolia* has many different characteristics (located near to the shore, near to the reed or to the open water) we selected six points to record all these specific situations. As the other habitats have more constant characteristics across the SP wetland we selected only one or two points located in each one, according to the habitat surfaces and visibility (in open water only one point was necessary to have a wide view).

To detect the species in spring migration, observations were conducted at each point between 23-26 April 2010 (on SR) and 17-21 April 2011 (on SP) during three days, three times during a day

(6:00-9:00, 13:00-16:00 and 20:00-23:00). We selected the end of April because this is the time when the migration period of all European rallid species overlaps (the end of migration period for Coot, Moorhen, Water Rail and Little Crane and the first part of migration period for Spotted Crane, Baillon's Crane and Corncrake) (Taylor 1998, Cramp 1980). In order to detect the species during the breeding season, we selected one day (19.07.2010 on SR and 15.07.2011 on SP) during which three observation times (6:00-9:00, 13:00-16:00 and 20:00-23:00) were conducted.

Considering that during the migration period the structure of the species communities from a specific habitat is changing continuously, we selected in this period three observation days to record these possible changes. Because during the breeding period the bird communities are more stable we performed the observation only one day, when the weather conditions were favourable.

To detect species presence, in each point, the spontaneous activity of each species was recorded for five minutes, after that we performed the Moorhen's, Water Rail's, Little Crane's, Spotted Crane's, Baillon's Crane and Corncrake's playbacks (from 20 to 30 seconds each one). After each playback, during 30 seconds we noted the presence or absence of a rallid species.

Results

A total of 90 observations, during spring migration and 30 during breeding season were conducted in each wetland. On SR, during the migration period we recorded five rallid species (Moorhen, Water Rail, Little crane, Spotted Crane and Corncrake). We mention that on May 3rd 2010, near to the first observation point on SR a Baillon's crane was also detected (Stermin, Pripon 2011). Considering this data, we have a total of six rail species located in SR during the spring migration, while on SP four species were recorded (Coot, Moorhen, Water rail and Little Crane).

The observation frequencies defined as number of observations for a single species/ total number of observations reveals a distinct presence of each rallid species considering both the type of habitat and phenological season (Tab. 1). During the breeding season, at SR two species were recorded (Water Rail and Moorhen) and in SP four species (Coot, Moorhen, Water rail and Little Crane), with different observation frequency (Tab. 1).

Taking into account only the species recorded during the breeding season, we calculated the percentage of species present in different types of habitats. In this way, in 80% of cases Coots were recorded in habitats dominated by *T. angustifolia* and 20% in open water; Moorhen was observed 88% inhabiting reed and 12% on open water. On the same criteria Little Crane was observed in 50% of cases in habitat dominated by *T. angustifolia* and 50% in *T. latifolia*, while Water Rail was recorded 91% in reed, 6% in *T. angustifolia* and 3% in *T. latifolia* (Fig. 2).

Discussions

We have chosen to study the presence-absence of rails only during the spring migration and breeding season, because during the post-breeding season many of the species (Water Rail, Little Crane, Spotted Crane and Baillon's crane) are less active vocally, being difficult to detect in dense vegetation (Taylor 1988).

According to our results, during the spring migration, all European rail species, except Coot were located on SR while in SP only four species were located. Species from SP remain there during the breeding season. Considering the ecological necessities of all rallid species recorded in SR during the spring migration only two of them remain there to breed.

Because Corncrake, which is a typical grassland rail and usually avoids marsh areas and reed beds during the breeding season (Taylor 1998) was recorded in SR, we conclude that this species is using the edge of reed beds during spring migration as a safety habitat for resting and optimal for feeding (Roe, Georges, 2007, Loughheed *et al.* 2008). Studies in areas from Southern Transylvania show that 32.32% of singing males were recorded in microhabitats represented by weed, sedge, small bushes as well as reed (Moga *et al.* 2010). A study from western Scotland reveals that in May or early June the birds are selecting the places with tall developed vegetation, like *Urtica dioica*, *Phragmites australis* or *Phalaris arundinacea*, the meadows being used only from the second part of June, when the grass is well developed (Cadbury 1980). These cases show the importance and implication of this homogenous wetland type in the grassland rail species migration and dispersal.

Also for Spotted Crane and Baillon's Crane, the homogenous reed beds are not specific; these birds being more dependent, during the breeding season,

on *Carex*, *Juncus*, *Scirpus* and *Typha* dominated habitats (Taylor 1998). During the spring migration these two species found optimal resting and feeding condition in the large reed beds (Tab. 1).

Even being one of the most specialized rails recorded in our study area during the breeding seasons (using only habitats dominated by *Typha angustifolia* and *T. latifolia*), Little Crane was also found in large reed beds during the spring migration (Tab. 1).

From all the recorded rails, Water Rail and Moorhen are the most generalist ones, nesting in homogeneous and heterogeneous wetland (Tab. 1). Moorhen, observed both in reed and open water, is using all types of wetland habitats monitored here (reed, cattail and open water). Water Rail is also using all wetland areas, except open waters, neither species being dependent on open water during spring migration or breeding season.

Regarding this, Coot is the most aquatic rail species, depending on open water not only in the breeding season but also during spring migration, it has not been detected in homogeneous reed beds.

Even considering that the large and homogeneous reed beds are untypical habitats for many rails (Spotted Crane, Little Crane, Baillon's Crane and Corncrake) (Cramp 1980, Taylor 1998) we assumed that this habitat is a typical one being used by species during a phenological season (spring migration) as important travel corridor and buffer zone. This proves that a homogeneous habitat can have, during migration seasons, higher rail diversity than a heterogeneous one.

Heterogeneous habitats are acknowledged as very important in rail conservation, being considerable areas in rail breeding. However, our results show that homogeneous habitats are also significant and necessary, being typical habitats in rail migration and dispersal. On a large landscape scale, the homogenous habitats, like large reed bed areas are stop-over points for all European rallids, being, during spring, dispersal points to the breeding habitats, having in this way an important role in rallid phenological seasons and wetlands networks.

Acknowledgements

We thank and are very grateful to Ligia I. Baştea for her suggestions in preparing this manuscript and improving the English. We also thank the anonymous reviewers, whose advices helped

improving the paper. This work was supported through the “Investing in people” PhD scholarship, co-financed by the European Social Found, Sectoral Operational Programme “Human

Resources Development” 2007-2013, Babeş-Bolyai University, Cluj-Napoca, Romania.

REFERENCES

- Adamik *et al.* 2003 Adamik Peter, Korňan Martin, Vojtek Juraj, *The effect of habitat structure on guild patterns and the foraging strategies of insectivores birds in forests*. In: *Biologia Bratislava* 58(2) (2003), p. 275-285.
- Brambilla, Jenkins 2009 Brambilla Mattia, Jenkins Richard, *Cost-Effective Estimates of Water Rail *Rallus aquaticus* Breeding Population size*. In: *Ardeola* 56 (2009) p. 95-102.
- Brooks, McLennan 1991 Brooks Daniel, McLennan Deborah, *Phylogeny, ecology and behavior: A research program in comparative biology*. In: *The University of Chicago Press*, Chicago (1991).
- Collar *et al.* 1994 Collar, Nigel, Crosby, Michael, Stattersfield, Allison, *Birds to Watch 2. The world list of throated birds*. In: *Birdlife conservation Series No. 4, Birdlife International*, Cambridge (1994).
- Cadbury 1980 Cadbury Charles, *The status and habitats of the Corncrake in Britain 1978/79*. In: *Bird Study* 27 (1980), p. 203-218.
- Cramp 1980 Cramp, Stanley, (ed.), *The Birds of the Western Palearctic*. In: *Oxford University Press* vol. II., London (1980).
- David 2008 David, Alin, *Ecologia populațiilor de păsări din Câmpia Fizeșului*. In: *Presa Universitară Clujeană*, Cluj-Napoca (2008).
- Haig *et al.* 1997 Haig Susan, Mehlman David, Lewis Oring, *Avian movements and wetlands connectivity in landscape conservation*. In: *Conservation Biology* 12 (1997), p.749-758.
- Jenkins, Ormerod 2002 Jenkins Richard, Ormerod Steve, *Habitat preferences of breeding Water Rail *Rallus aquaticus**. In: *Bird Study* 49 (2002), p. 2-10.
- Joyal *et al.* 2001 Joyal Lisa, McCollough Mark, Hunter Malcom. *Landscape ecology approaches to wetland species conservation: a case study of two turtle species in southern Maine*. In: *Conservation Biology* 15 (2001), p. 1755-1762.
- Lougheed *et al.* 2008 Lougheed Vanessa, McIntosh Mollie, Christian Parker, Stevenson Jan, *Wetland degradation leads to homogenization of the biota at local and landscape scales*. In: *Freshwater Biology* 53 (2008), p. 2402-2431.
- Moga *et al.* 2010 Moga Ioan Cosmin, Hartel Tibor, Öllerer Kinga, *Status, microhabitat use and distribution of the corncrake *Crex crex* in a Southern Transylvanian rural landscape, Romania*. In: *North-Western Journal of Zoology* VI (1) (2010), p. 63-70.
- Price *et al.* 1997 Price Trevor, Helbig Andreas, Richman Adam, *Evolution of breeding distributions in the old world leaf warblers (Genus *Phylloscopus*)*. In: *Evolution* 51 (1997), p. 552-561.
- Robinson, Horsfall 2004 Robinson Robert, Horsfall Joseph, *Rails*. In Perrins, Christopher (ed). *The New Encyclopedia of Birds*, Oxford University Press (2004) Oxford, p. 206-211.
- Roe, Georges 2007 Roe John, Georges Arthur, *Heterogeneous wetland complex, buffer zones and travel corridors: landscape management for freshwater reptiles*. In: *Biological conservation* 135 (2007), p.67-76.
- Stermin *et al.* 2011 Stermin Alexandru, Pripon Liviu, David Alin, Coroiu Ioan. *Wetlands management for Little Crane (*Porzana parva*) conservation in a “Natura 2000” site*. In: *ICESD2011* (2011), p. 91-94.
- Stermin, Pripon. 2011 Stermin Alexandru, Pripon Liviu, *Baillon’s Crane (*Porzana pusilla intermedia*, Herman 1804) Geographical and Historical distribution in Romania*. In: *Brukenthal. Acta Musei* VI (3), Sibiu (2011), p. 493-498.
- Taylor 1998 Taylor, Barry, *Rails. A guide to the rails, crakes, gallinules and coots of the world*. In *Yale University Press*, New Haven and London (1998).

LIST OF ILLUSTRATIONS

- Fig. 1.** The location of observation points and distribution of habitats in Sucutard Pond (SP) and Sic Reed beds (SR).
- Fig. 2.** Percentages of different habitats used by rail species during the breeding season.
- Tab. 1.** The observation frequency of rail species in Sic Reed beds (SR) and Sucutard Pond (SP) during spring migration and breeding season (with bold are values > 0; * Baillon's Crake was located on SR in May 3rd 2010 (Stermin, Pripon 2011))

LISTA ILUSTRĂȚIILOR

- Fig. 1.** Localizarea punctelor de observație și distribuția habitatelor în Heleșteul de la Sucutard (SP) și Stufărișurile de la Sic (SR).
- Fig. 2.** Procentajul diferitelor tipuri de habitate ocupate de speciile de ralde în timpul perioadei de reproducere.
- Tab. 1.** Frecvența observațiilor speciilor de Rallidae la "Stufărișurile de la Sic" (SR) și Heleșteele de la Sucutard (SP) în timpul migrației de primăvară și perioadei de reproducere ("**boldate**" sunt valorile >0; *Crestelul mic a fost identificat la "Stufărișurile de la Sic" în data de 3 mai 2010 (Stermin, Pripon 2011)).

Tab. 1. The observation frequency of rail species in Sic Reed beds (SR) and Sucutard Pond (SP) during spring migration and breeding season.

	SR		SP	
	Observation frequency during spring migration	Observation frequency during breeding period	Observation frequency during spring migration	Observation frequency during breeding period
Coot	0	0	0.25	0.33
Moorhen	0.40	0.23	0.07	0.03
Water rail	1	0.96	0.12	0.16
Spotted crane	0.02	0	0	0
Little crane	0.02	0	0.14	0.06
Baillon's crane	0*	0	0	0
Corncrake	0.01	0	0	0




Fig. 1. The location of observation points and distribution of habitats in Sucutard Pond (SP) and Sic Reed beds (SR).




Fig.2. Percentages of different habitats used by rail species during the breeding season.

SMALL MAMMALS (MAMMALIA: RODENTIA ET INSECTIVORA) FROM THE UPPER TISA RIVER BASIN (ROMANIA)

Ana Maria LAZĂR*
Ana Maria BENEDEK**
Georgiana MĂRGINEAN**

Abstract. During two field campaigns in August 2009 and May-June 2011 a series of investigations using live trapping was carried out in 5 stations along the rivers Vișeu and Iza, in the Tisa River Basin. The researches aimed to draw up a chorological list for the small mammal species and to characterize the community's structure and a similitude analysis of the research stations from this area. Up to the present 28 species are known from Upper Tisa River Basin, among them 8 insectivores and 20 rodents, one rodent (*Rattus rattus*) being probably extinct from the area.

Key words: insectivores, rodents, chorology, community structure, cluster analysis.

Rezumat. Pe parcursul a două campanii de teren, în august 2009 și lunile mai-iunie 2011, am întreprins o serie de investigații pe baza capturării animalelor vii, urmărind obținerea unei liste privind răspândirea acestora și o analiza de similitudine între stațiile investigate în această zonă. Până în prezent, sunt cunoscute 28 de specii din această arie, și anume 8 insectivore și 20 rozătoare. Dintre acestea un rozător (*Rattus rattus*) este probabil dispărut din zonă.

Cuvinte cheie: insectivore, rozătoare, corologie, structura comunității, analiză ierarhică

Introduction

The Upper Tisa River Basin covers in Romania the Maramureș region, including Maramureș Mountains, the northern slopes of Rodna Mountains, and Maramureș Depression. The tributaries of Tisa in Maramureș are: Vișeu, Iza, and Săpânța rivers, the first one having the largest basin.

The first data on the fauna of Maramureș region belong to Hanák (1848 ap. Szilágyi 1876), who describes the observation of *Marmota marmota* in Maramureș Mountains, as well as its whistling in Pietrosul Rodnei. The first faunistic list for Maramureș, including also mammals, was elaborated by Frivaldsky (1875). In 1960 Szabó published a paper on small mammals from the north-western part of Romania, including also data from Valea Vinului (Rodna Mountains). More recently some papers were published including data on the small mammal species from different areas of Maramureș region: Ardelean (1993) - Vaser Valley, Murariu and Răduleț (1998) - Maramureș Depression. The description of the habitat where one specimen of *Sicista betulina*

was captured from Fața Pietrosului Rodnei, along with a synthesis on the current knowledge on this species in Romania are presented by Murariu (1997). In a paper concerning the genus *Arvicola* in Romania (Murariu *et al.* 1997) information from Maramureș depression is also presented. Wagner (1974) published a PhD thesis on small mammals from the Inner Carpathian Basin, including data from Rodna Mountains. However, as there are no details concerning the sampling points, it is possible that at least some of the data come from the southern part of Rodna Mountains, outside the Tisa River Basin. Thus, we did not consider the information from this paper. In 2000 Ardelean and Béres published a synthesis on the vertebrate's fauna from Maramureș. In volume dedicated to the Maramureș Mountains Nature Park of the "Transylvanian Review of Systematically and Ecological Research" (Sibiu, 2008), Gurzău *et al.* published a paper on small mammals. In the same journal, in the volume dedicated to Rodna Mountains National Park (2010) two other papers were published on small mammal's fauna (Nae *et al.* 2010) and on the present distribution of *Marmota marmota* (Szabo 2010). Data concerning the distribution of the population of reintroduced marmots from Rodnei Mountains were previously published (Almășan 1981; Pânzariu 1993; Nădișan 2000).

*"Transylvania" University of Brașov and University of Bucharest, Faculty of Biology, anagurzau@yahoo.co.uk.

** Lucian Blaga University of Sibiu, Faculty of Sciences, benedek_ana@yahoo.com, georgiana.marginean@gmail.com

The present study is based on a series of field investigations carried out by the authors in several sites from the Upper Tisa River Basin, but also on the data collected and published by other authors. The faunistical data available up to the present from this area allows us to draw up a chorological list of the small mammals species from the Upper Tisa River Basin and to do a similarity analysis of the small mammal communities from different sites in the river basins of the three tributaries of Tisa River, namely Vișeu, Iza, and Săpânța.

Research area and methods

Investigations were carried out by live-trapping using hand-made Polish wooden box-traps (Fig. 1) and Fitch traps built according to the model described by Rose (1994). The traps were set either in a rectangular net (in forests) or in transect (along river banks), at 10 m distance one from another. They were baited using sunflower seeds and apple pieces. No pre-baiting was performed. Traps were checked twice a day, in the night and at dawn.

Captured animals were determined according to Pucek (1981), Murariu (2000), and Popescu, Murariu (2001), based on external morphological features. Individuals were marked by cutting their fur in different parts of the body, and then released.

The studies were accomplished during two field campaigns carried out in August 2009 and May-June 2011. Five research stations were established in the Upper Tisa River Basin. The sites were chosen in order to have a good geographical coverage of the area (considering also the data from the literature), both in terms of and altitude. Within a site one or more habitats were chosen to be researched, depending on the heterogeneity of the environment in the station.

1. Repede River (Rodnei Mountains, Vișeu River Basin) – the trapping area was situated at about 900 m altitude, in the spruce forest area. The traps were along the river bank, at 2 kilometres from entrance in Rodnei Mountains Nature Park, in a habitat with the herbaceous layer dominated by *Telekia speciosa*, *Urtica dioica* and *Petasites* sp., grasses and ferns.

2. Pietrosu weather station (Rodnei Mountains, Vișeu River Basin) – situated at 1760 m altitude, in the subalpine area. The traps were placed among the *Juniperus* shrubs in the vicinity of Rodnei Mountains Park Administration' chalet. No small mammal was captured in this station.

3. Săliște de Sus (Iza Valley) - situated at about 670 m altitude. Traps were set along Iza River bank a heterogeneous habitat, a sector with black elder (*Sambucus nigra*), blackberry (*Rubus* sp.) and other shrubs, one along the forest edge and the last with high vegetation dominated by grasses. Another transect was set in a wooded grassland with hawthorn (*Crataegus monogyna*) and wild cherry bushes (*Prunus avium*).

4. Bârsana village (Iza River Basin) – is located at 320 m altitude. Traps were placed in a mown hayfield. No small mammal was captured here.

5. Prislop Pass (Vișeu River Basin) – situated at 1413 m altitude in a coniferous woodland area.

Results and discussions

During the two field campaigns carried out in the Upper Tisa River Basin in August 2009 and May-June 2011, five trapping stations were established. Small mammals were captured only in three of the stations.

A synthesis of the trapping effort and the capture results is presented table 1.

A total of 507 trap nights (283 in 2009 and 224 in 2011) yielded 54 captured specimens belonging to 6 species: two shrews among insectivores - *Sorex araneus* and *Neomys fodiens*, and 4 mice among rodents - *Apodemus flavicollis*, *A. sylvaticus*, *A. agrarius*, and *Mus musculus*. Other two species, *Talpa europaea* and *Arvicola terrestris*, were identified based on visual observations. Among the captured individuals 8 were recaptured – four *A. agrarius* (in Săliște de Sus), three *A. flavicollis* (two in Repede Valley and one in Săliște de Sus), and one *A. sylvaticus* (from Repede Valley). The only species found in all the three sites where small mammals were captured is *A. flavicollis*. This result is consistent with other data from the literature (Hamar 1958, Wagner 1974) and from our previous studies (Benedek *et al.* 2002, Benedek, Drugă 2005, Benedek *et al.* 2005, Benedek 2006, Benedek 2008, Gurzău *et al.* 2008), indicating the wide distribution and high frequency of the yellow-necked mouse in mountain forested areas.

Up to the present 28 species of small mammals are known from the Maramureș Depression, 8 insectivores and 20 rodents. The faunistical and chorological catalogue is presented below. It represents an update of the list presented by (Ardelean, Béres 2000), adding the results of more

recent studies carried out by different authors (Nae *et al.* 2010; Szabo 2010), our present data and previously published (Gurzău *et al.* 2008) as well as the species *Clethrionomys glareolus*, overlooked in the older catalogue. The systematic is given according to Fauna Europaea version 2.5 (Bogdanowicz 2012).

Ord. Insectivora Bowdich 1821

Fam. Erinaceidae Bonaparte 1838

1. *Erinaceus concolor* Martin 1838 - was cited in Maramureş (Kardos ap. Szilágyi 1876, Călinescu 1931), but without mentioning the localities. Ardelean and Béres (2000) mention it as a common species in Maramureş Depression.

Vişeu River Basin: one specimen collected from Crasna, near Hei Valley (Murariu, Răduleţ, 1998), one individual was observed in the Coşnea - Poienile de Sub Munte area (Gurzău *et al.* 2008).

Fam. Soricidae Gray, 1821

2. *Sorex araneus* Linnaeus 1758 - first cited from Maramureş by Kardos (Szilágyi, 1876), is a relatively frequent species in this area (Ardelean, Béres, 2000).

Vişeu River Basin: cited from Repedea, Izvorul lui Dragoş – Faţa Pietrosului Rodnei area (Murariu, Răduleţ 1998), Coşnea - Poienile de Sub Munte (Gurzău *et al.* 2008). During our study one specimen was captured on Repede Valley.

Săpânţa Basin: encountered at Colibi (Murariu, Răduleţ 1998).

3. *Sorex minutus* Linnaeus 1766

Vişeu River Basin: collected from Izvorul lui Dragoş – Faţa Pietrosului Rodnei area (Murariu, Răduleţ, 1998). In the same area the pygmy shrew was captured in 2006 from the vicinity of the laboratory house (Nae *et al.* 2010).

Iza River Basin: cited from Poiana Brazilor, Vidraşca, Stedea, Runcului Valley (Murariu, Răduleţ 1998).

Săpânţa Basin: collected from Brustani (idem).

4. *Sorex alpinus* Schintz 1837

Vişeu River Basin: one specimen collected from Izvorul lui Dragoş – Faţa Pietrosului Rodnei area (Murariu, Răduleţ 1998).

5. *Neomys fodiens* (Pennant 1771) – first cited in Maramureş by Kardos (Szilágyi 1876).

Vişeu River Basin: mentioned from Repedea, Izvorul lui Dragoş (Murariu, Răduleţ 1998),

Iza River Basin: collected from Mara Valley (idem).

6. *Neomys anomalus* Cabrera 1907

Vişeu River Basin: collected from Hei Valley (Murariu, Răduleţ, 1998), during our study an individual was captured in August 2009 in Repede Valley from Rodnei Mountains.

Iza River Basin: found at Poiana Brazilor (idem).

Săpânţa Basin: collected from Colibi and Brustani (idem).

7. *Crocidura suaveolens* (Pallas 1811) - was collected by Béres from Sighetu Marmăţiei, Borşa, Mara, and other localities that are not mentioned (Ardelean, Béres 2000).

Fam. Talpidae Gray, 1825

8. *Talpa europaea* Linnaeus 1758 - was first cited from Maramureş by Kardos (Szilágyi 1876), is a very frequent species in Maramureş region, being spread from the depression area to the subalpine vegetation level (Ardelean, Béres 2000).

Vişeu River Basin: in Maramureş Mountains Nature Park its presence was noted based on the mole hills found in several areas (Gurzău *et al.* 2008). During our study the presence of moles was observed at Sălişte de Sus, also based on the mole hills.

Iza River Basin: During our study the presence of moles was observed at Sălişte de Sus, also based on the mole hills.

Săpânţa Basin: collected from Colibi and Brustani (Murariu, Răduleţ 1998).

Ord. Rodentia Bowdich 1821

Fam. Sciuridae Gray 1821

9. *Sciurus vulgaris* Linnaeus 1758 – was first cited from Maramureş by Szilágyi (1876). In present it is considered common and abundant in Maramureş (Murariu, Răduleţ, 1998), including the localities (e.g. Sighetu Marmăţiei in Grădina Morii park), up to the limit of the spruce forest (Ardelean, Béres, 2000).

Vişeu River Basin: Călinescu (1931) mentions the red squirrel from Făina forest range and Botizu, on Vaser Valley, localities cited later by Ardelean (1993). One specimen was observed on Vaser Valley, upstream Bardău forest range (Gurzău *et al.* 2008).

10. *Marmota marmota* (Linnaeus 1758)

Vişeu River Basin: was reintroduced in Pietrosul Rodnei Reserve in 1973. Two years later the marmots were dispersed in 10 colonies above Iezer Lake and around Buhăescu Lakes, up to the slopes of Buhăescu Peak (Almăşan 1981). In July 1995 the species was observed on the Pietrosul Rodnei slope, above Iezer Lake (Murariu, Răduleţ 1998). In the present the marmots are well established in

the subalpine vegetation level from several areas of Rodnei Mountains (Nădişan 2000). Above Iezer Lake the number of colonies increased from 4 to 7 (Szabo 2010). Four specimens were observed by the authors in August 2009 on the north-eastern slope of Pietrosul Rodnei, in the same area.

Fam. Muridae Gray 1821

11. *Clethrionomys glareolus* (Schreber 1780)

Vişeu River Basin: collected from Hei Valley and Izvorul lui Dragoş – Faţa Pietrosului Rodnei area (Murariu, Răduleţ 1998). Several specimens were captured on Vaser Valley at Bardău and Făina and one specimen was found on Bistra Valley and Valea Neagră (Gurzău *et al.* 2008). From the vicinity of Faţa Pietrosului Rodnei Laboratory House the bank vole was encountered also more recently (Nae *et al.* 2010).

Săpânţa Basin: found at Colibi and Brustani (Murariu, Răduleţ 1998)

12. *Arvicola terrestris* (Linnaeus 1758) - in Maramureş the species was cited for the first time from Sighetu Marmăţiei by Vasiliu and Sova (1968).

Vişeu River Basin: one specimen was observed in Coşnea - Poienile de Sub Munte area (Gurzău *et al.* 2008).

Iza Basin: found at Vidrişca (Murariu, Răduleţ 1998). During our study the presence of *A. terrestris* was observed in a seed bed from Sălişte de Sus, based on the galleries and the information from local people.

Săpânţa Basin: an important population of *A. terrestris scherman* is mentioned in the surroundings of Colibi Forest Range, in the Runcu Mare Valley, in the Colibi stream and Brustani Clearing (Murariu *et al.* 1997, Murariu, Răduleţ 1998).

13. *Microtus subterraneus* (de Selys-Longchamps 1836)

Vişeu River Basin: collected from Izvorul lui Dragoş (Murariu, Răduleţ 1998).

Săpânţa Valley: captured at Colibi (idem).

14. *Microtus agrestis* (Linnaeus 1761)

Vişeu River Basin: two individuals were captured in August 2007 on Vaserului Valley, at Făina and another was found in May 2008 (Gurzău *et al.* 2008).

Iza Basin: captured from Poiana Brazilor and Stedei Valley (Murariu, Răduleţ 1998).

Săpânţa Basin: found at Colibi (idem).

15. *Microtus arvalis* (Pallas 1778) – first cited from Maramureş by Kardos (Szilágyi, 1876), the common vole is considered a frequent species in Upper Tisa River Basin, from the river valleys to Pietrosul Rodnei Peak (Ardelean, Béres 2000).

Vişeu River Basin: collected from Izvorul lui Dragoş (Murariu, Răduleţ 1998).

Săpânţa Basin: found at Colibi and Brustani (idem).

16. *Microtus tatricus* (Kratochvil 1952)

Vişeu River Basin: was first mentioned from Romania by Flousek *et al.* (1985), who collected in 1984 several specimens from the surroundings of Borşa locality.

Iza River Basin: was collected by Murariu and Răduleţ, in 1998 from Runcului Valley, a tributary of Mara River.

17. *Ondatra zibethicus* (Linnaeus 1766) - according to Ardelean and Béres (2000) it was first mentioned from Maramureş in 1963, when a specimen was captured on Tisa River bank, and another from Sighetu Marmăţiei. Between 1960-1970 the species recorded an important expansion of its range, reaching Moisei (Vişeu Valley) and even the brooks from Poienile de sub Munte, followed by a significant retreat caused by drought in the 1980's. In present only a small population still inhabits the area, mainly the wetlands from Tisa River floodplain (idem).

Vişeu River Basin: it was observed at the confluence of the stream Bistra with Vişeu River (Murariu, Răduleţ 1998).

18. *Rattus rattus* (Linnaeus 1758) - was mentioned in Maramureş by Kardos, cited by Szilágyi (1876). In present it is probably extinct from the area, due to the competition with *R. norvegicus*.

19. *Rattus norvegicus* (Berkenhout 1769) - is a widely distributed species in the area, inhabiting especially human settlements. It was observed in all the researched localities, as well as in the railway stations and forest ranges in the vicinity of villages (Murariu, Răduleţ 1998).

Vişeu River Basin: One specimen was found dead in the vicinity of Coşnea - Poienile de Sub Munte (Gurzău *et al.* 2008).

20. *Mus musculus* Linnaeus 1758 - is first mentioned from Maramureş by Kardos (Szilágyi 1876). In present the house mouse is very frequent in Maramureş Depression (Ardelean, Béres 2000).

Iza Basin: captured from the vicinity of Sighetu Marmăţiei (idem), one specimen encountered at Sălişte de Sus, in June 2011.

Săpânța Basin: found at Colibi (Murariu, Răduleț 1998).

21. *Apodemus agrarius* (Pallas 1771) - was captured by Béres in 1967 in several areas from Maramureș, up to 800-1100 m altitude, but the localities are not mentioned (Ardelean, Béres 2000). Vișeu River Basin: captured at Bistra, Valea Neagră, Paltinul (Gurzău *et al.* 2008).

Iza Basin: collected from Vidrișca, Poiana Brazilor, and Mara Valley (Murariu, Răduleț 1998). During our study 25 specimens were captured at Săliște de Sus.

Săpânța Basin: found at Brustani (idem).

22. *Apodemus flavicollis* (Melchior 1834) - is a widespread species, being encountered in most of the researched stations.

Vișeu River Basin: collected from Făina, on Vaser Valley (Ardelean 1993) and by Béres in 1967 from several localities that were not mentioned (Ardelean, Béres 2000). Murariu and Răduleț (1998) collected 50 specimens from Izvorul lui Dragoș – Fața Pietrosul Rodnei area, Repedea, Hei Valley. Several specimens were captured by Gurzău *et al.* (2008) from Coșnea, Bistra Valley, Valea Neagră, Bardău, and Repedea. From the vicinity of Fața Pietrosului Rodnei Laboratory House the yellow-necked mouse was mentioned also more recently by Nae *et al.* (2010). 7 specimens were captured by the authors from Repede Valley in Rodnei Mountains in August 2009, and a specimen was captured in Prislop Pass in May 2011.

Iza Basin: encountered at Poiana Brazilor, Tătaru Chalet, Runcu and Mara valleys (Murariu, Răduleț 1998). 14 specimens were captured by the authors in May 2011 from Săliște de Sus.

Săpânța River Basin: found at Colibi and Brustani (idem).

23. *Apodemus sylvaticus* (Linnaeus 1758) – was first mentioned in Maramureș by Szilágyi in 1876. The wood mouse was collected in 1967 by Béres from different areas (Ardelean, Béres 2000).

Vișeu River Basin: cited from Vaser Valley (Ardelean 1993), several specimens were collected from Izvorul lui Dragoș – Fața Pietrosul Rodnei area, Repedea, Hei Valley (Murariu, Răduleț 1998), one specimen was captured in a beech forest near Vișeu de Sus town in May 2008 (Gurzău *et al.* 2008). From the vicinity of Fața Pietrosului Rodnei Laboratory House the wood mouse was mentioned also by Nae *et al.* (2010). During our study four individuals were captured in August 2009 from Repede Valley in Rodnei Mountains.

Iza Basin: collected from Rona de Sus, Vidrișca, Tătaru Chalet, Stedei, Runcu and Mara valleys (Murariu, Răduleț 1998).

Săpânța River Basin: found in Colibi and Brustani (idem).

24. *Apodemus uralensis* (Pallas 1811)

Iza River Basin: four specimens were collected in 1967 by Béres from the Iza River terrace, at 350 m altitude (Ardelean, Béres 2000).

25. *Micromys minutus* (Pallas 1771)

Iza River Basin: one specimen was collected from Mara Valley (Murariu, Răduleț, 1998), found in a cultivated field at Dobăieș (Sighetu Marmăției) (Ardelean, Béres 2000).

Fam. Myoxidae (Thomas 1897)

26. *Glis glis* Linnaeus 1766 – was first mentioned from Maramureș by Kardos (Szilágyi 1876)

Vișeu River Basin: the species was mentioned from the Vaser Valley (Ardelean 1993) and Repedea (Murariu, Răduleț 1998). Direct evidences of their presence in the area (broken tails) were found at Bistra, where it is trapped as a pest (Gurzău *et al.* 2008).

27. *Muscardinus avellanarius* (Linnaeus 1758) – according to Ardelean and Béres (2000) it is a common species in Maramureș, but rarely observed. Vișeu River Basin: recorded in the literature from Vaser Valley (Ardelean, 1993), Izvorul lui Dragoș – Fața Pietrosului Rodnei area and Repedea (Murariu, Răduleț 1998).

Iza Basin: cited from Coștiui and Agriș (Ardelean, Béres 2000).

Săpânța Basin: Câmpulung la Tisa (idem).

Fam. Dipodidae Gray 1821

28. *Sicista betulina* (Pallas 1779)

Vișeu River Basin: the species is known from the area based on a female specimen captured at Fața Pietrosului Rodnei, at 1375 m altitude (Murariu, 1997, Murariu, Răduleț 1998). The species is considered a glacial relict, in Romania having a reduced distribution in high montane and subalpine areas.

Due to the relatively low trapping effort, short capture sessions, and the random choice of sites we can not build a sound image on the small mammals community structure, and assess its dependence on different time (year) and space (river basin, site, altitude) variables. Besides, using only one capture method reduces the span of captured species. However, some characteristics of small mammal

communities may be revealed by the results of the present study.

The highest number of small mammals was captured in the lowest station from the three stations where they were captured (Săliște de Sus), situated at 670 m, while at Prislop Pass, located at 1400 m, only one specimen was found. At higher altitudes, in the subalpine vegetation level (Pietrosu weather station - 1760 m) no small mammal was captured. As the trapability of small mammals is unlikely to be significantly influenced by altitude, we consider that there is a decrease in the abundance of small mammal community along with the increase in altitude. This tendency was observed also in other mountain areas from the Romanian Carpathians (Ceahlău – Simionescu 1968; Retezat and Lotru mountains – Benedek 2008).

The increase in altitude influences not only the total abundance of the small mammal community, but also its specific structure and the density of particular populations. *Apodemus flavicollis* is the only species captured in all the three stations situated in the wooded area. It is a typical sylvic rodent, inhabiting all types of forests, both from lowlands and mountains. Its presence above the tree line is limited to the subalpine shrubs, and especially to the mugo pine (*Pinus mugo*). However, according to our previous results from Retezat Mountains and Râu Șes River Basin (Benedek 2008), its presence in the subalpine area is temporary, confined to the high density years.

The increase in altitude causes a shift in the specific structure of small mammal communities.

In Săliște de Sus, from Iza River Basin, situated in the beech forest area, the community is dominated by *Apodemus agrarius* (62.5%) (Fig. 2). The low altitude and the vicinity of the water course appear to favour the striped field mouse. In several mountain areas from Romanian Carpathians where studies on small mammals were carried out - Ceahlău Mountains (Simionescu, 1968); Retezat, Gurghiu, Harghita, and Gilău mountains (Wagner 1974); Retezat and Apuseni mountains, Râu Șes River Basin (Benedek 2008) - *A. agrarius* was not encountered. Wagner (1974) mentions it only from Rodna Mountains, at altitudes between 600 and 800 m a.s.l. Hamar (1958) notes the striped field mouse from Bâlea Valley in Făgăraș Mountains only in moist places in the vicinity of water courses, up to 600-650 m.

Mus musculus is another element characteristic for lower altitudes. Related to human settlements, although it can be present in inhabited buildings even above the tree limit, it is more frequent in lowlands, where it can survive also in natural habitats. Wagner (1974) found the house mouse in a building at 650 m, but it was absent from another at 1350 m. As we researched a natural habitat, the ratio of *M. musculus* is low, representing only 2.5% (one specimen) of the captured specimens. Due the low altitude of this station and the vicinity of the beech forest *A. flavicollis* records the highest abundance among the researched stations, but it is outnumbered by *A. agrarius*, which is favoured on the river bank by the high humidity. Thus, *A. flavicollis* represents in this habitat only 35% of the captured specimens.

In Repede Valley from Rodnei Mountains (Fig. 3), four species were captured, two rodents (*A. flavicollis* and *A. sylvaticus*) and two insectivores (*S. araneus* and *N. anomalus*), each of them with one specimen (7.7%). The community is dominated by rodents, and among them *A. flavicollis* represents more than half of the captured specimens (53.8%). In Făgăraș and Retezat Mountains (Hamar, 1968) the yellow-necked mouse populates the whole forested area. According to our previous results from other mountain areas (Benedek *et al.* 2002; Benedek, Drugă 2005; Benedek *et al.* 2005; Benedek 2006, Benedek 2008) *A. flavicollis* is the dominant rodent from mountain forest in the Romanian Carpathians. In Repede Valley the forest habitats which cover most part of the slopes provide favourable conditions to the yellow-necked mouse.

A. sylvaticus represents 30.8% of the community. The relatively high ratio of the wood mouse is in concordance with the results of Wagner (1974), who captured from Rodna Mountains 28 specimens of *A. flavicollis* and 19 specimens of *A. sylvaticus*. Murariu and Răduleț (1998) found the wood mouse (116 captured individuals) being more abundant in Maramureș Depression than the yellow-necked mouse (50 individuals). These results are probably due to the difference in the mean altitude of the research area in the two studies, Murariu and Răduleț focusing on the mountain valleys. During our previous studies in Maramureș, we encountered the wood mouse only in one station, near Vișeu de Sus (Gurzău *et al.* 2008). However, in other mountain areas from the Carpathians *A. sylvaticus* was not encountered at all by the authors, neither along the mountain valleys, nor on the slopes (Benedek *et al.* 2002;

Benedek, Drugă, 2005; Benedek *et al.*, 2005; Benedek 2006; Benedek 2008). The presence and relatively high abundance of *A. sylvaticus* along the valleys from the Upper Tisa River Basin, in the Maramureş and Rodna Mountains, is most likely due to the great changes occurred in the mountain landscape from this area, the expansion of human settlements along the valleys, and especially the massive forest cuttings which lead to large corridors of meadows and shrubs, *A. sylvaticus* usually avoiding compact forests (Hamar 1968).

Using our field data as well as the data from previously published faunistical studies (Murariu, Răduleţ 1998, Gurzău *et al.* 2008, Nae *et al.* 2010), a cluster analysis was done between the 23 stations along the main rivers from Upper Tisa Basin researched by different authors, based on the small mammal community structure. A synthesis of the investigated sites and habitats during previous studies, used for the cluster analysis, is presented in Tab. 2.

Due to the differences in the trapping technique, as well as in the trapping effort used in the various studies, we considered only presence-absence data. Besides, we considered only the small mammals species that enter traps set on the ground. Thus, we did not include in the analysis *E. concolor*, *T. europaea*, *S. vulgaris*, *M. marmota*, and *R. norvegicus*, which were identified based on direct or indirect visual observations. Among the dormice only the smallest species (i.e. *M. avellanarius*), which enters more frequently the traps for terrestrial small mammals, was included in the analysis. Among the research stations the three sites with no captured species were not included in the analysis. Table 3 contains the presence-absence data used for the cluster analysis.

The location of the 26 research stations (23 with captures and 3 blank), as well as their cluster are indicated in Fig. 4.

The terrestrial small mammals fauna from the Upper Tisa River Basin, as revealed by the trapping results is very heterogeneous, the similarity between the researched sites being low. Most of the stations join at a distance of 0.5 or greater (Fig. 5). Considering 0.7 as the cutting similarity level, the dendrogram illustrates six major groups. The most distinct research station (I), grouped at a distance of 0.91 in terms of Jaccard index, is Paltinul Chalet, on a river bank in beech forest, where only one species was captured, in this case *A. agrarius* (Gurzău *et al.* 2008). The second

cluster (II) is also distinct. It is formed of three sites. Among them, Vişeu de Sus and Rona de Sus, situated in mixed forest, in the vicinity of localities, have a 100% similarity, as only the wood mouse was captured (Murariu, Răduleţ, 1998; Gurzău *et al.* 2008). They are joined at a relatively great distance (0.66) by Vidrişca. Group III includes two stations from Iza River Basin, namely Valea Stedei and Poiana Brazilor, separated by a small distance, sharing half of their species (*S. minutus*, *M. agrestis*, and *A. agrarius*). The most diverse faunas are sheltered by the sites joined in group IV. It comprises the two localities from Săpânţa River Basin (Colibi and Brustani) and Izvorul lui Dragoş. The high diversity from the first two sites (10 and respectively 8 species) is probably due to the lower human impact in this area (Murariu, Răduleţ, 1998), while the great number of species (11 – the highest among all the research stations) from Izvorul lui Dragoş is also an effect of the higher capture effort. This is the only station where three field campaigns were carried out (*idem*). Another factor contributing to the increase in the number of captured species is the use of snap traps along with life traps. Cluster V is formed of 7 sites from Iza and Vişeu river basins characterised by the co-occurrence of *A. sylvaticus* and *A. flavicollis*. The last group (VI) joins 7 stations researched by the authors during the present study or previously. They are typical forest sites, with low specific diversity and *A. flavicollis* as the common element. Another characteristic is the absence of *A. sylvaticus*. Among these localities Bistra and Valea Neagră, from Vişeu River Basin, have a 100% similarity, all the three captured species being the same (*C. glareolus*, *A. agrarius*, and *A. flavicollis*). They are located at a small distance one from another, in similar habitats, consisting of river banks in beech forest (Gurzău *et al.* 2008). Bardău, in the mixed forest, and later Făina, in the spruce forest, are joined based on the presence of *C. glareolus*. Other two stations from this group, Prislop and Sălişte, have only the yellow necked mouse in common.

Conclusions

Up to the present 28 species of small mammals (8 insectivores and 20 rodents) are known from the Upper Tisa River Basin. One of them, *Rattus rattus*, is probably extinct. 6 species were captured during our research and other two were noted based on indirect observations.

The small mammals communities from the investigated area are dominated by the rodents belonging to *Apodemus* genus. *A. agrarius*, a

hygrophilous species inhabiting mainly open habitats along rivers in lower areas, and *A. flavicollis*, a typical forest species. *A. sylvaticus* has also an important share in the communities, indicating a change in the natural forest landscape from the Upper Tisa River Basin, especially through deforestation.

The grouping of research stations based on the small mammals fauna is influenced mainly by altitude and habitat type, but also by the trapping effort and research method. Stations from the same

river basin are sometimes joined together (Colibi and Brustani from Săpânța River Basin, Poiana Brazilor and Valea Stedei from Iza Basin), usually as an effect of small distance and habitat homogeneity.

Acknowledgements. The field investigations were accomplished with the indispensable help of Lazăr Cătălin Nicolae.

REFERENCES

- Almășan 1981 Almășan Ioan, *Situația actuală a răspândirii marmotei din România*. In: *Vânătorul și Pescarul Sportiv* 6, București (1981), p. 8-9.
- Ardelean 1993 Ardelean Gavrilă, *Fauna de vertebrate din stațiunea Făina (Vaser) și împrejurimi*. In: *Bul. Științ., seria B, fasc. Biol. - Chim.* 10, Baia Mare (1993), p. 84-90.
- Ardelean, Béres 2000 Ardelean Gavrilă, Béres Iosif, *Fauna de vertebrate a Maramureșului*. In: *Edit. Dacia*, Cluj-Napoca (2000), p. 283-314.
- Benedek *et al.* 2002 Benedek Ana Maria, Sîrbu Ioan, Coțofană Mihai, *Study on the small mammal communities from the Lotrioara River Basin (Lotru Mountains)*. In: *Trav. Mus. Nat. Hist. natl. "Grigore Antipa"* XLIV, București (2002), p. 455 - 464
- Benedek, Drugă 2005 Benedek Ana Maria, Drugă Marius, *Data regarding the small mammal communities (Mammalia: Insectivora et Rodentia) from Râu Șes River Basin (Țarcu and Godeanu Mountains, Romania)*. In: *Trav. Mus. Nat. Hist. natl. "Grigore Antipa"* XLVIII, București (2005), p. 321 - 329.
- Benedek *et al.* 2005 Benedek Ana Maria, Sîrbu Ioan, Vasile Mihai, *Small mammals (Ordo Insectivora and Ordo Rodentia) from the Apuseni Mountains Natural Park area (Western Carpathians, Romania)*. In: *Studii și Comunicări, serie Biologie, Muzeul Județean Satu Mare V-VI*, Satu Mare (2005), p. 116-121.
- Benedek 2006 Benedek Ana Maria, *Small mammals (Ordo Insectivora and Ordo Rodentia) from Retezat National Park (Romania)*. In: *Transylvanian Review of Systematical and Ecological Research* 3, Sibiu (2006), p. 139-146.
- Benedek 2008 Benedek Ana Maria, *Studii asupra mamiferelor mici (Ordinele Insectivora și Rodentia) în Transilvania - România*. In: *PhD thesis, University of Bucharest*, București (2008), p. 1-318.
- Bogdanowicz 2012 Bogdanowicz Wiesław, *Fauna Europaea: Mammalia. Fauna Europaea version 2.4* (2012), <http://www.faunaeur.org> (accessed 25 July 2012).
- Călinescu 1931 Călinescu Raul, *Mamiferele României. Repartiția și problemele lor biogeografice-economice*. In: *Bul. Min. Agric. Domenii* 251, București (1931), 1, p. 1-103.
- Flousek *et al.* 1985 Flousek Jirí, Flouskova Z., Tomasova K., *To the knowledge of small mammals in the Rodnei Mts. (Rumania)*. In: *Veřstník Cěskoslovenské společnosti zoologické (Acta Societatis Zoologicae Bohemoslovenicae)* 49, Praze (Prague) (1985), p. 6-17.
- Frivaldsky 1875 Frivaldsky János, *Adatok Maramoros vármegye faunájához*. In: *Mathematika es természettudományi Közlemenyek IX*, Budapest (1875), p. 183-232.
- Gurzău *et al.* 2008 Gurzău Anamaria, Benedek Ana Maria, Sîrbu Monica, Sîrbu Ioan, *Small mammals from the Maramureș Mountains Nature Park*. In: *Transylv. Rev. Syst. Ecol. Res.* 5, Sibiu (2008), p. 215 – 222.
- Murariu 1997 Murariu Dumitru, *La siciste de bouleux (Sicista betulina Pallas, 1779) (Rodentia, Zapodidae) dans les monts Rodna, Maramureș, Romania*. In: *Trav. Mus. Hist. Nat. "Gr. Antipa"* 37, București (1997), p. 147-157.
- Murariu *et al.* 1997 Murariu Dumitru, Răduț Năstase, Stănescu Mihai, *Contributions to the knowledge of the genus Arvicola Lapeyère, 1799 (Mammalia, Rodentia) from Romania*. In: *Trav.*

- Murariu, Răduleț 1998 *Mus. Nat. Hist. Natl. "Gr. Antipa"* 39, București (1997), p. 401-413.
Murariu Dumitru, Răduleț Năstase, *Mammalian fauna (Mammalia) from Maramureș Depression, Romania*. In: *Trav. Mus.Hist. Nat. "Gr. Antipa"* 40, București (1998), p. 609-621.
- Murariu 2000 Murariu Dumitru, *Insectivora*. In: *Fauna României, Mammalia XVI* (1), Editura Academiei Romane, București (2000), p. 1-142.
- Nae et al. 2010 Nae Ioana, Cobzaru Ioana, Cozma, Andreea Marina, *New data concerning the small mammal fauna (Insectivora, Rodentia) on the Rodna Mountains National Park (Eastern Carpathians, Romania)*. In: *Transylv. Rev. Syst. Ecol. Res.* 9, Sibiu (2010), p. 193-204.
- Nădișan 2000 Nădișan Ioan, *Pietrosul Rodnei – Rezervație a Biosferei*. In: *Editura Muzeului Județean Maramureș, Baia Mare* (2000), p. 1-187.
- Pânzariu 1993 Pânzariu Gheorghe, *Marmota (Marmota marmota L.) în Munții Rodnei, după 20 de ani de la colonizare*. In: *Ocot. Nat. Med. Înconj.* 37(1), București (1993), p.11-18.
- Popescu, Murariu 2001 Popescu Alexandrina, Murariu Dumitru, *Rodentia*. In: *Fauna României. Mammalia* Editura Academiei Romane XVI (2), București (2001), p. 1-214.
- Pucek 1981 Pucek Zdzislaw, *Keys to Vertebrates of Poland – Mammals*. In: *Polish Scientific Publishers, Warszawa (Warsaw)* (1981), p. 1-339.
- Rose 1994 Rose Robert, *Instructions for Building Two Live Traps for Small Mammals*. In: *Virginia Journal of Science* 43(3), Richmond (1994), p. 151-157.
- Simionescu 1968 Simionescu Viorica, *Contribuții la cunoașterea componenței specifice și repartiției pe verticală a mamiferelor mici de pe masivul Ceahlău*. In: *An. St. Univ. "Al. I. Cuza", Sect. II*, 14 (2), Iași (1968), p. 365-372.
- Szabo 2010 Szabo Beatrice Medea, *Data on the distribution of Marmota marmota (Rodentia, Sciuridae) from the Rodna Mountains National Park (Eastern Carpathians, Romania)*. In: *Transylv. Rev. Syst. Ecol. Res.* 9, Sibiu (2010), p. 205-210.
- Szabó 1960 Szabó I., *Contribuții la cunoașterea faunei de mamifere mici din partea nord-vestică a Republicii Populare Române*. In: *Stud. Univ. "Babeș-Bolyai", Ser. II, Biol. Cluj-Napoca* (1960), p. 119-125.
- Szilágyi 1876 Szilágyi János, *Maramoros egyetemes leírása*, Budapest (1876).
- Vasiliu, Șova 1968 Vasiliu George, Șova Constantin, *Fauna Vertebratica Romaniae (index)*. In: *Stud. Com. Muz. Șt. Nat. Bacău*, 1 partea II, Bacău (1968), p. 215-254.
- Wagner 1974 Wagner Otto, *Biogeografische Untersuchungen an Kleinsäugerpopulationen des Karpatenbeckens*. In: *PhD thesis, Saarbrücken* (1974) p. 1-179.

LIST OF ILLUSTRATIONS

- Fig. 1.** Polish wooden box-trap used in the research
Fig. 2. Specific structure of small mammal communities from Săliștea de Sus, Iza River Basin, expressed in terms of relative abundance (%)
Fig. 3. Specific structure of small mammal communities from Repede Valley, Rodnei Mountains, Vișeu River Basin, expressed in terms of relative abundance (%)
Fig. 4. Position of the 26 research sites from Upper Tisa River Basin and the results of the cluster analysis
Fig. 5. Cluster analysis of the research stations, based on Jaccard index, average linkage clustering
Tab. 1. The trapping effort and number of specimens of the small mammals species encountered in the investigated habitats, during the field researches in 2009 and 2011
Tab. 2. Investigated sites and habitats and capture method used during previous studies
Tab. 3. Presence-absence data for the captured species in the research sites

LISTA ILUSTRAȚIILOR

- Fig. 1.** Capcană-cutie de tip polonez utilizată în studiu
Fig. 2. Structura specifică a comunităților de mamifere mici din Săliștea de Sus, bazinul Izei, exprimată în termeni de abundență relativă (%).
Fig. 3. Structura specifică a comunităților de mamifere mici de pe Valea Repede, Munții Rodnei, exprimată în termeni de abundență relativă (%).
Fig. 4. Poziția celor 26 de stații de cercetare din bazinul superior al Tisei și rezultatele analizei ierarhice
Fig. 5. Dendrograma stațiilor de cercetare, întocmită pe baza indicelui Jaccard, prin metoda grupării la distanță medie
Tab. 1. Efortul de captură și numărul de indivizi din speciile de mamifere mici întâlnite în habitatele investigate în timpul campaniilor de teren din 2009 și 2011
Tab. 2. Stațiile investigate în studiile anterioare și metodele utilizate
Tab. 3. Datele de prezență-absență pentru speciile capturate în stațiile de cercetare



Fig. 1. Polish wooden box-trap used in the research




Fig. 2. Specific structure of small mammal communities from Săliște de Sus, Iza River Basin, expressed in terms of relative abundance (%).




Fig. 3. Specific structure of small mammal community from Repede Valley, Rodnei Mountains, Vișeu River Basin, expressed in terms of relative abundance (%).




Fig. 4. Position of the 26 research sites from Upper Tisa River Basin and the results of the cluster analysis (the roman figure indicates the cluster for each site, the empty circle is for the sites with no captures; the squares with arabic figures indicate the sites trapped during this study)




Fig. 5. Cluster analysis of the research stations, based on Jaccard index, average linkage clustering (The stations from Vișeu River Basin have the V letter preceding their name, the stations from Iza Basin, the letter I, and those from Săpânța Basin, the letter S)

Tab. 1. The trapping effort and number of specimens of the small mammals species encountered in the investigated habitats, during the field researches in 2009 and 2011.

Station	Habitat	Date	Trapping effort (traps*nights)	Species	No. ind.
Repede Valley	river bank	August 2009	52*5 = 260	<i>A. flavicollis</i>	7
				<i>A. sylvaticus</i>	4
				<i>S. araneus</i>	1
				<i>N. anomalus</i>	1
Pietrosu Rodnei	subalpine shrubs	August 2009	23*1 = 23	-	-
Săliște de Sus	river bank	May 2011	60*2 = 120	<i>A. flavicollis</i>	14
				<i>A. agrarius</i>	25
				<i>M. musculus</i>	1
	wooded grassland	May 2011	34*1 = 34	-	-
Bârsana	hayfield	May 2011	40*1 = 40	-	-
Prislop Pass	coniferous forest	June 2011	30*1 = 30	<i>A. flavicollis</i>	1
TOTAL			507 trap-nights		54

Tab. 2. Investigated sites and habitats and capture method used during previous studies

Author	Capturing method	Site	Investigated habitat types
Murariu, Răduleț (1998)	snap traps and life traps, some specimens being caught by hand (<i>Erinaceus concolor</i> , <i>Muscardinus avellanarius</i> , <i>Sicista betulina</i>),	Izvorul lui Dragoș	not specified
		Colibi	
		Brustani	
		Repedea	
		Hei Valley	
		Tătaru Chalet	
		Rona de Sus	
		Vidrișca	
		Poiana Brazilor	
		Valea Stedei	
		Runcu Valley	
		Mara Valley	
Gurzău et al. 2008	live trapping using Polish box traps and Fitch traps set 10 m apart, in line (along rivulets) or rectangular net (in forests)	Paltinul Chalet	beech forest, bank
		Repedea	beech forest edge, swamp, meadow
		Vinderel Lake	meadow, shrubs
		Coșnea	beech forest, bank
		Bărdău	mixed forest, bank, meadow, spruce forest
		Făina	spruce forest, bank, meadow
		Bistra	beech forest
		Valea Neagră	beech forest, bank
		Vișeu de Sus	mixed forest, bank
Nae et. al. 2010	box traps set in line	Fața Pietrosului Laboratory House	spruce forest, spruce forest edge, grazed pasture,

Tab. 3. Presence-absence data for the captured species in the research sites

	Izvorul lui Dragoș	Fața Pietrosului	Repedea	Repede Valley	Hei Valley	Bistra	Valea Neagră	Paltinul	Coșnea	Vișeu de Sus	Bardău	Făina	Prislop	Tătaru Chalet	Rona de Sus	Vidrișca	Poiana Brazilor	Stedei Valley	Runcului Valley	Mara Valley	Săliștea de Sus	Colibi	Brustani
<i>S. araneus</i>	+	-	+	+	-	-	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>S. minutus</i>	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	+	+	+	+	-	-	-	+
<i>S. alpinus</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>N. fodiens</i>	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>N. anomalus</i>	-	-	-	+	+	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	+	+
<i>C. glareolus</i>	+	+	-	-	+	+	+	-	-	-	+	+	-	-	-	-	-	-	-	-	-	+	+
<i>A. terrestris</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	-	+	+
<i>M. subterraneus</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-
<i>M. agrestis</i>	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-	+	+	-	-	-	+	-
<i>M. arvalis</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	+
<i>M. tatricus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-	-
<i>M. musculus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	+	-
<i>A. agrarius</i>	-	-	-	-	-	+	+	+	-	-	-	-	-	-	-	-	+	+	-	+	+	-	+
<i>A. flavicollis</i>	+	+	+	+	+	+	+	-	+	-	+	+	+	+	-	-	+	-	+	+	+	+	+
<i>A. sylvaticus</i>	+	+	+	+	+	-	-	-	-	+	-	-	-	+	+	+	-	+	+	+	-	+	+
<i>M. minutus</i>	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	+	-	-	-
<i>M. avellanarius</i>	+	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>S. betulina</i>	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

HERBARIUM VIVUM: A HISTORICAL COLLECTION OF MEDICINAL PLANTS FROM THE NATURAL HISTORY MUSEUM (SIBIU, ROMANIA)

Ghizela VONICA *
Aurelia HOROTAN **

Abstract. The herbarium book with medicinal plants - *Herbarium Vivum*, is one of the oldest herbarium from the Natural History Museum in Sibiu, dated at the beginning of the 19th century. The herbarium was organized by the chemist Joseph Oberth from Aiud (Alba Iulia County, Romania). The dried plants collection was a prerequisite to finish the pharmaceutical study and was intended to help the chemist in identifying plants from the wild flora. Oberth J., was the owner of the "Golden Star" Pharmacy (Aiud), and the tutor of the young chemist Ludwig Binder, to whom he gave his own herbarium with medicinal plants for further support in his practice. The dried plant collection was kept by Binder until 1910, when he passed it on to Karl Pissel. Following the Ministerial Decree (1949), the historical heritage from all the pharmacies around Romania, including the herbarium, was moved to Sibiu under inventory and stored at Pharmacy no. 20. After a short time, following another Ministerial Decree, the *Herbarium Vivum*, together with other pharmaceutical historical pieces, was moved to the Brukenthal National Museum from Sibiu, in order to keep and maintain a collection dedicated to the history of pharmacy. Today this pharmaceutical herbarium book is included in the botanical collection from the Natural History Museum of Sibiu. The *Herbarium* contains 130 pressed plants, and it has a scientific importance regarding the ethopharmacological flora from Romania.

Keywords: plant collection, dried specimens, medicinal properties, history of pharmacy.

Rezumat. Colecția cu plante medicinale de tip carte - *Herbarium Vivum*, este un ierbar de la începutul secolului al XIX-lea, realizat de farmacistul Joseph Oberth, proprietarul farmaciei "Steaua de Aur" (1857-1891) din Aiud. Ulterior acesta îi dăruiește ierbarul lui Ludwig Binder pentru a-l sprijini în stagiul său ca farmacist. Ierbarul este păstrat de Binder până în anul 1910, după care este înmănat lui K. Pissel, proprietar al farmaciei „La Coroana” din Sibiu și președinte al asociației „Gremiul farmaciștilor din Transilvania”, districtul Sibiu. În urma unui Decret Ministerial (1949), tot patrimoniul unităților sanitare și farmaciilor trec în subordinea statului, inclusiv *Herbarium Vivum*. După o scurtă perioadă de timp, printr-un alt decret ministerial, *Herbarium Vivum* împreună cu toate vestigile farmaceutice au fost cedate Muzeului Național Brukenthal în vederea realizării unei colecții de acest gen. În prezent acest ierbar de tip carte, cu 60 de coli se află în inventarul colecției de botanică din cadrul Muzeului de Istorie Naturală din Sibiu. Ierbarul conține 130 de specii, dintre care, mai mult de jumătate, sunt specii care nu se mai regăsesc în practicile farmaceutice și tradiționale de astăzi. Materialul ierborizat constituie o adevărată valoare pentru farmaciști și botaniști deoarece redă în parte cunoștințele și practicile farmaceutice ale acelor timpuri.

Cuvinte cheie: ierbar, plante presate, proprietăți farmaceutice, istoria farmaciei.

Introduction

In the 17th century, plants were studied by chemists and physicians who have contributed to the development of Botany as a science. In fact, the early botanical phytotherapy records are from the 17th century, when the first pharmacies were opened. Also, during that period, the first herbaria in the history of Botany were recorded. Transylvania's flora was also studied over time by famous chemists or physicians as: Fr. Kladny, G.

* Brukenthal National Museum, Natural History Museum, aghizela@yahoo.com; ghizela.vonica@brukenthalmuseum.ro

** Terezianum High School, Sibiu,

Kaysser, J. Bielz, J Römer, etc.

The chemist's main priority in achieving these herbaria was the preservation of as many plant parts as possible in order to facilitate the recognition of the species and to harvest more accurate the plants from the wild flora. Many of them, passionate botanists, have bequeathed to the Natural History Museum in Sibiu their own herbaria or collections of drugs. These herbaria are scientific proofs regarding Romanian ethnopharmacology and phytopharmacology because today's industrialized pharmacies do not use anymore many of this curative herbs. The

existence of these medicinal herbaria is also essential for plant taxonomy, species distribution, threatened species study and for the history of pharmacy and botany.

The aim of this paper is to assess the contribution of Oberth's *Herbarium Vivum* in the study of Romania's present-day flora from a botanical point of view and its importance to ethnopharmacology and pharmaceutical history. This paper clarifies also why there are two names attached to the *Herbarium Vivum* and who is the real author.

Materials and methods

All specimens were identified using the monographic work "Flora României" (Ciocârlan, 2009), updated with the Plant List (www.theplantlist.org) and NCBY- Taxonomy Browser (<http://www.ncbi.nlm.nih.gov>). To find out more about the curative properties of the plants, the authors revised recent literature on medicinal plants (Alexan *et al.* 1988, Oroian 2011, Panțu, 1929), ethnoiatry papers (Drăgulescu 1976, Oroian 2009, Pop, Olos 2004) and the Romanian Pharmacopoeia from 1993. The medicinal plants included in the herbarium were compared with the Pharmacy History Museum Collection inventory book and phytopharmacological literature (Ban 2007, Crișan 1996). The Romanian National Archives from Sibiu County (Archiv- Sibiu Direction) and pharmaceutical history literature were consulted in order to extract bibliographical data on the chemists, their scientific and professional activity (Acker 1970; Fabritius 1986; Gheorghiu 1981; Orient 1926, Paveleanu 2004; Roth 1970, 1973, 1976, manuscript).

Results and discussions

The Natural History Museum in Sibiu Botanical heritage includes numerous medicinal plants herbaria, one of these being Oberth's *Herbarium Vivum* (Fig.1.). At first this herbarium was a puzzle because its source was not known. The only evidence was the size of the book covers and the handwriting on it. The herbarium book covers are typical for the 18-19th century. Inside there are 130 species attached to 60 sheets of paper, all in a good conservation status (Fig.1.). On the front cover of the *Herbarium Vivum*, there is a small, black, handwritten note: "*Herbarium Vivum*", "Joseph Oberth" and the year "1839", meaning that the specimens were probably collected by Joseph Oberth (Fig.2.). On the inside of the cover and on the first sheet handwritten with red "*Apotheke zum Adler des Ludwig Binder in Mühlbach*" (Ludwig Binder' "Eagle Pharmacy" from Sebes) and on top of sheet is noted with black ink (other than the

cover) "*Ludwig Binder, Apotheker in Mühlbach, 28.02.1910*" (Fig.3.). The scientific name of each species is written with the same ink as on the front cover and this confirmed that the herbarium was prepared by Joseph Oberth (Fig. 4.). Starting from Joseph Oberth and Ludwig Binder's "Eagle Pharmacy" from Sebes the authors could find out the history of this herbarium and, in chronological order, the events that led to its presence in the Natural History Museum heritage.

Paveleanu (2004) describes with details the history of Pharmacy in Transylvania, the development of the pharmacies in the area, the chemists' activities starting with the 1700. According to Paveleanu (2004), Joseph Oberth (Iosif Oberth) was also a chemist as Ludwig Binder. He was the owner of the "Golden Star" (Aranycsillag) pharmacy from Aiud (Alba County). The pharmacy was founded in 1750 and was led by Oberth from 1857 until 1891 (Fabritius 1986). Comparing the year written on the front cover of the herbarium to the period in which he was practicing as a chemist in Aiud the authors concluded that the *Herbarium Vivum* was attained during his apprentice pharmacist period, as the herbarium was a necessary and required condition to finish the studies (Fig.2.). In that period, the private pharmacies were seen as a business and could be passed on from one generation to another and so Oberth J. left the pharmacy to his son Oberth J. Erben. The latter became the owner of the pharmacy between 1891-1912, but Oberth Erben could not practice the profession, probably due to the lack of a bachelor's degree (Fabritius 1986). Each owner could hire an apprentice or chemist assistant to help him in his work. Probably this is the reason why Joseph Oberth hired the young Ludwig Binder as an apprentice and later as an assistant. Unfortunately, no document that substantiates this assumption of the association, but the fact that the two chemists worked together or came in contact with each other at one point is proven by the herbarium inscriptions.

It was known that students were admitted to pharmaceutical school after obtaining the gymnasium certificate, at the age of 14 or 15. After concluding the studies they would become debutants and work under the supervision of a certified chemist in his pharmacy until they got acquainted with the practical aspects of the profession. At the end of the debutant period they were examined by a pharmaceutical board and received their final certificate (Gheorghiu 1981). It seems that Ludwig Binder completed his debutant period at the "Golden Star" pharmacy. Probably

Binder J. Ludwig received the medicinal plant collection from his supervisor to facilitate his work in the future.

A newly licensed chemist in order to exercise his profession, in his one pharmacy, needed to send a petition to the Hungarian Ministry and the petition had to be approved by three doctors (concession no. 1296/1878). For example, the Sebeş town's chemist Edward Müller submitted in 1874 the documentation for a second pharmacy in town. The three doctors did not concur and the opening was delayed. Also the negotiation with the Hungarian Ministry to open a second pharmacy in the area outlasted four years (Fabritius 1986; Roth 1976). On the other hand pharmacies in rural areas or small towns were a necessity. Ludwig Binder, native to Mediaş (Mediasch, Sibiu County) received on April 4th, 1878, from the Hungarian Internal Ministry, the authorization to open his one pharmacy with personal rights (no. 21.2525/1878) in Mediaş (Acker, 1970). The „Eagle” Pharmacy („Zum Adler”) was officially opened on November 11, 1878, in a small house on, that period, Lenin Street, number 44. Binder practiced for 32 years, until 1910 and he sold the pharmacy to chemist Walter Metz with the approval of the Hungarian Internal Ministry (no. 14005, 21st of February 1910) (Acker 1970; Fabritius 1986; Paveleanu 2004; Roth 1973). It seems that the medicinal herbarium received from Oberth Joseph has not been left to the new owner, as would be thought, as the data found in the herbarium does not mention Walter Metz (Inventory Book). Binder was an active member of the Chemists Association from Budapest (Pest Gremial), until he sold his pharmacy (Sibiu Archives). The Chemists Associations (Chemists Gremial) was founded in 1809 with the aim of providing a more serious and professional training to students, to oversee the pharmacies and drugstores operations and restrict the number of pharmacies especially in the urban areas (Roth 1976). For the same purposes, the Romanian authority divided Transylvania area into six regions across twenty-four districts and later in four districts without considering the administrative division of the country. Each region had a Gremial President and each district headquarters or branch-association was led by a secretary or Registrar (Roth 1976). Ludwig Binder was a member of the Medias Gremial association, Sibiu district administrated by the pharmacist Karl Pissel. The numerous association meetings held at district level, tied a friendship between pharmacist Ludwig Binder and the president of the association Karl Pissel, confirmed by the correspondence between the two (Sibiu Archives). Karl Pissell was

between the years 1903 and 1934, the owner of the “Crown” („Zur Krone”) in Sibiu.

The friendship between the two is what brought the medicinal herbarium in Sibiu. Perhaps, this Herbarium was a gift made by Binder to Pissell, when he reached the retirement age and sold his pharmacy. No document specifies when exactly the medicinal plant collection was brought to Sibiu, but it is very likely that Binder held it until 1918, when, because of the historical events in Romania, the association was dissolved. In the museum inventory books is noted that the herbarium did not come from the State Pharmacy no. 20, as many other historical pharmaceutical pieces did, but from the old "Crown" pharmacy, mentioned above (Inventory book). The „Crown” pharmacy was, in chronological order, the fourth opened in Sibiu with private owner (no. 25032), later on, with the implementation of the 1949 nationalization decree, it became a state pharmacy (Paveleanu 2004).

After the nationalization the Ministry of Health decided that during 1950 and 1951, all the historical pieces from the pharmacies or other health institutions from the country to be sent to the Brukenthal Museum from Sibiu. Thus, along with other pharmaceutical artifacts that became on 10.06.1951 part of the museum heritage, the *Herbarium Vivum* was assigned the inventory number F 2351/ F 817 (Inventory Book). In the museum inventory book at the origin of the herbarium is mentioned the "The Eagle" Pharmacy from Sebeş. Oberth's *Herbarium Vivum* remained in storage for 20 years, in the basement of the Brukenthal National Museum. In 1972 the Museum of Pharmacy was inaugurated in Sibiu, and the herbarium was transferred to this new location. Although herbarium was transferred in 1961 from the pharmacy museum's inventory to the Botanical collection inventory according to the accounting note 48/61, it still remained in the pharmaceutical collection. The inventory of the pharmaceutical collection done on the 30th of April 1970 registers the herbarium as part of the pharmaceutical history collection. After a period of reorganization the herbarium was officially handed to the Natural History Museum.

Conclusions

Herbarium Vivum is a medicinal herbarium from the 19th century that was used, at that time, as a medicinal plant guide. The Herbarium Book contains 130 medicinal plant species belonging to 35 families and 100 genera (tab.1.). The predominant plant families in this herbarium are *Fabaceae* and *Asteraceae*. Checking and updating

the scientific names of the medicinal plants revealed that most species are also present in Romania and a very small number are characteristic to other geographical areas. The plant species collected from other areas are: *Crepis discoridis* L., *Ferula communis* L., *Succisa kamerunensis* Engler ex Mildbraed and *Hesperis inodora* L., *Pseudolysimachion spurium* (L.) Rauschert.

Approximately 60% of the herbarium's plants are no longer used in today's traditional medicine, and future studies of these plants are recommended. The species identified in the herbarium whose medicinal use is unknown are: *Medicago prostrata* Jacq., *Lappula squarrosa* (Retz.) Dumort., *Cytisus nigricans* L., *Bifora testiculata* (L.) Spreng., *Monotropa hypopitys* L., *Cephalanthera longifolia* (L.) Fritsch, *Serratula tinctoria* L., *Leontodon hispidus* L., *Veronica anagallis – aquatica* L., *Gymnadenia odoratissima* (L.) Rich., *Trifolium ochroleucon* Huds., *Triglochin maritima* L., *Trifolium rubens* L., *Trifolium alpestre* L., *Lathyrus niger* (L.) Bernh., *Rhinanthus glacialis* Personnat., *Silene viscaria* (L.) Jess., *Inula hirta* L., *Campanula patula* L., *Pyrola media* Sw., *Cerinthe minor* L., *Vicia pannonica* Cranz, *Anthericum ramosum* L., *Veronica austriaca* subsp. *teucrium* L., *Nigella arvensis* L., *Dianthus carthusianorum* L., *Vicia cracca* L., *Galium glaucum* L., *Platanthera bifolia* (L.) L.C.Rich., *Persicaria bistorta* (L.) Samp., *Hypochaeris maculata* L., *Astragalus monspessulanus* subsp. *monspessulanus* L., *Plantago maritima* L., *Hesperis inodora* L., *Isopyrum thalictroides* L., *Tanacetum corymbosum* (L.) Sch.Bip., *Campanula spatulata* Sm., *Iris*

sibirica L., *Melampyrum arvense* L., *Orobanche caryophyllacea* Sm., *Salvia pratensis* L., *Lysimachia punctata* L., *Dactylorhiza incarnata* subsp. *incarnata* L. (Soó), *Barbarea vulgaris* R.Br., *Sisymbrium strictissimum* L., *Pseudolysimachion spurium* (L.) Rauschert, *Lathyrus vernus* (L.) Bernh., *Myosotis sylvatica* Hoffm.).

Due to the pharmaceutical industry development in artificially synthesizing substances that in the past were obtained only from plants and because of the quantitative reduction of species from the wild flora, 31, 53 % of the herbarium species are no longer used today. The following species were identified also in the Museum of Pharmaceutical History collection: *Dorycnium pentaphyllum* Scop., *Alyssum alyssoides* (L.) L., *Gladiolus communis* L., *Chaerophyllum bulbosum* L., *Chamaecytisus hirsutus* (L.) Link., *Leucanthemum vulgare* Lam., *Centaurea scabiosa* L., *Ornithogalum narbonense* L., *Lathyrus pratensis* L., *Veronica spicata* L., *Medicago falcata* L., *Stachys recta* L., *Sanicula elata* Buch.-Ham. ex D. Don., *Geranium pratense* L., *Prunella laciniata* (L.) L., *Anthriscus cerefolium* (L.) Hoffm., *Thalictrum aquilegifolium* L., *Filipendula vulgaris* Moench, *Anthemis tinctoria* L., *Scrophularia nodosa* L.

This high percentage includes plants used in skin diseases followed by plants used in digestive and respiratory diseases (Fig. 5).

Aknoledgements. The authors are grateful to Ana Maria Mesaroş for constructive comment on our manuscript and suggestions improving the language for the text. The authors would like to thanks to anonymous reviewer for helpful comment that improved the manuscript.

REFERENCES

- Acker 1970 Acker Michael, *Despre trecutul farmaciștilor din Sebeș și despre farmaciștii sebeșeni din trecut*. In: *Apulum* VIII, Alba-Iulia (1970), p. 271-281.
- Alexan et al. 1988 Alexan, Mircea, Bojor Ovidiu, Crăciun Florentin, *Flora medicinală a României*. In: *Editura Ceres*, București (1988), p. 224.
- Ban 2007 Ban Minodora, *Evidences regarding the medication based on resin substances of vegetal origin in the collection of the Pharmacy History Museum in Sibiu*. In: *Brukenthal Acta Musei* II.3, Sibiu (2007), p. 145-148.
- Ciocârlan 2009 Ciocârlan Vasile, *Flora ilustrată a României. Pteridophyta et Spermatophyta*. In: *Editura Ceres*, București (2009).
- Crișan 1996 Crișan Eva, *Materia Medica de Transilvania*. In: *Bibliotheca Musei Napocensis* XIV, Cluj-Napoca (1996).
- Drăgulescu 1976 Drăgulescu Constantin, *Note etnobotanice de la Păuca (jud. Sibiu) și considerații asupra plantelor medicinale folosite de poporul român*. In: *Studii și comunicări* 20, Sibiu (1976), p. 95- 112.
- Gheorghiu 1981 Gheorghiu Emil, *Preliminariile învățământului farmaceutic românesc*. In: Brătescu G. (eds), *Trecut și viitor în medicină. Ed. Medicală*, București (1981), p. 345-354.
- Fabritius 1986 Fabritius Guido, *Beiträge zur Geschichte der deutschen Apotheken und Apotheker in Sibenbürgen*, Stuttgart (1986), p. 425.
- Orient 1926 Orient Julius, *Aus pharmazeutischer Vergangenheit Siebenbürgens und des Banates. Gesellschaft für Geschichte der Pharmazie* (1926), p. 58.
- Oroian 2009 Oroian Silvia, *Flora și vegetația satelor săsești din sud-estul Transilvaniei*. In: *Editura University Press*, Târgu-Mureș (2009).
- Oroian 2011 Oroian Silvia, *Botanică Farmaceutică*. In: *Editura University Press*, Târgu-Mureș (2011).
- Panțu 1929 Panțu Zach C., *Plantele cunoscute de poporul Român. Vocabular botanic cuprinzând numirile române, franceze, germane și științifice*. In: *Editura Casei Școalelor* II, București (1929).
- Paveleanu 2004 Paveleanu Teodor, *Istoria farmaciilor din Transilvania III*. In: *Editura Universității Transilvania*, Brașov (2004).
- Pop, Olos 2004 Pop Maria, Olos Elisabeta, *Remedii din Farmacia Naturii II*. In: *Ed. Fiat Lux*, București (2004).
- Roth 1970 Roth Francisc, *Primele farmacii din Sibiu și evoluția lor până în secolul al XIX-lea. Studii și comunicări* 15, Sibiu (1970), p. 321-336.
- Roth 1973 Roth Francisc, *Registrul matricol al fostei asociații a farmaciștilor regiunii a VI-a Transilvania districtul 21 recte 22 Sibiu, de la înființare până în anul 1919 (încercare de reconstituire a unui document pierdut)*. *Studii și comunicări* 18, Sibiu (1973), p. 229-262.
- Roth 1976 Roth Francisc, *Organizarea farmaceutică din secolele trecute în Transilvania și în special la Sibiu*. In: *Studii și comunicări* 20, Sibiu (1976), p. 347-355.
- Roth- manuscript Roth Francisc, *Repertoriul fișierelor de farmacii și farmaciști din Ardealul de sud până la sfârșitul secolului al XIX-lea*. (Manuscript - Muzeul de Istoria Farmaciei). *Farmacopeea Română*. Ed. a X-a, In: *Editura Medicală*, București (1993).
- Farmacopeea Română 1993
- Decree No. 134/1949 Decretul privind naționalizarea unităților sanitare. Decretul nr. 134, 2.04.1949.
- Inventory Book Registrul de inventariere a patrimoniului Muzeului de Istorie a Farmaciei.
- Accounting note No. 48/61 Nota contabilă a Muzeului Național Brukenthal Sibiu, nr.48 din anul 1961.
- Archives- Sibiu Arhivele Naționale de Stat, Direcția Județeană Sibiu, Colecția Brukenthal, KK 1-31, no. 217
- Direction
- The Plant List *The Plant List* (2010). Version 1. Published on the Internet; <http://www.theplantlist.org/> (accessed 15th May, 2012).
- Taxonomy browser NCBI Taxonomy browser- <http://www.ncbi.nlm.nih.gov/> (accessed 15th of May 2012).

LIST OF ILLUSTRATIONS

- Tab. 1.** List of plant genera from *Herbarium Vivum*.
- Fig. 1.** The Herbarium Book from the Natural History Museum- Sibiu.
Fig. 2. Herbarium's cover- the year and the name of the collector can be seen.
Fig. 3. The herbarium's inside note.
Fig. 4. *Medicago prostrata* Jacq.- plant from Herbarium used in traditional medicine.
Fig. 5. The diseases for which were used the plants found in the herbarium.

LISTA ILUSTRAȚIILOR

- Tab. 1.** Clasificarea speciilor din *Herbarium Vivum*.
- Fig. 1.** Ierbar de tip carte din colecțiile Muzeului de Istorie Naturală din Sibiu.
Fig. 2. Coperta ierbarului - se poate vedea anul și numele celui care l-a realizat.
Fig. 3. Notele înscrise în interiorul ierbarului.
Fig. 4. *Medicago prostrata* Jacq.- plantă din ierbar folosită în medicina tradițională.
Fig. 5. Utilizarea medicinală a plantelor din ierbar.

Tab. 1. List of plant genera from *Herbarium Vivum*

Family	Genera	Sp.no.	Family	Genera	Sp.no.
Asclepiadaceae	Vincetoxicum	1		Chamaecytisus	20
Hyacinthaceae	Ornithogalum	1		Dorycnium	
Asparagaceae	Anthericum	1		Genista	
Asteraceae	Achillea	17		Lathyrus	
	Anthemis			Medicago	
	Calendula			Ononis	
	Centaurea			Trifolium	
	Chrysanthemum			Vicia	
	Crepis		Fumariaceae	Corydalis	1
	Eupatorium		Geraniaceae	Geranium	2
	Hypochaeris		Iridaceae	Gladiolus	2
	Inula	Iris			
	Leucanthemum	Juncaginaceae	Triglochin	1	
	Knautia	Lamiaceae	Marrubium	8	
	Leontodon		Mellitis		
	Matricaria		Prunella		
	Serratula		Salvia		
	Tanacetum		Scutellaria		
	Tragopogon		Stachys		
	Tussilago	Malvaceae	Althaea	1	
Apiaceae	Anthriscus	7	Orchidaceae	Anacamptis	5
	Bifora			Cephalanthera	
	Chaerophyllum			Dactylorhiza	
	Ferula			Gymnadenia	
	Heracleum			Platanthera	
	Pimpinella			Orobanchaceae	
	Sanicula		Orobanche		
	Berberidaceae		Berberis	1	Rinanthus
Boraginaceae	Cerinthe	6	Papaveraceae	Fumaria	1
	Echium		Plantaginaceae	Digitalis	7
	Lappula			Veronica	
	Myosotis			Pseudolysimachion	
	Pulmonaria		Plumbaginaceae	Limonium	1
	Symphytum		Polygonaceae	Persicaria	3
				Rumex	
Brassicaceae	Alyssum	7	Primulaceae	Lysimachia	2
	Barbarea		Ranunculaceae	Anemone	7
	Capsella			Clematis	
	Hesperis			Helleborus	
	Sinapis			Isopyrum	
	Sisymbrium			Nigella	
	Campanulaceae			Campanula	
Caryophyllaceae	Dianthus	5		Thalictrum	
	Silene		Rosaceae	Filipendula	1
	Stelaria		Rubiaceae	Galium	3
	Succisa		Scrophulariaceae	Scrophularia	3
Chenopodiaceae	Chenopodium	1			
Cornaceae	Cornus	1		Verbascum	
Ericaceae	Monotropa	2	Solanaceae	Hyoscyamus	1
	Pyrola		Valerianaceae	Valeriana	1
Euphorbiaceae	Euphorbia	3	Violaceae	Viola	1
Fabaceae	Anthyllis		Total: 35	100	130
	Astragalus				



Fig. 1. The Herbarium Book from Natural History Museum- Sibiu.



Fig. 2. Herbarium's cover- the year and the name of the collector can be seen.



Fig. 3. The herbarium's inside note.



Fig. 4. *Medicago prostrata* Jacq.- plant from Herbarium used in traditional medicine.




Fig. 5. The diseases for which were used the plants found in the herbarium.

**PROTECTED CULTURE STUDY OF CHINESE CABBAGE
(BRASSICA CAPESTRIS VAR. PEKINENSIS) VARIETIES AND HYBRIDS COLLECTION
GROWN IN THE TRANSYLVANIAN TABLELAND SPECIFIC CONDITIONS**

Enikő LACZI*
Alexandru Silviu APAHIDEAN*

Abstract. The Chinese cabbage (*Brassica campestris* var. *pekinensis*) collection of varieties and hybrids was established in the spring of 2011, in the polyethylene tunnel which belongs to the Vegetable Growing Department from the University of Agricultural Sciences and Veterinary Medicine from Cluj-Napoca. In this collection were studied and analyzed four hybrids (Michihli, Kingdom 80, Nepa F1 și Vitimo F1) and a variety (Granat). During the vegetation period, measurements were made regarding the growing and the development of these plants and upon the obtained yield. Using the recorded data was followed the evolution of plants height, diameter and number of leaves from planting until harvest, were analysed the interactions between the studied characteristics, but also the influence of hybrid upon the yield of Chinese cabbage. The processing of recorded data at harvest shows that the obtained yield varied between 77.40 and 110.40 t/ha, the Kingdom 80 hybrid reaching the highest yield. The obtained yields are satisfying, considering that the culture was an ecological one, no chemical products such as fertilisers or substances for prevention and control of pests and diseases were used.

Keywords: heading Chinese cabbage, protected culture, Chinese cabbage varieties, Chinese cabbage hybrids, Transylvanian Tableland.

Rezumat. Colecția de soiuri și hibrizi de varză chinezească (*Brassica campestris* var. *pekinensis*) a fost înființată în primăvara anului 2011, într-un solar care aparține Disciplinei de Legumicultură din cadrul Universității de Științe Agricole și Medicină Veterinară din Cluj-Napoca). În colecție s-a studiat un număr total de cinci cultivaruri, dintre care patru erau hibrizi, unul fiind un soi. Pe tot parcursul perioadei de vegetație s-au realizat măsurători în ceea ce privește creșterea și dezvoltarea plantelor, dar și asupra producției obținute. Utilizând datele înregistrate, s-a urmărit evoluția înălțimii și diametrului plantelor, a numărului de frunze de la plantare la recoltare; s-au analizat posibilele corelații care puteau exista între caracteristicile studiate, dar și influența hibrizului asupra producției de varză chinezească. Prelucrarea datelor înregistrate la recoltare, arată faptul că producția a variat între 77.40 și 110.40 t/ha, cea mai mare producție fiind realizată de către plantele aparținând hibrizului Kingdom 80. Producțiile obținute sunt satisfăcătoare având în vedere caracterul ecologic al culturii, nefolosindu-se îngrășăminte chimice sau diferite substanțe pentru prevenirea sau tratarea bolilor sau a dăunătorilor.

Cuvinte cheie: varza chinezească de căpățână, cultură protejată, soiuri de varză chinezească, hibrizi de varză chinezească, Podișul Transilvaniei.

Introduction

Chinese cabbage (*Brassica campestris* var. *pekinensis*, syn. *Brassica rapa* var. *pekinensis*) (Fig. 1 and Fig. 2), is a delicious vegetable that has been used extensively in Asian cooking for centuries. Large savoyed leaves with thick succulent midribs possess a sweet taste and crisp texture when eaten raw and it has a flavour somewhat milder than cabbage when cooked (Shattuck, Shelp 2004).

The development of Chinese cabbage in China was parallel with the European cabbage; both have evolved by cultivation from wild ancestors, both have been important foods since the remote past, and both now exist in numerous varieties which can be bought all the year round (Davidson, Jaine 2006).

Heading Chinese cabbage is a cold season, annual vegetable. It grows best under temperatures of 15–20°C degrees (Kalb, Chang 2005).

* University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, eniko.laczi@yahoo.com

These plants are amongst the fastest growing of all leafy vegetables, in favourable conditions heads can be cut ten weeks after sowing loose-headed types two to three weeks sooner, while seedlings four to five weeks after sowing (Larcom 2003).

The yield varies between 30 to 50 t/ha (Toxopeus, Baas 2004) or even between 40 to 100 t/ha (Choux, Foury 1994) according to the used hybrid and time and place of the culture).

Even if in our country is a less known vegetable, the Chinese cabbage is well known around the world and often used in western cuisine. The demand for this species, in Romania, could increase with the growth of the Asian population percentage (Ciofu *et al.* 2007), so it is very important to find an appropriate culture system for this species.

Material and method

The research took place in the polyethylene tunnel from the experimental field which belongs to the Vegetable Growing Department from the University of Agricultural Sciences and Veterinary Medicine Cluj - Napoca, in the spring of 2011.

The main purpose of this experiment was to study the behaviour of some of Chinese cabbage varieties in the Transylvanian Tableland specific conditions.

To achieve the objectives of this experiment a collection of varieties and hybrids was organized, which involved the following varieties:

- Michihli
- Kingdom 80
- Granat
- Nepa F1
- Vitimo F1

Each variety was placed into three repetitions.

The seeding was made in 25th February, the seeds were sown, one by one, in small nutrient pots and were transplanted in bigger pots in stage of 3-4 true leaves, in 21st of March, while planting was realized in 30th of the same month in the polyethylene tunnel.

During the vegetation period, there were not necessary any treatments or fertilizations. Harvest was realized in 18th of May at Granat variety and Nepa F1 and Michihli hybrids, in 26th of the same month at Kingdom 80 and 1st June at Vitimo F1 hybrid.

During growing season observations were made regarding plants growth and development (these were made at planting, at one month after planting and at harvesting), and on obtained production to.

Results and discussions

Seedlings development degree at planting

Fig. 3 shows that Michili hybrid registered the highest values of height (15.33 cm), diameter (19.00 cm), number of leaves (in average 7.67) and plants weight (10.67 g). On the other side is situated Vitimo hybrid, with the lowest height (10.67 cm) and plants diameter (19.00 cm). Seedlings from hybrids Kingdom 80, Nepa and Granat had an average weight of 10.17 g.

Plants development at one month after planting:

Measurements made one month after planting reveals the fact that plants height varied between 23.17 cm (at Kingdom 80 hybrid) and 43.83 cm (at Michihli hybrid), while their diameter between 46.50 cm (at Vitimo hybrid) and 58.33 cm (at Nepa hybrid). At this point of the vegetative period plants had between 14 (Nepa hybrid) and 21 leaves (Kingdom 80 hybrid) (Fig. 4).

Plants development before harvest

The highest plants (with an average height of 62.17 cm) and in the same time the fewest leaves (in average only 24.83) were measured at Michihli hybrid. Plants from Kingdom 80 hybrid formed the highest number of leaves, most than 51 leaves per plants, these being the most heaviest plants from the experiment (with an average yield of 1.69 kg).

Vitimo hybrid revealed with a low height (33.83 cm) and diameter (46.83 cm) but a high weight (an average of 1.32 kg) and a significant number of leaves (most than 49 leaves per plants) (Table 1).

Correlation between total weight and total number of leaves

The correlation coefficient in this case has a value of 0.82, which is lower than p (5%), so there is no relationship between the two studied factors. This means that the plants weight doesn't depend on the number of leaves, from statistical point of view (Fig. 5).

Cabbage heads development at harvest

The height of the harvested cabbage heads varied between 33.33 cm (at Vitimo hybrid) and 60.50 cm (at Michihli hybrid), the diameter between 29.17 cm (at Vitimo hybrid) and 43.33 cm (at Nepa hybrid).

The highest head weight was registered at Kingdom 80 hybrid (1.38 kg), in the other side with the lowest weight being situated hybrid Nepa, with an average weight of only 0.91 kg.

Regarding the number of leaves, it can be observed that a high number of leaves were formed by plants which belonged to Kingdom 80 hybrid (an average of 43.33 leaves), while with only 19.83 leaves stand out Michihli hybrid.

These results underlines the fact that Kingdom 80 and Vitimo hybrids forms very compact heads, which are not very tall, but are heaviest, than the other ones, the so called loose headed types (Table 2).

Correlation between head weight and height, diameter and number of leaves from the head

The value of p (5%), for five studied cases, is 0.88. Due to the fact that the correlation coefficient values are 0.45 (at the correlation between weight and height), 0.22 (at the correlation between weight and diameter), and 0.62 (at the correlation between weight and number of leaves), all of them being lower than 0.88, between these characteristics there is no interaction from the statistical point of view, therefore the plants weight does not depend on the plants height, diameter or the number of leaves (Fig. 6).

Comparison between total plants weight and heads weight

In Fig. 7 can be observed that between total plants weight and the head weights are differences ranging between 120 g, at Granat variety and 320 g, at Vitimo hybrid. This is a very important characteristic because after harvest only the compact heads can be send to market, so the outer leaves are cut away, therefore the total yield depends on the size of this lost.

Correlation between total and cabbage head weight

The correlation coefficient between the two studied characteristics had a value of 0.94. By comparing this value to the values of p (5%)=0.88, respectively p (1%)=0.96, the conclusion is, that between the two weights exists a very significant positive correlation, which indicates a strong dependence between the studied factors. With other words, with the increasing of the total weight, the head weight is increasing to (Fig. 8).

Leaf layout

The total number of leaves varied between 24.83 (at Michihli hybrid) and 51.83 (at Kingdom 80). The rosettes were formed in average from 4.67 leaves, at Granat variety and 9.00 leaves, at Vitmo hybrid.

The number of leaves which formed the heads varied according to the used variety, thereby at hybrid Michihli, the cabbage heads were formed by only 19.33 leaves, while those from Kingdom 80 had more than 43 leaves (Fig. 9).

Results regarding the bolting of the plants

The average bolting percentage, at one month after planting was 4.17%. The highest number of bolted plants was registered at hybrids Michihli and Nepa, where, at this moment, already 8.33% of the plants have bolted.

At harvest, more than 54.17% of the plants flowered at Michihli hybrid, which was followed by Granat variety, with a bolting percentage of 37.50%. Best results from this point of view were registered at Vitimo hybrid, where no plants emitted flowering stalks in the vegetative period.

The average bolting percentage, at harvest, was 24.17% (Fig. 10).

The cultivar influence upon the obtained yield

Data presented in table 3 shows that, in this experience, the yield of the five used cultivars varied between 72.40 t/ha, at Nepa hybrid and 110.40 t/ha, at Kingdom 80 hybrid. Compared to Granat variety, which was taken as control variant, only Kingdom 80 hybrid realized a higher yield. The difference, 22.27 t/ha, is very significant positive from the statistical point of view. Hybrids Michihli and Nepa realized lower yields than the control variant, with 10.93 and 15.73 t/ha, differences which are significant and distinct significant negatives.

The average yield of the five studied cultivars was 85.63 t/ha. Compared to this, a very significant positive difference was recorded at Kingdom 80 hybrid, where the yield increase was 28.9%. The lowest yield, 72.40 t/ha, was obtained at Nepa hybrid, this was lower with 13.23 t/ha than the average production, the difference being distinct significant negative.

Conclusions

The best developed seedlings were those which belonged to the Michihili variety, while at harvest those from Kingdom 80 hybrid registered the highest values considering the weight and number of leaves

By analyzing the interaction between some characteristics, the conclusion is that there exist a strong correlation only between head and total weight.

The loose headed cultivars had higher heights, while those which form a more compact head had higher weights and more leaves.

The average bolting percentage was almost 25%, while at hybrid Vitimo no plant emitted flowering stalks.

The obtained yields varied between 72.40 t/ha (at Nepa hybrid) and 110.40 t/ha (at Kingdom 80 hybrid).

The conclusion of this experiment is that Chinese cabbage can be easily cultivated in this area, in protected cultures.

REFERENCES

- | | |
|--------------------------|---|
| Chaux, Foury 1994 | Chaux Claude, Foury Claude, <i>Production légumières</i> , In: <i>TEC-DOC</i> , Paris, Paris (1994), p. 45-56. |
| Ciofu <i>et al.</i> 2007 | Ciofu Ruxandra, Stan Nistor, Popescu Victor, Chilom Pelaghia, Apahidean Silviu, Horgos Arsenie, Berar Viorel, Lauer Karl Fritz, Atanasiu Nicolae, <i>Tratat de legumicultură</i> . In: Editura Ceres, București (2004), p. 709-712. |
| Davidson, Jaine 2006 | Davidson Alan, Jaine Tom, <i>The Oxford Companion to Food</i> . In: <i>Oxford University Press</i> , Oxford, (2006), p.175. |
| Kalb, Chang, 2005 | Kalb Tom, Chang Lien-Chung, <i>Suggested cultural practices for heading Chinese cabbage</i> . In: <i>AVR DG pub.</i> , (2005), p. 5-642. |
| Larcom 2003 | Larcom Joy, <i>The organic salad garden</i> . In: <i>Frances Lincoln Limited</i> . London (2003), p. 31-33. |
| Shattuck, Shelp 2004 | Shattuck V., Shelp B., <i>Chinese Cabbage Production in Southern Ontario</i> . In: <i>Department of Horticultural Science, University of Guelph</i> , Guelph (2004), p. 1-7. |
| Toxopeus, Baas 2004 | Toxopeus H., Baas J., <i>Ressources végétales de l'Afrique tropicale 2</i> . In: <i>Fundation Prota / Backhuys Publishers / CTA</i> , Wageningen, (2004), p.162-169. |

LIST OF ILLUSTRATIONS

- Fig. 1.** Chinese cabbage (*Brassica campestris* var. *pekinensis*, syn. *Brassica rapa* var. *pekinensis*)
Fig. 2. Chinese cabbage (*Brassica campestris* var. *pekinensis*, syn. *Brassica rapa* var. *pekinensis*)
Fig. 3. Seedlings development at planting
Fig. 4. Plants development one month after planting
Fig. 5. Correlation between total number of leaves and total plants weight
Fig. 6. Correlation between head weight and height, diameter and total leaf number from cabbage head
Fig. 7. Comparison between total plant and head weight
Fig. 8. Correlation between total plant and head weight
Fig. 9. Leaf layout on plants
Fig.10. Bolting percentage
Tab. 1. Plants development before harvest
Tab. 2. Cabbage heads development at harvest
Tab. 3. The influence of cultivar upon the Chinese cabbage yield

LISTA ILUSTRAȚIILOR

- Fig. 1.** Varză chinezească (*Brassica campestris* var. *pekinensis*, syn. *Brassica rapa* var. *pekinensis*)
Fig. 2. Varză chinezească (*Brassica campestris* var. *pekinensis*, syn. *Brassica rapa* var. *pekinensis*)
Fig. 3. Dezvoltarea răsadurilor în momentul plantării
Fig. 4. Dezvoltarea plantelor la o lună de la plantare
Fig. 5. Corelația între numărul total de frunze și greutatea totală a plantelor
Fig. 6. Corelația dintre greutatea căpățânii și înălțimea, diametrul și numărul de frunze din căpățânile de varză
Fig. 7. Comparatie între greutatea totală a plantelor și greutatea căpăpățânilor
Fig. 8. Corelație între greutatea totală a plantelor și greutatea căpăpățânilor
Fig. 9. Dispunerea frunzelor pe plante
Fig.10. Procentul de plante înflorite prematur
Tab. 1. Gradul de dezvoltare al plantelor înainte de recoltare
Tab. 2. Gradul de dezvoltare al plantelor la recoltare
Tab. 3. Influența cultivarului asupra producției de varză chinezească

LIST OF ILLUSTRATIONS



Fig. 1. Chinese cabbage (*Brassica campestris* var. *pekinensis*, syn. *Brassica rapa* var. *pekinensis*)



Fig. 2. Chinese cabbage (*Brassica campestris* var. *pekinensis*, syn. *Brassica rapa* var. *pekinensis*)




Fig. 3. Seedlings development at planting




Fig. 4. Plants development one month after planting




Fig. 5. Correlation between total number of leaves and total plants weight ($p(5\%) = 0,88$, $p(1\%) = 0,91$)




Fig. 6. Correlation between head weight and height, diameter and total leaf number from cabbage head
($p(5\%)=0.88$, $p(1\%)=0.91$)




Fig. 7. Comparison between total plant and head weight




Fig. 8. Correlation between total plant and head weight




Fig. 9. Leaf layout on plants




Fig. 10. Bolting percentage

Tab.1. Plants development before harvest

Variant		Plant			
		Height (cm)	Diameter (cm)	Total weight (kg)	Number of leaves
No.	Variety / Hybrid				
1	Michihli	62.17	68.17	1.13	24,83
2	Kingdom 80	35.50	61.00	1.69	51,83
3	Granat	55.00	58.67	1.22	27,17
4	Nepa F1	56.17	69.00	1.04	30,17
5	Vitimo F1	33.83	46.83	1.32	49,50
Average		48,53	60.73	1.28	36.70

Tab. 2. Cabbage heads development at harvest

<i>Variant</i>		Cabage head			
<i>No.</i>	<i>Variety / Hybrid</i>	<i>Height (cm)</i>	<i>Diameter (cm)</i>	<i>Total weight (kg)</i>	<i>Number of leaves</i>
1	Michihli	60.50	40.33	0.97	19,83
2	Kingdom 80	38.67	42.50	1.38	43,33
3	Granat	60.17	40.00	1.10	22,50
4	Nepa F1	57.33	43.33	0.91	25,00
5	Vitimo F1	33.33	29.17	1.00	40,50
<i>Average</i>		50,00	39.07	1.07	30.23

Tab. 3. The influence of cultivar upon the Chinese cabbage yield

<i>Variant</i>	<i>Average yield (t/ha)</i>	<i>Relative yield (%)</i>	<i>Difference (t/ha)</i>	<i>Significance</i>	<i>Relative yield (%)</i>	<i>Difference (t/ha)</i>	<i>Significance</i>
<i>Variety / Hybrid</i>							
Granat	88.13	100.0	0.00	Ct.	102.9	2.50	-
Kingdom 80	110.4	125.27	22.27	***	128.9	24.77	***
Michihli	77.20	87.60	-10.93	o	90.2	-8.43	-
Nepa F1	72.40	82.15	-15.73	oo	84.6	-13.23	oo
Vitimo F1	80.00	90.77	-8.13	-	93.4	-5.63	-
<i>Average</i>	85.63	-	-	-	100.0	0.00	C.t.
<i>LSD (p 5%)</i>				8.88			8.88
<i>LSD (p 1%)</i>				12.92			12.92
<i>LSD (p 0,1%)</i>				19.38			19.38

THE FLORA OF THE BREITE WOOD-PASTURE (SIGHIȘOARA, ROMANIA)

Kinga ÖLLERER*

Abstract. The Breite wood-pasture located on a plateau nearby Sighișoara (Schäßburg, Romania) is considered one of the most representative sites for this type of habitat in Central-Eastern Europe. As a result of the field study begun in 2005, a list of 476 vascular plant species and subspecies is provided together with considerations on their ecology, and a small discussion regarding dynamics, distribution and biodiversity conservation issues.

Keywords: vascular plants, Breite, wood-pasture, biodiversity conservation, Romania.

Rezumat. Pajiștea cu stejari seculari de pe platoul Breite de lângă Sighișoara (Schäßburg, România) este considerată una din cele mai reprezentative pentru acest tip de habitat din Europa Centrală și de Est. În urma studiilor de teren începute în anul 2005, a fost întocmită o listă cuprinzând 476 specii și subspecii de plante vasculare, prezentată împreună cu date privind ecologia acestora și o scurtă discuție privind aspecte de dinamică, distribuție și conservarea biodiversității.

Cuvinte cheie: plante vasculare, Breite, pajiște împădurită, conservarea biodiversității, România.

Introduction

Wood - pastures represented one of the determinant land-use practices across Europe throughout centuries, but nowadays these sites are prone to degradation and disappearance mainly because of the abandonment of traditional practices, a threat affecting semi-natural habitats across the world (Bergmeier *et al.* 2010).

Wood-pastures are bearers of a characteristic biodiversity and living documents of land-use and cultural history. These deliberately shaped semi-natural habitats require specific management, its seizure or modification leading to major changes in their character (Rois-Díaz *et al.* 2006, Bergmeier *et al.* 2010). From the perspective of their conservation, the issues that are characteristically addressed target mainly the (veteran, standing or fallen) trees and the associated flora and fauna (Kirby *et al.* 1995, Rich *et al.* 2005, Stiven 2009). The ground vegetation is most often neglected, despite the fact that where not agriculturally improved, it can also be of considerable significance, increasing the biodiversity of the site, and the survival of the trees (Crofts, Jefferson 1999). It is therefore important to consider also the ground vegetation of wood-pasture habitats, addressing issues like the small-scale impacts, disturbance, grazing, browsing *etc.*, besides the

common research and monitoring usually focused on the trees. The first phase of such studies has to be a qualitative evaluation, including the identification of the species present in a specific site, which is also the topic of this paper.

While the flora and vegetation of the Sighișoara area is well documented due to several botanical works (Baumgarten 1816, 1846, Fronius 1858, Fuss 1866, Schur 1866, Simonkai 1886, Höhr 1914, Csűrös, Kovács 1962, Sanda *et al.* 1976), there are only a few studies discussing the Breite specifically. Out of these, the earliest is *Flora von Schäßburg*, published by Friedrich Fronius in 1858, in which the author provides a list of 1204 species, including 43 from the Breite. Heinrich Höhr, teacher and naturalist from Sighișoara, collected some plant species from the Breite, which are kept in his herbarium, publishing also several papers regarding this area (Höhr 1914, 1930, 1949).

In the last years the Breite and also other wood-pastures from the area benefited of biodiversity conservation initiatives and an increased scientific interest on which occasion several data were published (*e.g.* Bucșa 2007; Moga *et al.* 2007; Moga *et al.* 2009; Hartel, Moga 2010; Uhde 2011). A part of these studies included some aspects regarding the vegetation, while other papers and reports provided partial data regarding the vascular plants specifically (Akeroyd 2003, Mihai Eminescu Trust 2005, 2007, Mititiuc 2006, Csergő 2007, Muică 2009, Öllerer 2009, 2010, 2012, Öllerer *et al.* 2008a, b). Despite all these

* Institute of Biology – Romanian Academy, Ecology, Taxonomy and Nature Conservation Department
kinga.ollerer@gmail.com

contributions, the flora of the Breite remained only partially known. Other naturalists, including Eckhard Hügel (1908-1977) and more recently Erika Schneider-Binder and Hedwig Deppner wrote on different occasions about the biodiversity values of the Breite, mentioning also a few plant species.

The aim of this study was therefore to obtain a detailed view regarding the ground vegetation of the Breite, providing an up-to-date list of present plant species and make a baseline evaluation in order to understand plant species dynamics and distribution.

Study area and methods

Based on the data provided by Bergmeier *et al.* (2010) regarding wood-pasture habitats from Europe, the Breite is one of the largest and most representative enclosed wood-pastures in Central and Eastern Europe.

The Breite is situated on a 500 m (504 m in the northern part and 530 m in the southern part) elevation plateau bearing the same name, which is reflecting very well the open character of this habitat (*Breite* = „wideness” in German) nearby the town of Sighișoara (Schäßburg, Romania, center point coordinates: lat. 46.2011, lon. 24.7606). The 133 ha open area, with over 630 scattered or locally grouped mature oaks (*Quercus robur* and *Q. petraea*), out of which around 400 are several centuries old, is surrounded by a mixed deciduous Sessile Oak and Hornbeam with Beech forest (Mihai Eminescu Trust 2005, 2007, Fig. 1). The wood-pasture was formed partially from the opening of this forest and most probably also from maintenance plantings, which was a widespread land-use practice in this part of Transylvania since the Middle Ages (Oroszi 2004), reflected also by the relatively high number of such habitats in Southern Transylvania (Hartel, Moga 2010; Öllerer *et al.* 2010).

The Breite plateau is part of the Transylvanian Tableland and is located in the middle section of the Târnava Mare River. The soil types on the plateau are: brown forest soils, podzols and clayey soils, clayey-illuvial and alluvial (Mihai Eminescu Trust 2007). The high clay content and increased debasification rate resulted in a very compact and almost impermeable soil on the Breite. This explains the fact that the Breite was a humid, even marshy area in the past (Siegmund 1948). This character is kept nowadays only in its southern part due to the significant water loss following the

construction of a drainage system in the 1970s, that also resulted in the formation of small sites with excessive humidity in the drainage ditches (Mihai Eminescu Trust 2007).

The climate in this area is temperate continental; the annual mean temperature is 8.2°C, while the thermal amplitude between the minimum in January (-4.3°C) and the maximum in July (18.6°C) is 22.9°C. The nebulosity is high especially during winter and spring and precipitations are not uniform, being more abundant in April-October, when 70 % of the total annual rainfall is recorded. The rainiest months are May and June, with 90-100 l/sqm (Mihai Eminescu Trust 2007).

Besides the general threats that affect wood-pastures across Europe (lack of tree regeneration, change of use, abandonment and the resulting modifications of the vegetation and of the overall species composition - Bergmeier *et al.* 2010), the Breite was also affected by the results of improper management interventions that targeted the transformation of the area into a military airport and later into a crop field (Mihai Eminescu Trust 2007).

After the recognition of its scientific and historical values in the mid-1970s, the Breite became gradually protected, while its overall management for conservation purposes became an issue only in 2005 (Mihai Eminescu Trust 2005, 2007). Nowadays 70 ha of the Breite are considered as nature reserve under the name „Stejarii seculari de la Breite” (The Breite ancient oak tree reserve), being listed in the position 2.635 in the Romanian Law 5/2000 regarding protected areas. The entire plateau is included in the Natura 2000 European network of protected areas, in the Sighișoara-Târnava Mare SCI and the larger Podișul Hârtibaciului SPA sites, the latter partially overlapping the former. In 2005, the Romanian Academy, through the Commission for the Protection of Natural Monuments approved the extension of the protected area in order to cover the entire 133 ha of the plateau.

The study of the ground vegetation cover started in 2005, including both field studies and consulting historical and bibliographical data (*e.g.* Fronius 1858; Oroszi 2004; Teșculă, Goța 2007, ordinance surveys). The aim was to obtain a clearer view of the history of this site and of other wood-pastures in the area that might help to explain the presence and distribution of plant species and formulate recommendations for further management from a biodiversity conservation perspective.

In the following, the resulting list of species is provided including only those present in the wood-pasture (open) area of the Breite plateau. Plant species were identified with the help of illustrated guides in use for the Romanian flora (Săvulescu 1952-1976, Ciocârlan 2009). Nomenclature follows Flora Europaea (2001). Systematics follows Ciocârlan (2009). Data regarding the ecological characteristics of species were obtained from Ciocârlan (2009).

Results

List of species

PTERIDOPHYTA – ferns and horsetails

Equisetaceae

1. *Equisetum arvense* L.

Dennstaedtiaceae

2. *Pteridium aquilinum* (L.) Kuhn – HH (1914)¹

Aspleniaceae

3. *Athyrium filix-femina* (L.) Roth
4. *Dryopteris dilatata* (Hoffm.) A. Gray (*Dryopteris austriaca* Woyen.)
5. *Dryopteris filix-mas* (L.) Schott

SPERMATOPHYTA

PINOPHYTINA (*Gymnospermae*) – conifers

Pinaceae

6. *Larix decidua* Mill. cult.
7. *Picea abies* (L.) H. Karst. cult.
8. *Pinus nigra* J.F. Arnold cult.
9. *Pinus sylvestris* L. cult.

MAGNOLIOPHYTINA (*Angiospermae*) – flowering plants

Cls. Magnoliopsida (*Dicotyledonatae*)

Ranunculaceae

10. *Anemone nemorosa* L.
11. *Anemone ranunculoides* L.
12. *Ranunculus acris* L.
13. *Ranunculus auricomus* L.
14. *Ranunculus bulbosus* L.
15. *Ranunculus ficaria* L. s. l.
16. *Ranunculus polyanthemus* L.
17. *Ranunculus repens* L. s. l.
18. *Ranunculus strigosus* Schur

Papaveraceae

19. *Chelidonium majus* L.

Urticaceae

20. *Urtica dioica* L.

Juglandaceae

21. *Juglans regia* L.

Fagaceae

22. *Fagus sylvatica* L. – HH herb. (1908)²
23. *Quercus petraea* (Matt.) Liebl. – FF (1858) [949] (under *Q. sessiliflora* Salisb.)³
24. *Quercus robur* L. – FF (1858) [950] (under *Q. pedunculata* Ehrh.)

Betulaceae

25. *Betula pendula* Roth – HH herb. (1909)

Corylaceae

26. *Carpinus betulus* L. – HH herb. (1908)

Caryophyllaceae

27. *Arenaria serpyllifolia* L.
28. *Cerastium fontanum* Baumg. subsp. *fontanum* – FF (1858) [175] (under *Cerastium vulgatum* L.)
29. *Cerastium glomeratum* Thuill.
30. *Cerastium pumilum* Curtis subsp. *pumilum*
31. *Dianthus armeria* L. subsp. *armeria* – HH herb. (1910)
32. *Dianthus carthusianorum* L.
33. *Dianthus deltoides* L.
34. *Gypsophila muralis* L.
35. *Lychnis flos-cuculi* L. – FF (1858) [160]; HH herb. 1908
36. *Moehringia trinervia* (L.) Clairv.
37. *Myosoton aquaticum* (L.) Moench (*Stellaria aquatica* (L.) Scop.)
38. *Sagina procumbens* L.
39. *Scleranthus annuus* L.
40. *Scleranthus perennis* L.
41. *Silene latifolia* Poir. subsp. *alba* (Mill.) Greuter & Burdet
42. *Silene noctiflora* L.
43. *Stellaria graminea* L.
44. *Stellaria holostea* L.
45. *Stellaria media* (L.) Vill. s.l.
46. *Stellaria nemorum* L.
47. *Stellaria uliginosa* Murray (*Stellaria alsine* Grimm)

Amaranthaceae

48. *Amaranthus powellii* S. Watson
49. *Amaranthus retroflexus* L.
50. *Spergularia rubra* (L.) J. Presl & C. Presl

Chenopodiaceae

51. *Atriplex patula* L.
52. *Chenopodium album* L.

Polygonaceae

53. *Fallopia dumetorum* (L.) Holub
54. *Polygonum aviculare* L. s. l.
55. *Polygonum bistorta* L.
56. *Polygonum hydropiper* L.
57. *Polygonum lapathifolium* L.
58. *Polygonum minus* Huds.
59. *Polygonum persicaria* L.
60. *Rumex acetosa* L.
61. *Rumex acetosella* L. s. l. – HH herb. (1912)

62. *Rumex conglomeratus* Murray – MM (2006)⁴

63. *Rumex crispus* L.

64. *Rumex obtusifolius* L. s. l.

65. *Rumex sanguineus* L.

Crassulaceae

66. *Sedum sexangulare* L.

Rosaceae

67. *Agrimonia eupatoria* L. – OS (2009)⁵

68. *Crataegus monogyna* Jacq. – HH herb. (1908)

69. *Filipendula ulmaria* (L.) Maxim. subsp. *ulmaria*

70. *Filipendula vulgaris* Moench

71. *Fragaria vesca* L.

72. *Geum urbanum* L.

73. *Malus sylvestris* Mill. – HH herb. (1908)

74. *Potentilla anserina* L.

75. *Potentilla argentea* L.

76. *Potentilla erecta* (L.) Rausch.

77. *Potentilla recta* L.

78. *Potentilla reptans* L.

79. *Prunus cerasifera* Ehrh.

80. *Prunus spinosa* L.

81. *Pyrus pyrastra* Burgsd.

82. *Rosa canina* L. – FF (1858) [352]; OS (2009)

83. *Rosa micrantha* Borrer ex Sm.

84. *Rosa obtusifolia* Desv.

85. *Rubus caesius* L.

86. *Rubus hirtus* Waldst. & Kit.

87. *Rubus hirtus* Weihe & Nees

88. *Rubus idaeus* L.

89. *Rubus macrophyllus* Weihe & Nees

90. *Sanguisorba officinalis* L.

Fabaceae (Leguminosae)

91. *Anthyllis vulneraria* L. s. l.

92. *Astragalus glycyphyllos* L.

93. *Chamaespartium sagittale* (L.) P.E. Gibbs (*Genistella sagittalis* (L.) Gams) – FF (1858) [240]

94. *Coronilla varia* L.

95. *Galega officinalis* L.

96. *Genista tinctoria* L. subsp. *tinctoria*

97. *Lathyrus hirsutus* L.

98. *Lathyrus pratensis* L.

99. *Lathyrus vernus* (L.) Bernh.

100. *Lotus corniculatus* L.

101. *Medicago lupulina* L.

102. *Medicago minima* (L.) Bartal.

103. *Medicago sativa* subsp. *falcata* (L.) Arcangeli (*Medicago falcata* L.)

104. *Melilotus alba* Medik.

105. *Melilotus officinalis* (L.) Pall.

106. *Ononis arvensis* L.

107. *Robinia pseudacacia* L.

108. *Trifolium arvense* L.

109. *Trifolium aureum* Pollich

110. *Trifolium campestre* Schreb. – FF (1858) [271]

111. *Trifolium dubium* Sibth.

112. *Trifolium fragiferum* L. s. l.

113. *Trifolium hybridum* L. s. l.

114. *Trifolium medium* L. subsp. *medium*

115. *Trifolium montanum* L.

116. *Trifolium ochroleucon* Huds.

117. *Trifolium pratense* L. s. l.

118. *Trifolium repens* L.

119. *Vicia cracca* L.

120. *Vicia hirsuta* (L.) S.F. Gray

121. *Vicia sativa* subsp. *nigra* (L.) Ehrh. (*Vicia angustifolia* L.)

122. *Vicia sepium* L.

123. *Vicia tetrasperma* (L.) Schreb.

124. *Vicia villosa* Roth

Lythraceae

125. *Lythrum portula* (L.) D.A. Webb

126. *Lythrum salicaria* L. – MM (2006)

Onagraceae

127. *Circaea lutetiana* L.

128. *Epilobium angustifolium* L.

129. *Epilobium montanum* L.

130. *Epilobium palustre* L. – MM (2006)

131. *Epilobium tetragonum* L. subsp. *lamyi* (F.W. Schultz) Nyman

Cornaceae

132. *Cornus sanguinea* L. – HH herb. (?)

Loranthaceae

133. *Viscum album* L. – HH herb. (1908)

134. *Loranthus europaeus* Jacq.

Celastraceae

135. *Euonymus europaeus* L.

136. *Euonymus verrucosus* Scop.

Euphorbiaceae

137. *Euphorbia amygdaloides* L.

138. *Euphorbia angulata* Jacq.

139. *Euphorbia cyparissias* L.

140. *Euphorbia epithymoides* L. – FF (1858) [925]

141. *Euphorbia exigua* L.

142. *Euphorbia helioscopia* L.

143. *Euphorbia platyphyllos* L.

144. *Euphorbia serrulata* Thuill.

Rhamnaceae

145. *Frangula alnus* Mill.

Aceraceae

146. *Acer campestre* L.

147. *Acer negundo* L.

148. *Acer platanoides* L.

149. *Acer pseudoplatanus* L.

150. *Acer tataricum* L. – HH herb. (1913)

Oxalidaceae

151. *Oxalis acetosella* L.

152. *Oxalis fontana* Bunge (*Oxalis europaea* Jord.) – HH herb. (1908)
153. *Oxalis stricta* L. (*Oxalis dilenii* Jacq.)
- Geraniaceae**
 154. *Erodium cicutarium* (L.) L'Hér.
 155. *Geranium pratense* L.
 156. *Geranium pusillum* L.
 157. *Geranium robertianum* L. – HH herb. (1941)
- Balsaminaceae**
 158. *Impatiens noli-tangere* L.
- Polygalaceae**
 159. *Polygala comosa* Schkuhr
 160. *Polygala major* Jacq.
 161. *Polygala vulgaris* L.
- Apiaceae** (Umbelliferae)
 162. *Aegopodium podagraria* L. – HH herb. (1913)
 163. *Angelica sylvestris* L.
 164. *Anthriscus cerefolium* (L.) Hoffm.
 165. *Anthriscus sylvestris* (L.) Hoffm.
 166. *Carum carvi* L.
 167. *Chaerophyllum aromaticum* L.
 168. *Chaerophyllum bulbosum* L.
 169. *Chaerophyllum temulentum* L.
 170. *Daucus carota* L. subsp. *carota* – MM (2006)
 171. *Eryngium campestre* L.
 172. *Eryngium planum* L.
 173. *Heracleum sphondylium* L.
 174. *Pastinaca sativa* L. s. l.
 175. *Pimpinella major* (L.) Huds.
 176. *Pimpinella saxifraga* L.
 177. *Sanicula europaea* L.
 178. *Selinum carvifolia* (L.) L.
 179. *Torilis arvensis* (Huds.) Link s. l. – MM (2006)
- Hypericaceae**
 180. *Hypericum hirsutum* L. – FF (1858) [201]
 181. *Hypericum humifusum* L.
 182. *Hypericum perforatum* L.
- Malvaceae**
 183. *Alcea pallida* (Willd.) Waldst. & Kit.
- Violaceae**
 184. *Viola arvensis* Murray
 185. *Viola canina* L.
 186. *Viola hirta* L. – HH herb. (1908)
 187. *Viola mirabilis* L.
 188. *Viola reichenbachiana* Jord. ex Boreau
- Cistaceae**
 189. *Helianthemum nummularium* (L.) Mill. subsp. *obscurum* (Čelak.) J. Holub
- Brassicaceae** (Cruciferae)
 190. *Alliaria petiolata* (M. Bieb.) Cavara & Grande (*Alliaria officinalis* Andr. ex Bieb.)
191. *Barbarea vulgaris* R.Br. – HH herb. (1909)
192. *Capsella bursa-pastoris* (L.) Medik.
193. *Cardamine flexuosa* With.
194. *Cardamine impatiens* L.
195. *Cardamine pratensis* L. subsp. *pratensis*
196. *Cardaria draba* (L.) Desv. (*Lepidium draba* L.)
197. *Cardamine bulbifera* (L.) Crantz
198. *Descurainia sophia* (L.) Webb ex Prantl
199. *Rorippa palustris* (L.) Besser
200. *Rorippa pyrenaica* (Lam.) Rchb.
201. *Rorippa sylvestris* (L.) Besser subsp. *sylvestris*
202. *Sinapis arvensis* L.
203. *Sisymbrium officinale* (L.) Scop.
- Salicaceae**
 204. *Populus alba* L.
 205. *Populus tremula* L.
 206. *Salix alba* L.
 207. *Salix caprea* L.
 208. *Salix cinerea* L.
 209. *Salix fragilis* L.
 210. *Salix purpurea* L.
 211. *Salix triandra* L.
- Primulaceae**
 212. *Anagallis arvensis* L.
 213. *Anagallis foemina* Mill.
 214. *Lysimachia nummularia* L. – HH (1914)
 215. *Lysimachia vulgaris* L.
 216. *Primula veris* L.
- Gentianaceae**
 217. *Centaurium erythraea* Rafn. subsp. *erythraea*
 218. *Gentiana pneumonanthe* L. – HH herb. (1941)
- Asclepiadaceae**
 219. *Asclepias syriaca* L.
- Oleaceae**
 220. *Ligustrum vulgare* L. – HH herb. (1943)
- Convolvulaceae**
 221. *Convolvulus arvensis* L.
 222. *Cuscuta europaea* L.
- Boraginaceae**
 223. *Echium vulgare* L.
 224. *Myosotis arvensis* (L.) Hill subsp. *arvensis* – HH herb. (?)
 225. *Myosotis discolor* Pers.
 226. *Myosotis scorpioides* L.
 227. *Myosotis stricta* Link ex Roem. & Schult.
 228. *Myosotis sylvatica* Hoffm.
 229. *Pulmonaria mollis* Wulfen ex Hornem.
 230. *Symphytum officinale* L. subsp. *officinale*
 231. *Symphytum tuberosum* L. – FF (1858) [708]
- Verbenaceae**
 232. *Verbena officinalis* L. – HH herb. (1941)
- Lamiaceae** (Labiatae)

233. *Ajuga reptans* L.
234. *Ballota nigra* L.
235. *Clinopodium vulgare* L.
236. *Galeopsis bifida* Boenn.
237. *Galeopsis speciosa* Mill.
238. *Galeopsis tetrahit* L.
239. *Glechoma hederacea* L.
240. *Lamiastrum galeobdolon* (L.) Ehrend. & Polatschek – FF (1858) [836] (under *Galeobdolon luteum* Huds.)
241. *Lamium album* L.
242. *Lamium maculatum* L.
243. *Lamium purpureum* L.
244. *Lycopus europaeus* L.
245. *Lycopus exaltatus* L.f.
246. *Marrubium vulgare* L.
247. *Mentha arvensis* L.
248. *Mentha longifolia* (L.) Huds. – MM (2006)
249. *Mentha pulegium* L.
250. *Origanum vulgare* L.
251. *Prunella grandiflora* (L.) Scholler – MM (2006)
252. *Prunella laciniata* (L.) L.
253. *Prunella vulgaris* L. – OS (2009)
254. *Salvia glutinosa* L.
255. *Salvia pratensis* L.
256. *Salvia verticillata* L.
257. *Scutellaria galericulata* L.
258. *Stachys annua* (L.) L.
259. *Stachys officinalis* (L.) Trevis. (*Betonica officinalis* L.)
260. *Stachys sylvatica* L.
261. *Teucrium chamaedrys* L.
262. *Thymus glabrescens* Willd.

Callitrichaceae

263. *Callitriche palustris* L.

Plantaginaceae

264. *Plantago lanceolata* L. – MM (2006)
265. *Plantago major* L. s. 1.
266. *Plantago media* L. s. 1.

Scrophulariaceae

267. *Digitalis grandiflora* Mill. – HH herb. (1910)
268. *Euphrasia rostkoviana* Hayne – FF (1858) [791] (under *E. officinalis* L.)
269. *Euphrasia stricta* J.P. Wolff ex J.F. Lehm.
270. *Linaria vulgaris* Mill. – OS (2009)
271. *Melampyrum bihariense* A. Kern. – HH herb. (1943)
272. *Rhinanthus minor* L.
273. *Rhinanthus rumelicus* Velen.
274. *Scrophularia nodosa* L.
275. *Verbascum nigrum* L. subsp. *nigrum*
276. *Verbascum phlomoides* L.
277. *Veronica arvensis* L.
278. *Veronica beccabunga* L.

279. *Veronica chamaedrys* L. s. 1.
280. *Veronica hederifolia* L. subsp. *hederifolia*
281. *Veronica officinalis* L.
282. *Veronica persica* Poir.
283. *Veronica polita* Fr.
284. *Veronica serpyllifolia* L. subsp. *serpyllifolia* – FF (1858) [769]; HH herb. 1912

Solanaceae

285. *Atropa bella-donna* L. – HH (1914); HH herb. (?)
286. *Solanum dulcamara* L.
287. *Solanum nigrum* L.

Campanulaceae

288. *Campanula patula* L. subsp. *patula*
289. *Campanula persicifolia* L.
290. *Campanula rapunculoides* L.
291. *Campanula trachelium* L.

Rubiaceae

292. *Cruciata glabra* (L.) Ehrend.
293. *Cruciata laevipes* Opiz – HH herb. (?)
294. *Galium album* Mill. s. 1.
295. *Galium aparine* L.
296. *Galium mollugo* L.
297. *Galium odoratum* (L.) Scop. (*Asperula odorata* L.) – FF (1858) [474]; HH herb. 1908
298. *Galium palustre* L.
299. *Galium schultesii* Vest
300. *Galium verum* L.

Caprifoliaceae

301. *Lonicera xylosteum* L.
302. *Sambucus ebulus* L.
303. *Sambucus nigra* L.
304. *Viburnum opulus* L.

Valerianaceae

305. *Valeriana officinalis* L.

Dipsacaceae

306. *Dipsacus laciniatus* L.
307. *Knautia arvensis* (L.) Coult. – MM (2006)
308. *Succisa pratensis* Moench – HH herb. (1941)

Asteraceae (Compositae)

309. *Achillea collina* J. Becker ex Rchb.
310. *Achillea millefolium* L. – MM (2006); OS (2009)
311. *Achillea ptarmica* L.
312. *Anthemis ruthenica* M. Bieb.
313. *Arctium lappa* L.
314. *Artemisia absinthium* L.
315. *Artemisia vulgaris* L.
316. *Bellis perennis* L.
317. *Bidens tripartita* L. – HH herb. (1911)
318. *Carduus acanthoides* L.
319. *Carduus crispus* L.
320. *Centaurea jacea* L. – HH herb. (1908)

321. *Centaurea phrygia* L. subsp. *phrygia* (*C. austriaca* Willd.) – MM (2006)
 322. *Chamomilla recutita* (L.) Rauschert (*Matricaria chamomilla*)
 323. *Chamomilla suaveolens* (Pursh) Rydb. (*Matricaria discoidea* DC.)
 324. *Cichorium intybus* L. – OS (2009)
 325. *Cirsium arvense* (L.) Scop.
 326. *Cirsium canum* (L.) All.
 327. *Cirsium furiens* Griseb. & Schenk
 328. *Cirsium oleraceum* (L.) Scop. – HH herb. (1943)
 329. *Cirsium palustre* (L.) Scop.
 330. *Cirsium rivulare* (Jack.) All. – OS (2009)
 331. *Cirsium vulgare* (Savi) Ten. – MM (2006)
 332. *Conyza canadensis* (L.) Cronq. (*Erigeron canadensis* L.)
 333. *Crepis biennis* L.
 334. *Crepis capillaris* (L.) Wallr.
 335. *Erechtites hieracifolia* (L.) Rafin. ex DC.
 336. *Erigeron annuus* (L.) Pers. subsp. *annuus*
 337. *Erigeron annuus* (L.) Pers. subsp. *strigosus* (Muhl. ex Willd.) Wagenitz
 338. *Eupatorium cannabinum* L.
 339. *Filago vulgaris* Lam.
 340. *Filaginella uliginosa* (L.) Opiz subsp. *uliginosa* (*Gnaphalium uliginosum* L.)
 341. *Galinsoga parviflora* Cav.
 342. *Hieracium aurantiacum* L.
 343. *Hieracium cymosum* L. s. l.
 344. *Hieracium murorum* L. s. l. – FF (1858) [646]
 345. *Hieracium pilosella* L. s. l.
 346. *Hieracium umbellatum* L. s. l.
 347. *Hypochaeris radicata* L. – FF (1858) [625]
 348. *Inula britannica* L. – MM (2006)
 349. *Inula helenium* L.
 350. *Lactuca serriola* L.
 351. *Lapsana communis* L. subsp. *communis*
 352. *Leontodon autumnalis* L. subsp. *autumnalis* – MM (2006)
 353. *Leontodon hispidus* L.
 354. *Leucanthemum vulgare* Lam.
 355. *Logfia minima* (Sm.) Dumort. (*Filago minima* (Sm.) Pers.)
 356. *Matricaria perforata* Mérat
 357. *Mycelis muralis* (L.) Dumort.
 358. *Omalotheca sylvatica* (L.) Sch. Bip. & F.W. Schultz (*Gnaphalium sylvaticum* L.)
 359. *Picris hieracioides* L.
 360. *Pulicaria dysenterica* (L.) Bernh.
 361. *Pulicaria vulgaris* Gaertner
 362. *Rudbeckia laciniata* L.
 363. *Senecio aquaticus* Hill subsp. *barbareifolius* (Wimm. & Grab.) Walters (*Senecio erraticus* Bertol.)
 364. *Senecio erucifolius* L.
 365. *Senecio jacobaea* L.
 366. *Solidago canadensis* L.
 367. *Solidago gigantea* Aiton subsp. *serotina* (O. Kuntze) McNeill
 368. *Solidago virgaurea* L.
 369. *Sonchus oleraceus* L.
 370. *Tanacetum vulgare* L.
 371. *Taraxacum officinale* Weber
 372. *Telekia speciosa* (Schreber) Baumg.
 373. *Tragopogon pratensis* L. subsp. *orientalis* (L.) Čelak. (*Tragopogon orientalis* L.)
 374. *Tussilago farfara* L.
 375. *Xanthium strumarium* L. – OS (2009)
- Cls. LILIOPSIDA (*Monocotyledonatae*)**
- Alismataceae**
376. *Alisma plantago-aquatica* L. – HH herb. (1908)
- Liliaceae s. l.**
377. *Colchicum autumnale* L.
 378. *Maianthemum bifolium* (L.) F.W.Schmidt
 379. *Paris quadrifolia* L.
 380. *Polygonatum latifolium* (Jacq.) Desf.
 381. *Polygonatum multiflorum* (L.) All.
 382. *Scilla bifolia* L. s. l.
- Alliaceae**
383. *Allium oleraceum* L.
- Iridaceae**
384. *Iris pseudacorus* L. – HH herb. (1908)
- Orchidaceae**
385. *Dactylorhiza incarnata* (L.) Soó
 386. *Gymnadenia conopsea* (L.) R. Br.
 387. *Listera ovata* (L.) R.Br.
 388. *Orchis morio* L. subsp. *morio* – HH herb. (1908)
 389. *Orchis ustulata* L.
 390. *Platanthera bifolia* (L.) Rich.
- Juncaceae**
391. *Luzula campestris* (L.) DC.
 392. *Luzula luzuloides* (Lam.) Dandy & Wilmott
 393. *Juncus articulatus* L.
 394. *Juncus bufonius* L.
 395. *Juncus conglomeratus* L. – MM (2006)
 396. *Juncus effusus* L.
 397. *Juncus inflexus* L. – MM (2006)
 398. *Juncus tenuis* Willd.
- Cyperaceae**
399. *Carex acuta* L.
 400. *Carex acutiformis* Ehrh.
 401. *Carex appropinquata* A. Schumach.
 402. *Carex brizoides* L.
 403. *Carex caryophylllea* Latourr.
 404. *Carex cespitosa* L.
 405. *Carex depressa* Link
 406. *Carex divulsa* Stokes

407. *Carex elata* All.
 408. *Carex hirta* L.
 409. *Carex michelii* Host
 410. *Carex nigra* (L.) Reichenb.
 411. *Carex ovalis* Gooden.
 412. *Carex pallescens* L. – FF (1858) [1107]
 413. *Carex paniculata* L.
 414. *Carex pendula* Huds.
 415. *Carex pilosa* Scop.
 416. *Carex remota* L.
 417. *Carex spicata* Huds.
 418. *Carex sylvatica* Huds. – FF (1858) [1106]
 419. *Carex tomentosa* L. – FF (1858) [1113]
 420. *Carex vesicaria* L.
 421. *Carex vulpina* L.
 422. *Eleocharis palustris* (L.) Roem. & Schult.
 423. *Scirpus sylvaticus* L.

Poaceae (Gramineae)

424. *Agrostis canina* L.
 425. *Agrostis capillaris* L. (*Agrostis tenuis* Sibth.)
 426. *Agrostis gigantea* Roth subsp. *gigantea*
 427. *Agrostis stolonifera* L. – MM (2006)
 428. *Alopecurus aequalis* Sobol. – HH herb. (1943)
 429. *Alopecurus pratensis* L.
 430. *Anthoxanthum odoratum* L. – HH herb. (1908)
 431. *Apera spica-venti* (L.) P. Beauv.
 432. *Arrhenatherum elatius* (L.) P. Beauv. ex J. Presl & C. Presl subsp. *elatius*
 433. *Brachypodium sylvaticum* (Huds.) P. Beauv.
 434. *Briza media* L. – HH herb. (1911)
 435. *Bromus commutatus* Schrad.
 436. *Bromus erectus* Huds. s. l.
 437. *Bromus hordeaceus* L.
 438. *Bromus inermis* Leyss. – FF (1858) [1191]
 439. *Bromus japonicus* Thunb.
 440. *Bromus squarrosus* L.
 441. *Calamagrostis arundinacea* (L.) Roth
 442. *Calamagrostis canescens* (Weber) Roth
 443. *Calamagrostis epigejos* (L.) Roth
 444. *Cynosurus cristatus* L.
 445. *Dactylis glomerata* L. subsp. *aschersoniana* (Graebn.) Thell.
 446. *Deschampsia caespitosa* (L.) P. Beauv. s. l. – FF (1858) [1151] (under *Aira caespitosa* Muhl.)
 447. *Digitaria sanguinalis* (L.) Scop.
 448. *Echinochloa crus-galli* (L.) P. Beauv.
 449. *Elymus repens* (L.) Gould
 450. *Eragrostis minor* Host
 451. *Festuca arundinacea* Schreb. s. l.
 452. *Festuca drymeja* Mert. et W.D.J. Koch
 453. *Festuca gigantea* (L.) Vill.

454. *Festuca heterophylla* Lam.
 455. *Festuca pratensis* Huds. subsp. *pratensis*
 456. *Festuca rubra* L.
 457. *Festuca valesiaca* Schlecht. ex Gaudin
 458. *Glyceria fluitans* (L.) R.Br.
 459. *Holcus lanatus* L. – MM (2006)
 460. *Holcus mollis* L.
 461. *Hordelymus europaeus* (L.) Harz
 462. *Leersia oryzoides* (L.) Sw.
 463. *Lolium perenne* L. – HH herb. (1943)
 464. *Melica nutans* L.
 465. *Milium effusum* L.
 466. *Molinia caerulea* (L.) Moench
 467. *Nardus stricta* L.
 468. *Phleum pratense* L.
 469. *Poa annua* L.
 470. *Poa compressa* L.
 471. *Poa palustris* L.
 472. *Poa pratensis* L.
 473. *Poa trivialis* L.
 474. *Setaria pumila* (Poir.) Schult.
 475. *Trisetum flavescens* (L.) P.Beauv.

Lemnaceae

476. *Lemna minor* L.

¹ Höhr H. (1914) *Schäffburgs Archegoniaten (Moos- und Farnpflanzen)*.

² Data regarding the species from the herbarium of Henrich Höhr (noted as *HH herb. year* in the list of species) were provided by Klaus Gross, and were not checked by the author (KÖ). (?) stands for unknown year.

³ Fronius F. (1858) *Flora von Schäffburg*. Values in square brackets [] represent the position of the species in the list of Fronius.

⁴ Mititiuc M. (2006) *Rezervația naturală „Stejarul seculari de pe platoul Breite”*.

⁵ Oroian S. (2009) *Flora și vegetația satelor satești din Transilvania*.

Besides those included in the above list, there are a few plant species mentioned by Fronius (1858) as existing on the Breite. Despite the fact that their actual presence in the open area is questionable, they are listed in the following. Species names and authors are given as in *Flora Europaea* (2001), while values in square brackets [] represent the position of the species in the list of Fronius.

1. *Alchemilla vulgaris* L. [360]
2. *Astrantia major* L. [404]
3. *Carex flava* L. [1120]
4. *Cerastium brachycephalum* Desp. (accepted name *Cerastium brachypetalum* Pers.) [176]

5. *Cypripedium calceolus* L. (accepted name *Cypripedium calceolus* L.) [1015]
6. *Clematis flammula* L. [2]
7. *Cytisus capitatus* Scop. (syn., accepted name *Chamaecytisus supinus* (L.) Link.) [245]
8. *Erythronium dens-canis* L. (present on the Breite, but not seen by KÖ in the open area) [1042]
9. *Gentiana amarella* L. (probably *Gentianella amarella* (L.) Börner) [689]
10. *Gnaphalium dioicum* L. = *Antennaria dioica* (L.) Gaertn. [538]
11. *Helleborus purpurascens* Waldst. & Kit. (present on the Breite, but not seen by KÖ in the open area) [40]
12. *Hieracium praemorsum* L. (ext., accepted name *Crepis praemorsa* (L.) Walther.) [649]
13. *Iris caespitosa* Pall. ex Link (syn., accepted name *Iris ruthenica* Ker Gawl., and noted by Fronius as „=” *Iris graminea* L.) [1022]
14. *Linum flavum* L. [177]
15. *Melampyrum barbatum* Waldst. & Kit. ex Willd. [786]
16. *Orchideae maculata* L. (probably *Orchis maculata* L., syn., accepted name *Dactylorhiza maculata* (L.) Soó) [1000]
17. *Orobis niger* L. (syn., accepted name *Lathyrus niger* (L.) Bernh.) [310]
18. *Pimpinella magna* L. (syn., accepted name *Pimpinella major* (L.) Huds.) [414]
19. *Poa nemoralis* L. [1170]
20. *Scabiosa columbaria* L. [501]
21. *Srophularia glandulosa* Waldst. & Kit. (syn., *Scrophularia scopoli* Hoppe) [747]
22. *Vinca minor* L. (present on the Breite, but not seen by KÖ in the open area) [681]

Analysis of the flora of the Breite wood-pasture

Following the field studies begun in 2005, 476 species and subspecies belonging to 248 genera and 64 families were identified on the Breite. Additional species are to be expected based also on the unpublished list provided by Klaus Gross. The most abundant families (>5 %) are: Asteraceae (*Compositae*) – 14.08 %; Poaceae (*Gramineae*) – 10.92 %; Fabaceae (*Leguminosae*) – 7.14 %; Lamiaceae – 6.30 %; Cyperaceae – 5.25 % and Rosaceae 5.04 %, while 22 families are represented by only one species.

Considering their region of origin (geoelements), the identified plant species belong to the following groups: Eurasian (194 species: 40.76 %), European (74: 15.55 %), Circumpolar (52: 10.92 %) mainly Central European (49: 10.29 %; including for

example species that are predominantly Central European but also Mediterranean), Cosmopolite (37: 7.77 %), while the remaining 70 species (Mediterranean, Pontic, Balcanic *etc.*) total 14.71 %. Out of this last category, 17 are adventive species, requiring special attention from a biodiversity conservation perspective. Most of these species can be found nearby the forestry road that crosses the plateau in its length, and especially around the football field, while they are most abundant in the SE part of the plateau, where in 2007 a clear-cut was done in the surrounding forest (Fig. 1). According to the classification used by Anastasiu, Negrean (2009), these 17 species can be included in the following groups from the perspective of their abundance on the Breite:

- Common: *Erigeron annuus* (L.) Pers. subsp. *annuus*; *Erigeron annuus* (L.) Pers. subsp. *strigosus* (Muhl. ex Willd.) Wagenitz; *Galinsoga parviflora* Cav.; *Juncus tenuis* Willd.
- Scattered: *Amaranthus powellii* S. Watson; *Amaranthus retroflexus* L.; *Chamomilla suaveolens* (Pursh) Rydb.; *Erechtites hieracifolia* (L.) Rafin. ex DC.
- Rare: *Acer negundo* L.; *Asclepias syriaca* L.; *Conyza canadensis* (L.) Cronq.; *Oxalis fontana* Bunge; *Oxalis stricta* L.
- Locally abundant: *Rudbeckia laciniata* L.; *Solidago canadensis* L.; *Solidago gigantea* Aiton subsp. *serotina* (O. Kuntze) McNeil; *Veronica persica* Poir.

In order to limit their presence and future spread on the Breite, the individuals of the following species have been manually removed (pulled out, including roots) on different occasions: *Asclepias syriaca* L.; *Rudbeckia laciniata* L.; *Solidago canadensis* L.; *Solidago gigantea* Aiton subsp. *serotina* (O. Kuntze) McNeil.

The identified plant species were grouped according to their predominant life form (Fig. 2) (e.g. a hemicryptophyte, but secondary chamaephyte species “H(Ch)” was considered as hemicryptophyte). Hemicryptophytes are the most abundant group, being represented by 236 species (49.58 %), followed by therophytes with 103 species (21.64 %) The two epiphyte species are *Loranthus europaeus* Jack. and *Viscum album* L.

Considering their life span, perennials are the most abundant group, being represented by 289 species (60.71 %), followed by annuals (76: 15.97 %), trees (27: 5.67 %), biennials and shrubs (19: 3.99 % each), while the rest of species, with secondary life span (e.g. annual-perennials) total 9.66 %.

The ecological characterization of the flora according to the humidity requirements (Fig. 3) shows an overall mesophilous character and a significant presence of hygrophilous and hydrophilous elements, reflecting the historical (humid, even marshy - Siegmund 1948, Mihai Eminescu Trust 2005) characteristics of the plateau. The xerophilous species were found mainly in the more open areas of the wood-pasture (especially in the northern part), but also in some spots of the southern part, probably as a result of the desiccation process and unusual lack of precipitations during 2006-2009 in comparison with the multiannual averages (personal observations). The two hemiparasite species are *Loranthus europaeus* Jack. and *Viscum album* L.

When considering the biodiversity conservation value of the ground vegetation, it can be stated that it is represented by its general character (semi-natural habitat, wood-pasture in particular), and by the few species whose presence is interesting in this area (ex. the mountainous-alpine *Potentilla erecta* (L.) Rausch. and *Hieracium aurantiacum* L. subsp. *aurantiacum*) rather than in the presence of species of conservation interest.

The species included in the National Red Lists are: *Achillea ptarmica* L. (Vulnerable - Oltean *et al.*, 1994; Rare - Negrean 2001); *Dactylorhiza incarnata* (L.) Soó (R - Oltean *et al.* 1994, Negrean 2001); *Gentiana pneumonanthe* L. (V - Boşcaiu *et al.*, 1994; Negrean 2001); *Gymnadenia conopsea* (L.) R. Br. (R - Oltean *et al.* 1994); *Listera ovata* (L.) R.Br. (R - Oltean *et al.* 1994); *Orchis morio* L. (R - Oltean *et al.*, 1994; Negrean 2001); *Orchis ustulata* L. (R - Oltean *et al.* 1994, Negrean 2001); *Pinus sylvestris* L. (R - Oltean *et al.* 1994, planted on the Breite); *Platanthera bifolia* (L.) Rich. (R - Oltean *et al.* 1994); *Rosa micrantha* Borrer ex Sm. (R - Oltean *et al.* 1994); *Senecio aquaticus* Hill (R - Oltean *et al.*, 1994). Out of these, only *Dactylorhiza incarnata* (L.) Soó, *Gymnadenia conopsea* (L.) R. Br. and *Platanthera bifolia* (L.) Rich. are included in the European Red List of Vascular Plants (Bilz *et al.* 2011), all three as Least Concern. None of the species is listed in the Red Data Book of Vascular Plants for Romania (Dihoru, Negrean, 2009).

Discussion

The flora of the Breite wood-pasture contains a large number of species with very diverse ecological requirements, a situation explained probably by the special conditions created by the presence of the veteran trees (providing shadow

and maintaining humidity), the closeness of the surrounding forest, the hydrological regime, but also by the numerous interventions and modifications in the land-use of the area that occurred throughout the years. Therefore in the current species composition several species that characteristically do not belong to this territory have a transitory presence, while others are still present despite the fact that the ecological conditions are not favourable for their existence any more (Van der Maarel 1996). Beside the specific dynamics and distribution determined by these conditions, the reintroduction of grazing in 2008 as part of the management plan (Mihai Eminescu Trust 2007) will most probably influence the distribution and presence of some species.

For the overall management, interventions aiming to restore the former hydrological regime are recommended, together with controlling the spread of adventive species and shrub encroachment. In order to meet this, the enclosure of the drainage ditches started in 2007, together with removal of Hornbeam saplings and, as mentioned before, removal of adventive species (Mihai Eminescu Trust 2007).

Because, following the abandonment in 2004, grazing was reintroduced in 2008 as part of the conservation management activities, in the following some specific recommendations are presented regarding this matter from the perspective of wood-pastures, and therefore relevant for this type of habitat in general.

Grazing must be applied through rotation, not more than three-four days in the same spot, followed by 10-14 days of resting (Földes 1895, Bíró 1910).

Where scything is applied, although this is not a characteristic practice in wood-pasture habitats (Crofts, Jefferson 1999; Oroszi, 2004), grazing must be avoided especially during the dry season (Bíró, 1910). Sheep grazing should be generally avoided during the dry season because on torrid days animals pull out the plants instead of grazing, causing severe loss in biomass and regenerative parts of plants (Zsarolyáni 1897).

Weeds that appear in more intensely grazed areas (e.g. *Cirsium arvense* (L.) Scop. and *C. vulgare* (Savi) Ten., species that had an expansion in 2009) must be cut with a spade or hoe twice a year (in June and August) before flowering. Cattle droppings should be left to dry out and later spread across the area in order to avoid the concentration of nutrients (Bíró 1910). Because fallen leaves have a high mineral content that is very important

for herbaceous plants, cattle should be left to walk across the area during humid days in late autumn in order to trample the leaves, helping in this way the formation of a thick humus layer (Zsarolyáni 1897).

Although interventions might be considered by many as contradictory with nature conservation principles, it is of key importance to keep in mind that wood-pasture habitats are dependent on such practices, being the result of human activities, the seizure of these leading to their disappearance in the long run (Introduction and Bergmeier *et al.* 2010 for a recent review on this topic).

Acknowledgements

The author is grateful to Tibor Hartel, Gavril Negrean, Cosmin Ioan Moga and Anna-Mária Csergő for their help during fieldwork, to Vasile

Sanda and Gheorghe Dihoru for their help in the identification of several species and for Klaus Gross for providing his unpublished list regarding the flora of the Breite, including the herbarium data of Heinrich Höhr. Different stages of this study were funded by the Mihai Eminescu Trust, the Swedish Biodiversity Centre, the Deutsche Bunesstiftung Umwelt and Alfred Toepfer Stiftung through the NatuRegio Project, and the Institute of Biology Bucharest - Romanian Academy, through the project “Taxonomical and chorological researches on flora and fauna species from Romania”. Gavril Negrean and Constantin Drăgulescu provided valuable comments and observations during the preparation of the manuscript.

REFERENCES

- Akeroyd 2003 Akeroyd John, A *Transylvanian wood-pasture*. In: *Plant Talk* 34 (2003) United Kingdom, p. 34–37.
- Anastasiu, Negrean 2009 Anastasiu Paulina, Negrean Gavril, *Neophytes in Romania*. In: Rákossy László, Momeu Laura (Ed.) *Neobiota din România*. Presa Universitară Clujeană, Cluj Napoca (2009), p. 66–97.
- Baumgarten 1816 Baumgarten Johann Christian Gottlob, *Enumeratio stirpium Magno Transsilvaniae principatui praeprimis indigenarum in usum nostratum botanophilorum conscripta inque ordinem sexuali-naturalem concinnata*. In: *Vindobonae* (1, 2 in 1816, 3 in 1817), Libraria Camesinae), Sibiu (1816, 1817).
- Baumgarten 1846 Baumgarten Johann Christian Gottlob, *Enumeratio stirpium Magno Transsilvaniae principatui praeprimis indigenarum in usum nostratum botanophilorum conscripta inque ordinem sexuali-naturalem concinnata*. In: *Tomus Quartus*, Fuss Michael (Ed.) Typis Haeredum M. Nobilis de Hochmeister (Theodor Steinhausen), Sibiu (1846),
- Bergmeier *et al.* 2010 Bergmeier Erwin, Petermann Jörg, Schröder Eckhard, *Geobotanical survey of wood-pasture habitats in Europe: diversity, threats and conservation*. In: *Biodiversity and Conservation* 19, (2010), p. 2995–3014.
- Bilz *et al.* 2011 Bilz Melanie, Kell Shelagh P., Maxted Nigel, Lansdown Richard V., *European Red List of vascular Plants*. Publications Office of the European Union, Luxembourg (2011).
- Bíró 1910 Bíró János, *A legelő-erdők berendezéséről és gyepesítéséről, különös tekintettel a Székelyföldre*. In: *Erdészeti Lapok* 49-2, Budapest (1910), p. 38–69.
- Boşcaiu *et al.* 1994 Boşcaiu Nicolae, Coldea Gheorghe, Horeanu Clement, *Lista roşie a plantelor vasculare dispărute, periclitare, vulnerabile şi rare din Flora României*. In: *Ocotirea Naturii şi Mediului Înconjurător* 38, Bucureşti (1994), p. 45–56.
- Bucşa 2007 Bucşa Livia, *Macromycetes of the Breite nature reserve of ancient oaks (Transylvania, Romania)*. In: *Transylvanian Review of Systematical and Ecological Research* 4, Sibiu (2007), p. 33–42.
- Ciocârlan 2009 Ciocârlan Vasile, *Flora ilustrată a României*. Editura Ceres, Bucureşti (2009).
- Csergő 2007 Csergő Anna-Mária, *Vegetaţia ierboasă a platoului Breite*. In: Mihai Eminescu Trust (Ed.) *Planul de management al rezervaţiei “Stejarii seculari de la Breite”*, Sighişoara (2007), p. 14–19.
- Crofts, Jefferson 1999 Crofts A., Jefferson R.G. (Ed.), *Grassland management in wood-pasture and parkland*. In: *The lowland grassland management handbook*. 2nd edition. English Nature / The Wildlife Trusts section 14.3., (1999).
- Csűrös, Kovács 1962 Csűrös Ştefan, Kovács Attila, *Cercetări asupra vegetaţiei din raioanele Sighişoara şi Agnita*. In: *Contribuţii Botanice*, Cluj-Napoca (1962), p. 205–232.
- Dihoru, Negrean 2009 Dihoru Gheorghe, Negrean Gavril, *Cartea roşie a plantelor vasculare din România*. Editura Academiei Române, Bucureşti (2009).
- Flora Europaea 2001 Flora Europaea Database extracted from the digital version of the *Flora Europaea*, held at the Royal Botanic Garden Edinburgh (2001).
- Földes 1895 Földes János, *A legelő-erdők berendezése, okszerű kezelése, használata és felújítása*. In: *Erdészeti Lapok* 34, Budapest (1895), p. 30–72, 157–212, 278–329, 395–432, 526–536.
- Fronius 1858 Fronius Friedrich Franz, *Flora von Schäßburg. Ein Beitrag zur Flora von Siebenbürgen*. In: *Program des evangelischen Gymnasiums in Schässburg und der damit verbundenen Lehranstalten zum schluss des Schuljahres 1857/8*, Sighişoara (1858), p. 3–95.
- Fuss 1866 Fuss Michael, *Flora transsilvaniae Excursoria*. Munificentia Societatis pro illustranda Transsilvaniae cognitione et excellentissimi D. Archiepiscopi Dr. L. Haynald edidit societas naturae curiosorum Transsilvanica Cibiniensis. Typis Haeredum Georgii de Closius, Sibiu (1866).
- Hartel, Moga 2010 Hartel Tibor, Moga Cosmin Ioan, *Manual de bună practică în managementul*

- habitatelor de pajiște cu arbori seculari*. Fundația Mihai Eminescu Trust, Sighișoara (2010).
- Höhr 1914 Höhr Heinrich Wilhelm, *Schäßburgs Archegoniaten (Moos- und Farnpflanzen). Ein Beitrag zur Siebenbürgens Kryptogamenflora*. In: *Festschrift anlässlich der vom 30. August bis 2. September 1914 in Hermannstadt stattfindenden XXXVII. Wanderversammlung ungarischer Ärzte und Naturforscher*. Hrsgg. Von der Med. Sektion des Ver. Für Naturw. Hermannstadt. Michaelis i. Komm. Gr. 8, Sibiu (1914), p. 82–139.
- Höhr 1930 Höhr Heinrich Wilhelm, *Unsere Breite zur Herbstzeit*, In: *Groß-Kokler Bote*, Sighișoara (1930).
- Höhr 1949 Höhr Heinrich Wilhelm, *Die Pflanzengesellschaften des Großkokelgebietes bei Schäßburg und ihre Pflanzenarten. Ein vegetationskundliche Studie mit einer Vegetationskarte 1: 25.000* (1949). (unfinished manuscript).
- Kirby *et al.* 1995 Kirby K.J, Thomas R.C., Key R.S., McLean I.F.G., Hodgetts N. *Pasture woodland and its conservation in Britain*. In: *Biological Journal of the Linnean Society* 56, suppl., London (1995), p.135–153.
- Mihai Eminescu Trust 2005 Mihai Eminescu Trust (Ed.), *Documentația științifică pentru propunerea extinderii rezervației naturale „Stejarii seculari de la Breite”*, Sighișoara (2005).
- Mihai Eminescu Trust 2007 Mihai Eminescu Trust (Ed.), *Planul de management al rezervației “Stejarii seculari de la Breite”*, Sighișoara (2007).
- Mititiuc 2006 Mititiuc Mihai, *Rezervația naturală „Stejarii seculari de pe platoul Breite”*, In: *Catalog de semințe și spori oferite pentru schimb în anul 2006*. Grădina Botanică „Anastasia Fătu” a Universității „Alexandru Ioan Cuza” 83, Iași (2006) p. 53–54.
- Moga *et al.* 2007 Moga Cosmin Ioan, Öllerer Kinga, Hartel Tibor, *The breeding avifauna of the “Breite Plateau” Natural Reserve and the surrounding forest*. In: *Studia Universitatis Babeș-Bolyai, Seria Biologia* 52, Cluj-Napoca (2007), p. 33–43.
- Moga *et al.* 2009 Moga Cosmin Ioan, Hartel Tibor, Öllerer Kinga, *Ancient oak wood-pasture as a habitat for the endangered Tree Pipit, Anthus trivialis*. In: *Biologia* 64, (2009), p. 1011–1015.
- Muică 2009 Muică Adina Stela, *Fitodiversitatea platoului Breite-Sighișoara (jud. Mureș)*. “Lucian Blaga” University of Sibiu, Faculty of Sciences, Specialization Biology (2009). Licentiate thesis (manuscript).
- Negrean 2001 Negrean Gavril, *Lista roșie a plantelor din România existente în pajiști (inclusiv endemite și subendemite)*. In: Coldea Gheorghe, Negrean Gavril, Sârbu Ion, Sârbu Anca (Ed.) *Ghid pentru identificarea și inventarierea pajiștilor seminaturale din România*. Editura Alo, București (2001), p. 30–58.
- Oroian 2009 Oroian Silvia, *Flora și vegetația satelor sătești din Transilvania*. In *Editura University Press, Târgu-Mureș* (2009).
- Oroszi 2004 Oroszi Sándor, *Az erdélyi szászok erdőgazdálkodása*. In: *Erdészettörténeti közlemények* 63, Budapest (2004), p. 1–153.
- Oltean *et al.* 1994 Oltean Mircea, Negrean Gavril, Popescu Aurel, Roman Nicolae, Dihoru Gheorghe, Sanda Vasile, Mihăilescu Simona, *Lista roșie a plantelor superioare din România*. In: *Studii, sinteze, documentații de ecologie*. Academia Română, Institutul de Biologie, București (1994).
- Öllerer 2009 Öllerer Kinga, *Raport de cercetare privind ecologia covorului vegetal al Platoului Breite* (2009) (unpublished report for the Mihai Eminescu Trust).
- Öllerer 2010 Öllerer Kinga, *Ecologia covorului vegetal al Platoului Breite, importanța sa la nivel național și internațional și actualitatea măsurilor de conservare inițiate de Fundația Mihai Eminescu Trust* (2010) (unpublished report for the Mihai Eminescu Trust).
- Öllerer 2012 Öllerer Kinga, *Előzetes adatok a Breite legelőerdő flórájáról (Segesvár, Románia)*. In: *Kitaibelia - Journal of Botany and Nature Conservation* 17, Debrecen (2012), p. 131.
- Öllerer *et al.* 2008a Öllerer Kinga, Csörgő Anna-Mária, Hartel Tibor, *Vegetation studies towards the management of the Breite woodpasture (Sighisoara, Romania)*. In: *Kitaibelia - Journal of Botany and Nature Conservation* 13, Debrecen (2008), p. 184.

- Öllerer *et al.* 2008b Öllerer Kinga, Hartel Tibor, Moga Cosmin Ioan, Csergő Anna-Mária, *Conservation management activities in the Breite wood pasture natural reserve (Sighisoara, Romania)*. In: 6th European Conference on Ecological Restoration, 8-12 September, Ghent, Belgium (2008). <http://ser.semico.be/ser-pdf/092.pdf>
- Öllerer *et al.* 2010 Öllerer Kinga, Moga Cosmin Ioan, Samoilă Ciprian, *The wood-pastures from Saxon Transylvania, Romania: knowledge, conservation status and trends*. In: *II International Conference "The Oak - Ecology, History, Management and Planning"*, Isparta, Turkey. *Book of abstracts*, (2010), p. 81–82.
- Rich *et al.* 2005 Rich T., Peterken G., Tucker G., McMeechan F., Dobson D. *Habitat requirements and issues. Lowland wood-pastures and parkland*. In: Hill David, Fasham Matthew, Tucker Graham, Shewry Michael, Shaw Philip. (Ed.) *Handbook of Biodiversity Methods. Survey, Evaluation and Monitoring*. Cambridge University Press, (2005), p. 114–117.
- Rois-Díaz *et al.* 2006 Rois-Díaz Mercedes, Mosquera-Losada Rosa, Rigueiro-Rodríguez Antonio, *Biodiversity Indicators on Silvopastoralism across Europe*. In: *Technical Report 21*. European Forest Institute, (2006).
- Sanda *et al.* 1976 Sanda Vasile, Popescu Aurel, Doltu Marcel Ion, *Contribuții la cunoașterea vegetației din bazinul mijlociu al Tîrnavei Mari*. In: *Studii și Comunicări, Științele Naturii, Muzeul Brukenthal* 20, Sibiu (1976), p. 65–93.
- Săvulescu 1952-1976 Săvulescu Traian. (Ed.) *Flora României. Flora Romaniae*. Editura Academiei Române, vol. I–XIII, București (1952–1976).
- Schur 1866 Schur Philipp Johann Ferdinand, *Enumeratio plantarum Transsilvaniae exhibens: stirpes phanerogamas sponte crescentes atque frequentius cultas, cryptogamas vasculares, characeas, etiam muscos hepaticasque*. In: *Vindobonae*, G. Braumüller, (1866).
- Siegmund 1948 Forstrat Konrad Siegmund, *Der Stadtwald von Schäßburg, seine Lebensgeschichte*. Das Archiv der Siebenbürgischen Bibliothek in Gundelsheim. Tiposcript (1948).
- Simonkai 1886 Simonkai Lajos, *Erdély edényes flórájának helyesbitett foglalata (Enumeratio Florae Transilvanicae vesiculosae critica*. In: *Ex mandato Societas Scientiarum Naturalium Regiae Hungaricae*). Kir. Magyar Természettudományi Társulat, Budapest (1886-1887).
- Stiven 2009 Stiven Roland, *Management of ancient wood pasture. Guidance note*. Forestry Commission Scotland, Edinburgh (2009).
- Teșculă, Goța 2007 Teșculă Nicolae, Goța Alexandru, *Istoria ariei protejate*. In: Mihai Eminescu Trust (Ed.) *Planul de management al rezervației "Stejarii seculari de la Breite"*, Sighișoara (2007), p. 51–55.
- Uhde 2011 Uhde Britta, *How to protect the wood pasture "Breite" (Sighișoara, Romania)? - Analysis on the oak regeneration at the forest edge*. Fakultät Forst-, Geo- und Hydrowissenschaften der Technischen Universität Dresden, Dresden (2011). Licentiate thesis (manuscript).
- Van der Maarel Van der Maarel Eddy, *Pattern and process in the plant community: Fifty years after A.S. Watt*. In: *Journal of Vegetation Science* 7, (1996), p. 19–28.
- Zsarolyáni 1897 Zsarolyáni Márton Sándor, *A legelő-erdők berendezésénél figyelembe veendő tényezőkről*. In: *Erdészeti Lapok* 36-8, Budapest (1897), p. 619–647.

LIST OF ILLUSTRATIONS

- Fig. 1.** Localization and overview of the Breite wood-pasture based on the Google Earth imagery
- Fig. 2.** The life form categories of vascular plants from the Breite wood-pasture:
Ch – chamaephyte;
Ep – epiphyte;
G – geophyte;
H – hemicryptophyte;
Hh – helohidatophyte (hydrophyte);
Ht – hemitherophyte;
Ph – phanerophyte;
T – therophyte.
- Fig. 3.** The ecological categories of vascular plants from the Breite wood-pasture:
higr, hydr – hygrophilous, hydrophilous;
eurif – euriphyte;
mes – mesophilous;
mesohigr – mesohygrophilous;
mesoxer – mesoxerophilous;
semipar – semiparasite;
subsp, cultiv, rud – subsponaneous, cultivated, ruderal;
xer – xerophilous;
xeromes – xeromesophilous.

LISTA ILUSTRAȚIILOR

- Fig. 1.** Localizarea și aspectul general al platoului Breite pe baza imaginii Google Earth.
- Fig. 2.** Spectrul bioformelor pe platoul Breite:
Ch – camefite;
Ep – epifite;
G – geofite;
H – hemicriptofite;
Hh – helohidatofite (hidrofite);
Ht – hemiterofite;
Ph – fanerofite;
T – terofite.
- Fig. 3.** Caracterizarea ecologică a florei platoului Breite:
higr, hydr – higrofite, hidrofite;
eurif – eurifite;
mes – mezofite;
mesohigr – mezohigrofite;
mesoxer – mezoxerofite;
semipar – semiparazite;
subsp, cultiv, rud – subsponane, cultivate, ruderale;
xer – xerofite;
xeromes – xeromezofite.



Fig. 1. Localization and overview of the Breite wood-pasture based on the Google Earth imagery




Fig. 2. The life form categories of vascular plants from the Breite wood-pasture




Fig. 3. The ecological categories of vascular plants from the Breite wood-pasture

GEOLOGICAL AND HYDROLOGICAL CONSIDERATIONS IN THE SECAȘUL MARE BASIN (SOUTH – WESTERN PART OF THE TRANSYLVANIAN DEPRESSION)

Marioara COSTEA*
Rodica CIOBANU**

Abstract. *In the current paper we intend to present the main geological factors leading to or restricting the evolution of the underground waters in the Secașul Mare Basin (the structural and lithological characteristics of the Secașul Mare basin), hydrological aspects of the store rocks and the underground waters resources in the given basin. Since the underground waters represent an important natural resource, their quality, mineralization degree and chemical content are all very relevant. Any form of natural water (rain fall waters, surface waters or subterranean waters) contains chemical substances, some of which may be harmful, either to man or to other living organisms. The modification of the water chemistry can be due to the mineral or rock dissolution phenomenon, as a result of the water – rock contact, as well as to the anthropic pollution. We also point out the necessity of permanent monitoring of the quality of underground waters, and especially of those used in agriculture, fish – farming, house management and consumption.*

Keywords: *Geology, Hydrology, Secaș Basin, Transylvanian Depression.*

Rezumat. *În lucrare ne propunem să prezentăm factorii geologici majori care concură sau restricționează evoluția apelor subterane în bazinul Secașului Mare (caracteristicile structurale și litologice ale bazinului Secașului Mare), aspecte hidrogeologice ale rocilor magazin și resursele de ape subterane din bazinul Secașului Mare. În condițiile în care apele subterane reprezintă o resursă naturală importantă, calitatea acestora - gradul de mineralizare, conținutul chimic este foarte important. Orice apă naturală - apa din precipitații, apele de suprafață sau cele subterane conțin substanțe chimice dintre care unele sunt dăunătoare omului sau altor forme de viață. Modificarea chimismului apelor se poate datora atât fenomenului de dizolvare a mineralelor, rocilor la contactul rocă-apă, cât și poluării antropice. Procesele chimice la contactul apă-rocă sunt necontrolabile, complicate și de multe ori necunoscute. Evidențiem în lucrare necesitatea unei monitorizări permanente a calității apelor subterane, și în special a celor utilizate în agricultură, piscicultură, gospodării și în consum.*

Cuvinte cheie: *geologie, hidrologie, Bazinul Secașului, Depresiunea Transilvaniei.*

Introduction

Regarded an important geospheres of the Earth, the lithosphere and the hydrosphere represent at the same time the environment and the compounding elements ensuring and realizing the material, informational and energetic network in the geosphere. The study of the organization and manifestation forms in the continental field, commonly points out the complex and at times – common genesis of the geological and hydrological processes and phenomena, and not only. In this context, the

organization of the hydrological network, the current hydrological phenomena and manifestation forms, all require a cross – disciplinary approach and a correlative time – space analysis, particularly with regards to the geological conditions, as well as to other environmental elements (relief, weather condition, vegetation coverage degree, fields' utilization).

There are strong and at times complicated relations between water, in all its forms, and the rocks and geological structures, respectively tectonic structures of the lithosphere. The evolution of the geological and geomorphological processes from within and from the surface of the Earth crust is strongly connected to the presence of water. Water in its liquid state, is highly significant, on one hand, as a morphogenetic factor, as a factor for setting a hierarchy of the landforms, and on the other hand

* Lucian Blaga University of Sibiu, Science Faculty,
marioara_costea@yahoo.com

** Brukenthal National Museum,
rodi_ciobanu2005@yahoo.com,
rodica.ciobanu@brukenthalmuseum.ro

due to the economic significance it has based on its quality.

The presence and evolution of the underground water resources is influenced by the geological factors, both from a quantitative and qualitative point of view (due to the physical and chemical properties), as well as from the point of view of its accessibility and distribution. The rocks, the geological and tectonic structures which characterize them, are all in a continuous inter-conditional relation with water. The outcomes of this biunivocal relation take a concrete form both in what concerns the evolution of the hydro – geological resources, as well as that of the ancient and especially recent local or regional geology. The evolution and the spread of underground water resources are conditioned by the circulation of water within the rock, among rocks, on the stratification levels, reaching from the surface towards the inner part, through infiltration and in reverse, through capillarity. Equally important are the physical properties of the store rocks, the behaviour of the rock and the physical and chemical processes, taking place in the presence of water – all of which represent conditioning factors for the genetic characteristics passed on subsequently to the underground water.

In the current paper our goal is to present the major geological factors which contribute to or restrict the evolution of underground water in the Secaşul Mare basin.

The structural and lithological characteristics of the Secaşul Mare basin

In order to properly and correctly perform a geological analysis and to integrate the forming conditions of the studied region, in the tectonic and lithological background of southern Transylvania, our analysis is done in correlation to the regional context and is related to an area greater than the one within the limits of the hydrographic basin of Secaşul Mare.

The hydrographic basin of Secaşul Mare represents a geographic space of geological and geomorphological interference, in which the elements of the landform interact and are interdependent with the anthropic element, conditioning each other. It is from this interference that a complex of relations emerges, which individualizes and personalizes the hydrographic basin among the landform units partially entering its limits: the Cindrel Mountains in the south, the Sebeş – Apold depressionary passage in the center

and the Secaşelor Plateau in the northern part of the basin (Costea 2006).

The genesis and evolution of the Secaşul Mare Basin have permanently been conditioned by its geographical overlapping an interference zone of two areas with diverse genetic and evolutionary tendencies: subsiding and accumulation in the Transylvanian Basin, while on the other hand – rising and erosion in the Cindrel Mountains. From a geologic point of view, the Secaşelor Plateau, the Apoldu Depression and the contact Depression Sebeş – Alba Iulia represent sub – units of the Transylvanian Depression, situated at its contact with the mountain ridge of the Southern Carpathians (Carpații Meridionali) and of the Western Mountains (Munții Apuseni). The former came to be as a result of the following two phases: the basin phase – in which the Transylvanian Depression was formed and sedimented, and the sub aerial modelling phase – in which the landform of the Apoldu Passage and of the limitrophe areas is defined, evolving into the current general line (Sandu 1998, Ciobanu 2002).

The foundation of the region in the south – west of the Transylvanian Depression, a region upon which the Secaşul Mare Basin overlaps, is crystalline – Mesozoic, made up of: green sericite – calcarous schists, with quartz lenses, quartzites, paragneiss with muscovite and biotite and micaceous quartzite schists with granites. The pre – Laramie sediments¹ were deposited over the crystalline itself, which is covering the fundament and which date from Mesozoic times, belonging to two cycles: a Permian – Triassic one and a Lower Jurassic – Cretaceous one. The oldest sedimentary deposits belong to the Lower Permian² and can be encountered only in the eastern part of the Transylvanian Depression, while in the south – western part these such deposits are lacking.

The Triassic deposits are located along the regional Alba Iulia – Șeica Mare fault and are represented by: dolomite, calcareous marl, limestone and reddish conglomerate. This facies is also present on the southern side, at the contact point with the Sebeş Mountains, therefore appearing similar to the Triassic series of the “Codru” Domain (Ilie 1958). Following the deposit of the Transylvanian

¹The pre-laramie sedimentary and the crystalline schists form the crystalline – Mesozoic foundation.

²Based on a paleontological approach, only the presence of the Aptian and of the Senonian were identified, but the geognostic approach reveals that the pre-laramie sediment includes also Permian deposits: breccia and conglomerates with crystalline schists elements.

Depression in Triassic, an exondation took place, lasting until the Lower Jurassic. In the Middle and Superior Jurassic, massive calcareous conglomerate were then deposited, along with greyish blackish or pink limestone and coralline limestone. The Lower Cretaceous is represented by Barremian – Aptian deposits (grey compact marl, blackish marl, reddish brown clay with gritstone intercalations, *Orbitolina lenticularis* limestone) and albian deposits (greyish – yellowish compact limestone, grit marl, greyish hard gritstone, greyish – greenish conglomerates, compact tectonic marl).

The proper sedimentary deposits of the Transylvanian Depression and of the south – western part of the depression belong to the sedimentation cycles, starting from Upper Cretaceous and until Quaternary, with certain stratigraphic lacunas due to the exondation suffered by the Transylvanian Block (Fig.1).

The Upper Cretaceous is present in the south – western part of the Transylvanian Depression in Senonian, Cenomanian and Vraconnian deposits. To the south of Sebeş-Alba, the Senonian lies transgressively over the crystalline: at the bottom of the deposits there is clay and greyish micaceous gritstone, with sand and light cemented grovel intercalations and small conglomerates, followed by rough micaceous sand, gritstone, conglomerates and greyish – greenish marls. The complex also contains coal lenses and sulphur efflorescence. The series is wrapped up by massive polygenic conglomerates, with rolled up quartzite elements, micaschists, gneiss and seldom crystalline blocks, whose diameter does not reach higher than 0.5 m. In the Cacova-Săsciori area, on the touch point line of the Sebeş-Alba Iulia Depression with the Şureanu and Cindrel Mountains, the cretaceous deposits reach the surface (Ciupagea *et al.* 1970).

The Cenomanian and the Vraconnian are present on the southern part of the Secaşul Mare basin, on the Poiana Sibiului – Rod – Apoldu de Sus alignment through: rough calcareous – micaceous gritstone with thin intercalations of greyish marls with foraminiferae, marly - limestone, light cemented gritstone. Towards the north of the Secaş hydrographical basin, the upper Cretaceous is to be found in flysch facies and the Cretaceous deposits outcrop, in the Gosau facies, on a 1,300 m wide area, similar to the one from Săsciori.

The Paleogene rises transgressively over the Upper Cretaceous and the crystalline. The Transylvanian Basin started to evolve as an inter – mountain accumulation basin towards the end of the

Cretaceous, after the finalization of the Laramie climax. The basin was not unitary; emerging areas carried on evolving (towards the centre and eastern part of the depression, the Miocene deposits lay directly on top of the crystalline), and areas where sunken Paleogene deposits are present (the early accumulation basins). The deposits are marked by great facies and thickness variations, due to the Laramie and post – Laramie movements, which have influenced the sedimentation. The Paleogene is represented by Eocene and Oligocene deposits.

The Eocene appears transgressively in the Sebeş-Alba Iulia area over the Upper Cretaceous and is made up of gritstone, conglomerates, marl and clay with thin intercalations of organogenetic limestone with nummulites. The Eocene reaches 100 – 870 m in thickness in the Secaşelor Plateau and is represented by deposits of: reddish clay at the bottom, followed towards the upper level by grit limestone with nummulites, of the Porceşti type.

The Oligocene appears in the Alba Iulia area in: siliceous sands, fossiliferous gritstone and bituminous limestone. During the Eocene and Oligocene sedimentation, the Transylvanian Depression was continuously connected to the northern part of the Pannonian and of the Maramureş Basin. The Oligocene sedimentation phase was followed by an exondation and erosion phase, thus resulting in the absence of the Oligocene deposits from the higher areas of the Secaşelor Plateau.

Towards the southern and south – western part, the connection with the Pannonian Basin was interrupted ever since the Oligocene, when the exondation was stopped by strong erosion, thus making it possible for the Cretaceous deposits in the Sebeşului area to reach the surface.

The Lower Miocene is represented by deposits from the Aquitanian – Burdigalian stage. The Aquitanian deposits from the Sebeş-Alba Iulia are transgressively lying over the Cretaceous deposits in the eastern site of the Apuseni Mountains, or in patches over the Eocene deposits. The Burdigalian is present in gritstone with *Operculina* (Ciupagea *et al.* 1970) in the Alba Iulia – Sebeş region, on Secaşul Mare Valley and on its tributary streams on the right and transgressive gritstone over the red Aquitanian complex. The Helvetian emerges especially in the slopes of the Secaşelor Plateau, through a clayey marls complex with gritstone intercalations, sand and seldom, with gravel. Towards the north of Sebeş, Râpa Roşie, is an opening in the slope, highlighting an alternation of

grovel with rolled up quartzite of different colours, micaschists, diabase, porphyrite, jasper with red and white quartzite sand, brick – coloured and greenish clay (Fig. 2). By cementation, they developed a conglomerate appearance. The age of these deposits has been intensely discussed: Paucă (1971) attributes them to the Tortonian, while Mutihac (1990) is of the opinion that they belong to the Aquitanian.

The *Upper Miocene* deposits (*Badenian* – *Buglovian* - *Sarmatian*) are characterized by an extremely rich and varied fauna, similarly to the Sub Carpathian deposits of the same age; from the point of view of the succession, namely: tuff horizon at the bottom, salt horizon, schists horizon with radiolarians, marl upper horizon with *Spiralis*.

Towards the south – west of the Transylvanian Depression the dacite tuff horizon is replaced by the tuffaceous marls complex with *Globigerine*, with dacite tuff intercalations and calcareous marls. In the Alba Iulia – Sebeş zone, a conglomerate horizon of about 5 m in thickness was located under the horizon of this tuffaceous marls, and which was made up of calks elements with nummulites (the Porceşti calk) and of crystalline rocks (Ciupagea *et al.* 1970).

The salt horizon is very well developed towards the north and east of the hydrographical basin of the Secaşul Mare (Ocna Mureş, Ocna Sibiului) with variable thickness, which would later on become unnoticeable in the Miercurea Sibiului – Daia Română area, with no exploitation possibilities. Its presence is pointed out at the surface by the “salty water” springs (the springs from Miercurea Băi and the Slatina spring to the north of Daia Română). On top of the salt horizon there lie the Badenian deposits, which in the Sebeş – Apoldu de Sus area are transgressive and take the form of patches over the Upper Cretaceous ones, and even over the crystalline schists in the Cindrel Mountains. The Badenian patches are represented by tuffs and tuffaceous marls, blackish marls with gypsum efflorescence (Fig. 4). One notices the ever higher development of the Badenian in lagoon facies. There are lagoon deposits present, made up of gypsum, salty marls, salty springs (Ilie 1955), as well as in the constant debit and high salt level in the fountains.

To be noticed also the Badenian in reef coast facies, under the form of isolated limestone blocks, from Apold and Rod, as well as the shallow neritic facies (white greyish marls with *Globigerine*), accompanying the Aquitanian from Alba Iulia and the crystalline on the mountain edges. Likewise, in

the Bişbocuş steep from Daia Română, there are brownish clays with sulfur and gypsum efflorescence and with yellowish – whitish sand intercalations, while at Cut there are greyish blackish marls.

This have a local spread in the Secaşul Mare basin, especially on the Slatina Valley (toponym indicating the presence of the salty horizons) and at Miercurea Sibiului, where the salt manifestations appear in the axle of several anticlines. Furthermore, there are salty efflorescence also at Apoldu de Jos, in the openings on the southern slope of the Potca hill (Fig.4), where there is also vegetation pointing towards this fact, composed of halophytic plants: *Salsola*, *Salicornia* and *Tamarix*. These efflorescences can be the result of a certain gradual thinning of the salt lenses from the Ocna Sibiului mountain or they can also be due to an intense capillary circulation of the water from the substratum, which ends up dissolving the salt from the salty lenses and then, as a result of intense evaporation, to be deposited under the form of an efflorescence on the stratified surfaces³.

The Sarmatian emerges in the Sebeş – Apoldu de Sus area in Volhynian and Bessarabian deposits of marls and clays, with grit stone and sand intercalations. To the south of Miercurea Sibiului, as well as in the Dobârca – Câlnic area, the Sarmatian appears transgressively over the crystalline (Fig.1, 3), over the Senonian or over the Badenian, within conglomerate sandy facies, with marl intercalations. The Sarmatian sediments display various concretions, cuasi-spherical or ovoidal connected by limy cement, within the slightly cemented sand horizon (Vancea 1960). These deposits were brought to the light at Răhău, Reciu, Gârbova, Dobârca and Apoldu de Sus, and are covered by Pannonian deposits Apold. We are referring here to the fossil site on the Câlnicului Valley to the east of the fortress (grey and greyish – yellow sand, grey micaceous marl) and on the Dobârca Valley, to the north of the settlement (yellowish – grey sand), with a matching fauna, composed of *Ervilia*, *Cardium*, *Piranella* etc.

³The concretions are present in all the areas where there are saliferous horizons. One believes that the tectonic movements of minor scale, due to the movement of salt towards the surface, determine the modification of the passing direction of carbonated water through sand, thus resulting, by means of cementation, certain unusual forms – the trovants.

The *Pliocene*⁴ is present in the area in the fossil sites of the Pontian (in the Miercurea Sibiului area, at Cunța and Șpring), which is highlighted in the Secaşelor Plateau, in the Arini ravine from Daia (micaceous sand, sand with clay intercalations, greyish or yellowish clay deposited in a torrential structure), as well as at the bottom of the Ghirbomului Hill, also at Daia, in an opening caused by a torrential organism (grey blackish clay), in which a *Melanopsis* and *Congeria* fauna was identified. Pontian deposits were also identified at Cunța (gravels with quartzite elements and greyish - violet marls with organic traces of *Limnocardium*, *Congeria* and *Melanopsis*), at Șpring (thoroughly rolled quartzite grovels with clay, sand and marls containing traces of *Congeria*), at Miercurea Sibiului on Morii Hill (grovels caught in a mass of ferruginous sands with traces of *Limnocardium*, *Congeria* and *Valenciennius*).

At the bottom of the Apold Depression, the Pontian deposits are further crumbled, so that at Apold and Sângătin one comes across fine grovel and white micaceous and ferruginous sand, displayed in a torrential structure. There is small gravel with sandy cement, with clay and grey marl enclaves, with ferruginous alterations, at Apoldu

⁴The latest research in the field reveal that only the Dacian and the Romanian belong to the Pliocene (Mutihac 1990) while the Meotian and the Pontian belong to superior Miocene, these levels being transgressively and not harmoniously located over the Sarmatian. In the southern part of the Transylvanian Basin, within the studied area, the Meotian was not identified, and Huică and his collab. (1972) pointed out that the lower part of the Pannonian can be labelled with the superior Bessarabian, the Kersonian and the Meotian from the Euxino-Caspic basin, while the superior part can be labelled with the extra - carpathian Pontian.

Lubenescu (1977) pointed out that the Pannonian occupies the south - western part of the Transylvanian Basin, lying transgressively and unharmoniously on the sarmatian and badenian deposits, or even on the crystalline. This particular distribution can be noticed on the geological map (1: 200 000, ed. 1978), in the presence of the Pannonian in the western part of the Secaşul Mare basin, both to the north, as well as to the south of the hydrographic artery, being represented by thick packs of yellowish sands, alternating thinner clay and marl horizons, visible in the openings from Miercurea Sibiului, the Gorganu Drașov hill, at Sângătin, Ludoș and Doștat. Posea (2002) finds the Pannonian to be equivalent with the Pontian, while Mutihac (1990) notices that there is no Pannonian in the Transylvanian Depression.

de Sus, on the Rod spring, transgressively over the sandy Sarmatian. The marl deposits are covered with a complex layer of gravel, with crystalline schists and quartzite elements, alternating with sand (Ilie 1955, Ciupagea *et al.* 1970).

The *Quaternary deposits* are of significant thickness in the touch point area, cross by Secaş and by its left side tributaries.⁵ These are represented by sand and slightly rolled grovel, located in the glacis, terraces and river beds of the Sebeș and Secaşul Mare rivers, thus forming permeable strata with spring lines at the bottom of glacis and of the front parts of the terraces.

The morpho - tectonics of this sector completes the lithological and structural picture. Thus, the following anticline structures, on the north - east and south - west general orientation and with small variations from this direction, were identified by Dumitru (1992), quoted by Raboca (1995): the Daia Română anticline (Sarmatian and Pliocene deposits), the Slatina anticline, following the course of the Slatina Valley (Sarmatian deposits with salt occurrences), the Sebeș anticline (lower Miocene, Badenian and Sarmatian) and the Miercurea Băi - Doștat anticline (Pliocene deposits with salt occurrences). The orientation of the anticline structures generally follows the morphology of the landform, with their aksum being overlapped on the valley aisles, except for the Doștat - Sângătin - Miercurea anticline, which is not adapted to the morphology of the landform. These structures include synclines and are both part of the virgation folding system, located between Mureșului Valley, Cindrel Mountains and the Ohaba - Ocna Sibiului anticline (Ilie 1955, Ciupagea *et al.* 1970, Raboca 1995).

To sum up, the lithological formation and palaeo - geographic evolution of the Sebeș basin stand proof to a series of tight connections between the depressionary unit and the Carpathian mountain unit neighbouring the basin and occurring from Paleogene and until present day.

Hydrogeological aspects of the reservoir rocks

The existence and evolution of the water resources is connected to the water behaviour within the rock, to the processes taking place in the presence of water and depending on their formation or later on

⁵ Downstream, these rivers developed an orientation contrary to the Mureș mainstream. This orientation is due to a small block within the fundament having sunk in the area corresponding marginal threshold to the north - western part of Sebeș.

gained characteristics. Within the rocks, water is under the action of two universal forces – the *gravitational* force, under whose action water moves downwards, and the *molecular* attraction taking place between the molecules of the mineral (as component of a rock). The rock – water complex is under the influence of these forces, which impact both the flow of the water through the rocks, as well as their physical – mechanical characteristics.

According to the nature of the forces taking place between water and the mineral, the water can be *connected* or *free*. The quantitative existence and evolution of the chemically and physically *connected water* therefore depends on the mineralogical composition of the rock, on the environment conditions (endogenous – temperature, pressure and exogenous – the infiltration water). Within the analyzed area, We can notice that in the category of the physically connected water, the hygroscopic water has a decisive importance in inducing terrain degradation due to gravitational processes, such as the sliding. Thus, the quantity of hygroscopic water retained at the surface of the particles is of 10% for clay and of only 2% for sand. The quantity of connected water from the surface of the particles of clay depends on: the mineralogical composition, the cementation degree of the rock (the dispersion of the mineral granules), the presence of soluble salts in the water. Given the increase of humidity, the areas with clay deposits are considered to be “sensitive deposits”, since the hygroscopic water covers the particles completely, thus slightly crossing into a plastic state (Ciocârdel 1957).

The unconnected (free) water is the water which is maintained in deposits, respectively in the rocks due to the gravitational and capillary forces. The movement of the gravitational water is done through the sub- capillary pores, cracks, fractures, while the capillarity is conditioned by the reduced dimensions of the pores and of the solid particles⁶. Therefore, if the presence of the gravitational water and of the hydrodynamic processes connected to it is conditioned by the tectonic and geological structures, as far as the capillary water is concerned, the primary geological structures connected to the genesis of the rocks play a determinant role.

The hydrogeological characteristics of the aquifer strata depend, first of all, on the structure and texture of the reservoir rocks. Thus, in the Secașul Mare basin, there are two types of rocks which can store water: rocks with pores/ interstitial and cracked rocks. The interstitial rocks are the sedimentary detritus and light cemented rocks, with granule of very varied forms and dimensions, closing within small gaps in the shape of pores, which can be occupied by water and gas, especially in the plateau side. The cracked rocks are the compact rocks developed in the southern part of the basin, hard and cracked, in which the most important gaps are connected to the presence of the open cracks.

Regarding the permeability, the rocks from the geologic substratum of the basin are both *permeable* (the porous rocks – sand, gravel, marl and those which are genetically impermeable but once cracked, become permeable) and *impermeable* (clay, clayey schists, marl, un – cracked metamorphic rocks, compact limestone, residual deposits, etc.) (Băncilă *et al.* 1980). The water circulation is enabled by the gravity force, by the capillarity, but also by the horizontal transmissibility. Water circulates through the cracks, by the separation sides of the strata to the plateau monocline structure (stratification level), on the separation levels between the crystalline fundament and the sedimentary cover (the Călnic basin, Garbova, where the crystalline is very close to the surface or even outcrop at the surface). As for the deposits with fine granulation, the circulation of the water is done through the pores, which can communicate to each other, being in such close proximity.

The circulation of underground water within the rock is done first and foremost through the pores. The volume of the pores related to the volume of the rock is expressed by the quantifiable physical measure of the *porosity* (Zamfirescu 1997). The value of the porosity varies a lot for the different rock categories, as well as within the same category (Tab. 1).

The development of the underground water sources is directly dependent upon the character of the infiltration. In Secașului Mare Basin, the geological factors – the structure and the lithology – are among the most important factors determining the character of the water infiltration in the alteration cover. The most permeable deposits are the quaternary ones, made up of the superficial sediments of macro – porous or cracked sand and gravel, which have a high capacity to

⁶ By experimental means, we discovered the following capillar heights for different types of rocks: sand 0.1-0.5 m; loess 2-50 m; lutite rock 5-15 m; clay 20 and even over 50 m.

retain water and ensure the optimal conditions for infiltration. The Pliocene clay, the clay deposits, Sarmatian and Badenian marl deposits are limiting the infiltration process, therefore decreasing the contribution of surface water. The alternation of these deposits with sand, conglomerates, fine gravel lenses, etc. and the stratification both bring and extra permeability, as a result either of the infiltration in the stratification levels, or of the lack of rock homogeneity.

As a result, the macro – porous or cracked rocks foster the development of underground water resources, due to the increase of the infiltration water, such as the cracked limestone from the south – western part of the basin, the detritus un – cemented rocks, like the sand, gravel or cemented – such as the marl, the conglomerates, etc. which are to be found particularly in the Apoldu Depression and the piedmont hills. On the other hand, the water repellent rocks, the water repellent rocks are compact and without cracks, preventing the infiltration process: the clay, predominant in the plateau sector, the compact metamorphic rocks, in the shape of blocks, from the southern extremity of the basin, etc. (Fig.1).

Underground water resources in the Secaşul Mare basin

The distribution and characteristics of the underground water reserves are subject to the current relations between the alimentation through rainfall, infiltration, surface and underground leakage, the petrographic nature and the structure (*the geological location* - Romanescu 2003), differentiated by the two sectors of the basin: the sub - mountain sector and the plateau sector (Fig. 5).

In the southern part of the Secaşul Mare basin, which is taking the lower level of the Cindrel Mountains and the sub - mountain hills at their bottom, the underground water appears in the form an aquifer complex, whose formation is conditioned by the type of rock. In the Rod basin, the **aquifer complex of the metamorphic rocks** (crystalline schists, mica schists, micaceous gneiss, etc.) has the highest percentage (Trufaş 1978; Trufaş, Ştef 1998). These rocks are part of the Getic Nape and are practically impermeable, however, the complex system of faults, cracks and schist planes, some of which are open and unclogged, of clayey material, allowing the infiltration and the underground gravitational circulation of snow and rainfall.

The epimorphic complexes situated on the northern part of the Cindrel Mountains (at the touch point with the Apoldu Depression) have, generally, the same characteristics, except for certain areas reduced in surface, situated in the south – western part of the Secaşul Mare basin, which is dominated by limestone and gritstones. Thus, in the Săsciori – Piatra Varului - Sebeşel - Petreşti sector, the **calcareous - gritstone aquifer complex**, which was formed due to infiltration on the diacase lines and on the stratification planes of water coming from rainfall and from snow melting, according to a permeability coefficient of 0,2 – 0,5 m/day (I.S.P.H. 1998). The hydrogeological importance of these deposits is limited, and the accumulation possibility for large volumes of underground water is low. The hydrologic regime of the springs is conditioned by the rainfall regime, most of the appearances being situated at the touchpoint of the geologic elements and having a low debit: 0,1 – 0,2 l/s. the diluvia, the coluvia and the proluvia formed as a result to the rock alteration have a high permeability coefficient (20–25 m/day) and allow the quartering and swift circulation of the sheets and lenses of phreatic water, which is supplying the springs.

In the Apoldu Depression, the phreatic water appears in the form of continuous sheet in the **river meadow** of Secaşul Mare and its affluents, the **alluvial deposits** and discontinuously in the **terrace deposits**. In the river meadows, the level of the phreatic water is close to the surface (0,50 m), thus resulting under light drainage into swamp areas. The direct or indirect tributaries of Secaşul Mare fragment the glacis and the terraces, thus revealing, by means of erosion the ground water from the aquifer horizons. The thickness of the phreatic strata differs according to the local conditions: the inclination angle of the strata, the report to the hydrographic surface arteries, the volume of water carried by the rivers, the length of the course upstream, the climatic conditions, etc. the level oscillations of the grounds water from the southern part of the Secaşul Mare Basin are under the direct incidence of the pluvial and pluvio – nivale alimentation regime, with significant increases of the level due to the meltdown of snow and heavy rainfall (during spring time) and an inward withdrawal in the summer and autumn months.

At the northern border of the Cindrel Mountains there lies the **aquifer complex of the piedmont deposits**, with aquifer strata captive at depths of 10 – 15 m, with a basic pH (7,4) and a yellowish

colour, highlighted in the drinkable water springs on the Călnic – Dobârca direction (Trufaş, Ştef 1992). The petrographic constitution is given by gravel and blocks of rocks of large dimensions, encompassed in a sandy and clay – sandy composition, with a high permeability. The permeability coefficient of the alluvial deposits on the Rod and Poienii Valley was registered at 300 – 400 m/day (I.S.P.H. 1998), while the hydrostatic level of ground water was identified of the delluvio – proluvial deposits at depths of over 4 m.

The hydrogeological characteristics are completely different in the hills in comparison to those from the plateau, due to the geological constitution, formed of sedimentary rocks: marl, clay, gravel and sand. The Secaş Plateau is over all lacking underground water reserves and has a few accumulations in a **clayey marls aquifer complex**, with waters manifesting upwards, in an artesian form. These belong to the monocline marginal sub - unit of the Transylvanian Depression (the peripheral monocline, Ujvari 1972) and can be characterized by strong mineralisations and hydrocarbon content (Geografia României, vol. I, 1983). The drillings done by C.Z.M.H. Sibiu (1970) do not indicate the presence of significant aquifer deposits. The petrographic constitution, mainly clay with a low infiltration coefficient concerning the saturation of clay, as well as the little rainfall in the plateau area (compared to the southern part of the basin) both create weak accumulation conditions for underground water in the Badenian, Sarmatian and Pliocene deposits; the only exception is the Quaternary deposits.

Compared to the sub – mountain frame to the Secaşul Mare Basin the density of the plateau springs is further reduced. Their debits are conditioned by the regime of the rainfall and the impermeability of the saturated clay. The reduced slope of the mountain sides from the discharge area of the springs confers them a swamp character. These springs are present between Sebeş and Daia Română (on the Slatina Valley), at Draşov, and at Miercurea Sibiului.

From a qualitative point of view, the natural waters always have several substances dissolved in them, even the rain water contains a solution of substances and gas (carbon dioxide and oxygen). The dissolved substances, especially the gases, increase its dissolution capacity. The chemical composition of subterranean water is fairly varied in the Secaşul Mare basin. The basic processes of the forming, of the evolution and of spreading the mineralisation of the waters in the geological

stratum are governed by hydrogeological and hydrochemical laws. Changes in the waters chemical composition, of their quality depends mainly on the rocks lithology under the chemical substances and the substance exchange between the water – the universal dissolvent – and the parental rock. The direction of the chemical reactions, products of the diverse exogenous chemical processes, but especially of the endogenous ones, depends on the other environmental conditions.

The subterranean waters have a poor quality in the plateau, as a result of the high mineralisation and of the rich NaCl content, caused by the washing of salty horizons on the edges of the diapir structures laid on the Ocna Mureş – Miercurea Sibiului - Ocna Sibiului direction. The emerging springs brought up by erosion in the cuesta of the Secaş Plateau have low debits (0,2 l/s) and are frequently salted. Marosi (1980), points out that in the axial area of the Daia Română – Ocna Sibiului anticline, the tortonian deposits, rich in salt, almost reach the surface and influence the salinity and the type of mineralisation of underground waters. The same author points out, based on several chemical analyses of the drilling waters and of the studies done in this area by Vancea (1960), that contrary to the chlorosodic waters on Daia Valley, which are “simple levigation waters of the tortonian salt (M tot. = 18,1 – 36,7 g/l), the shallow drilling water and those captured in the basins from Miercurea Sibiului (M tot. = 19,3 – 40,2 g/l, type Cl – Na) have a relatively high content of Ca^{++} , Br^- , I^- and NH_4^+ , which indicates the mixture of tortonian levigation water with the waters of the Miocene deposits, or even the predominance of the fossil waters components in the hydro – chemical total of these waters. The few methane gas traces, identified in the superficial capture points of the respective water, also plead in favour of the mixed genesis“. According to the chemical composition, one will find in the Secaşul Mare Basin ferruginous, sulphate, bicarbonate and sodic waters at Vingard⁷, sodic chlorate water at Daia, Miercurea Sibiului and sodic chlorate water enriched with other chemical elements at Cunţa and Cut (Preda, Marosi 1971; Pricăjan 1985). The presence of gas-bearing deposits in the substratum is signaled by the emanations which encounter

⁷ The water has a low mineralisation (Mtot = 11,1 g/l), of pure sulphatic – sodium type, with a high level of Ca and Mg as a result of the presence of diagenetic gypsum and of the lack of upward alimentation of the horizons in the pannonian deposits by the tortonian water (Marosi 1980).

lenses of phreatic line in their way, pushing towards the surface. The latter soften the clay in they upward movement and contribute to the surface appearance of the Boz – Sarata basin muddy volcanoes.

The anthropic intervention in the Secaşul Mare basin, mainly by setting up the fish farming exploitations on the Daia valleys (Slatina and Boz) have had and keep on having a significant impact on the hydrogeological natural regime (Costea 2005, Ştef, Costea 2006). This impact can be seen in the following aspects identified based on the visual observations and on the information provided by the locals:

- the dynamics of the phreatic level in the slope and interfluvial deposits in the vicinity of the accumulation lakes and from their area of influence, is conditioned by the rainfall regime and the downstream transfer;
- the oscillations of the ground water within the main river meadow of Secaşul Mare downstream from Daia and its alimentation are depend on the rainfall regime and on the variation of the debits of the hydrographic artery;
- the water storage in the accumulation lakes lead to: the reduction of the water flow downstream from the lakes, the decrease of the water level from the aquifer stratum and therefore to the decrease of the hydrostatic level with up to 1.5 m in the main river meadow and in its area of influence, especially at the bottom of the slopes;
- the water level in the fish – farming lakes on the Daia and Boz Valleys influences the dynamic of the ground water and its circulation in the slope deposits, as well as in the alluvial deposits form the merging upstream, through their permanent alimentation circuit from the lake;
- the decrease of the phreatic line can also be noticed in the plateau area, where the locals resorted to capturing the springs from the bottom of the slopes and building water accumulation basins – “troughs” – on the pastures and on the streets of the settlements;
- at the bottom of the depression there is also a direct hydrodynamic report between the ground water from the alluvial deposits and the hydrographic arteries, with rises of the piezometric level highlighted by the increased level of fountain water (1m) and by the basements touched by dampness; to all these, the water surplus from the rainy periods is to be added.

Conclusions

The hydrogeological analysis requires a fine tuning of the genetic and evolutionary aspects of underground water to the palaeogeographical evolution, as well as a permanent correlation to the geological elements of the studied region. The geological elements condition the evolution of the landform and its modulation, and therefore represent the key element of several hydrogeological inter conditioning relations, due to the typology of the aquifer deposits present within the landform elements. Given this context, the lithological diversity, the structure and the tectonics are all elements which explain the distribution and evolution of underground water; namely, the following aspects have been identified in the Secaşul Mare basin:

- the petrographic influence, to which the base level, triggering off the erosion - determined by the Transylvanian Depression, respectively by the Mureş Passage - are strong arguments for the morphogenetic shaping of underground water;
- the degree of mineralization, the chemistry of underground water depends on the physical – chemical characteristics of the rocks in the geological substratum;
- the physical properties of the rocks and the stratification manner, the slopes, the compactness, etc.- all determine the intensity of the hydrodynamic exchanges among the rock deposits, the aquifer and surface water;
- the petrographic differentiations are reflected in the landform at various levels storage levels of the underground water, which generate by association, a series of elementary or complex geomorphological processes at the surface, leading to the degradation of the fields (saltiness, streaming, gorging, torrentiality, gravitational sliding);
- the previously described paleogeographical conditions have led to a differential preservation of the landform elements and of the underground water resources: the level surfaces in the plateau, where the phreatic water is to be located at higher depths, the accentuated fragmentation and the springs discharge, permeable piedmont accumulations with significant aquifer resources, but also with active drainage, and the loss of underground water due to the increased infiltration enabled by the friability of the rocks;
- the geological element influences the sliding of the slopes (we quote slope figures between 15 – 75 degrees on the monocline structure) and the

emergence of springs on cuesta fronts or on structural surfaces;

- the hardness, the compactness and high impermeability of the rocks in the Carpathian area explain the low presence of underground water, while the high porosity and low permeability of clay on the lower part of the basin explain the intense dynamic of underground water in the plateau region, which, when associated with thermic processes and lack of rainfall in the drought years, leads to the decrease of underground water resources;

- the geological elements (due to the impermeability of the saturated clay) reduce the infiltration and increase the surface leakage, as well as the accumulation of significant debits, which enables the utilization of the hydrographic network in the plateau for piscatorial purposes, through the setting up of ponds.

Regarding the mineralization, underground water reflects, from a chemical point of view, the entire geological evolution of a certain structural unit. The passing of the minerals from the rocks, by means of levigation (partially) and/or the dissolution, are due not only to the chemical composition of the mineral, but also to several complex chemical processes taking place at the ion level. Any form of natural water – rainfall water, surface or underground water all contain chemical substances, some of which are harmful to humans or any other life forms.

The modification of water chemistry might be due to the dissolution phenomena undergone by minerals and rocks when in contact with water, as well as to the anthropic pollution. The chemical processes taking place at the water – rock contact cannot be controlled, are complicated and often unknown. For example, a significant percentage of the water used for irrigations dissolves the chemical manure from the soil and by means of infiltration it can then reach the rocks, leading later on to unpredictable chemical reactions, rarely beneficial for the quality of the phreatic water. Therefore a permanent monitorization of the quality of underground water, especially of the water utilized in agriculture, fish-farming, households and consumption, is more than necessary.

Acknowledgements

This work was co-financed from the European Social Fund through Sectorial Operational Programme Human Resources Development 2007-2013, project number POSDRU/89/1.5/S/63258 "Postdoctoral school for zootechnical biodiversity and food biotechnology based on the eco-economy and the bio-economy required by eco-san-genesys". The authors are grateful to Ana Maria Mesaroş for constructive comment on our manuscript and to Irina Ciobanu for the useful suggestions improving the language for the text. The authors would like to thanks to anonymous reviewer for helpful comment that improved the manuscript.

REFERENCES

- Albu 1981 Albu Mihai, *Mecanica apelor subterane*. In: *Editura Tehnică*, Bucureşti (1981).
- Băncilă *et al.* 1980 Băncilă Ioan *et al.*, *Geologie inginerească*. In: *Editura Tehnică*, Bucureşti (1980).
- Castany 1972 Castany George, *Prospectarea şi exploatarea apelor subterane*. In: *Editura Tehnică*, Bucureşti (1972).
- Chivu 1978 Chivu C., *Contribuţii la cunoaşterea geologiei şi metalogenezei părţii de nord a Munţilor Sebeş*. In: *Dări de seamă ale şedinţelor IGPSMS LXIV (1976-1977)*, Bucureşti (1978).
- Ciobanu 2002 Ciobanu Rodica, *Depresiunea Sibiului. Studiu geologic*. In: *Editura Universităţii „Lucian Blaga”*, Sibiu (2002).
- Ciobanu 2006 Ciobanu Rodica, *Elemente geologice favorabile şi restrictive în evoluţia reţelelor hidrografice*. În: *Hidrologie aplicată*, Ştef Vasile, Costea Marioara (eds.), Editura Universităţii “Lucian Blaga”, Sibiu (2006), p. 21-49.
- Ciocârdel 1957 Ciocârdel Radu, *Hidrogeologie*. In: *Editura Tehnică*, Bucureşti (1957).
- Ciortu, Păcurar 1998 Ciortu Ioan, Păcurar Vasile, *Curs de Hidrologie şi hidrogeologie*. In: *Universitatea „Transilvania” Braşov, Facultatea de Silvicultură şi Exploatare Forestiere*, Braşov (1998).
- Ciupagea *et al.* 1970 Ciupagea, D., Paucă Mircea, Ichim Traian, *Geologia Depresiunii Transilvaniei*. In: *Editura Academiei Române*, Bucureşti (1970).
- Codarcea 1962 Codarcea Dessilla Marcela, *Încercare de reconstituire paleogeografică şi orogenetică a Carpaţilor Meridionali Centrali*. In: *St.şi cercet.de geologie VII*, Bucureşti (1962), p.3-4.
- Costea 2005 Costea Marioara, *Bazinul Sebeşului, Studiu de peisaj*. In: *Editura Universităţii “Lucian Blaga”*, Sibiu (2005).
- Huică *et al.* 1972 Huică Ioan, Lubenescu Victoria, Clemens Antoaneta, Cehlarov Aura, Bârlea Lidia, *Date noi asupra geologiei regiunii Ocna Sibiului – Alba Iulia – Sebeş (Sud – vestul bazinului Transilvaniei)*, In: *SCGGG – Geol.* 17 (2), Bucureşti (1972).
- Ilie 1955 Ilie Mircea, *Cercetări geologice în bazinul Transilvaniei, (Regiunea Alba Iulia, Sibiu, Făgăraş, Rupea)*. In: *An.Com.Geol. XXVIII*, Bucureşti(1955).
- Marosi 1980 Marosi Petre, *Hidrogeologie, Capitole alese pentru uzul studenţilor*. In: *UBB, Cluj-Napoca* (1980).
- Mutihac 1990 Mutihac Vasile, *Structura geologică a teritoriului României*. In: *Editura Tehnică*, (1990) Bucureşti.
- Paucă 1971 Paucă Mircea, *Criterii noi în stratigrafia miocenului din bazinul Transilvaniei*. In: *SCGGG –Geol.* 16 (2), Bucureşti (1971).
- Paucă 1972 Paucă Mircea, *Etapale genetice ale Depresiunii Transilvaniei*. In: *SCGGG –Geol.* 17 (2), Bucureşti (1972).
- Pavelescu 1955 Pavelescu Gheorghe, *Cercetări geologice în Munţii Sebeşului*. În: *An. Com.Geol. XXVIII*, Bucureşti (1955).
- Pişota 1995 Pişota Ioan, *Hidrologie*. In: *Editura Universităţii*, Bucureşti (1995).
- Popa 2011 Popa Dana Cătălina, *Consideraţii generale asupra impactului detectării unor nutrienţi în apele de suprafaţă şi de fântână*. In: *Colecţia de cărţi de "Cercetare Postdoctorală în Biodiversitate zootehnică şi Biotehnologii alimentare pe baza Economiei şi bioeconomiei"*, (2011), p. 125-136.
- Posea 2002 Posea Grigore, *Geomorfologia României*. In: *Editura Fundaţiei „România de Măine”*, Bucureşti (2002).
- Preda, Marosi 1971 Preda Ioan, Marosi Petre, *Hidrogeologie*. In: *Editura Didactică şi Pedagogică*, Bucureşti (1971).
- Pricăjan 1985 Pricăjan Aurel, *Substanţe minerale*. In: *Editura Ştiinţifică şi enciclopedică*,

- București (1985).
- Raboca 1995 Raboca Nicolae, *Podișul Secașelor. Studiu de dinamica versanților*. In: Editura „Sarmis”, Cluj – Napoca (1995).
- Rodeanu 1926 Rodeanu Ioan, *Observații geomorfologice la zona de contact a bazinelor Oltului și Mureșului în județul Sibiu*. In: *LIGUC* (1926) II, București (1924 – 1925).
- Romanescu 2003 Romanescu Gheorghe, *Hidrologie Generală*. In: Editura Terra Nostra, Iași (2003).
- Sandu 1998 Sandu Maria, *Culoarul depresionar Sibiu – Apold. Studiu geomorfologic*. In: Editura Academiei, București (1998).
- Savu 1998 Savu Haralambie, *Origin and evolution of a Carpathian triple junction*. In: *AUB, Geol.* XLVII, București (1998).
- Semenenko et. al 1963 Semenenko N.D., Dessila - Codarcea Marcela, Demidenko S. G., Zeidis B., *Interpretarea geologică a datelor de vârstă absolută a formațiunilor cristalofoliene din Carpații Meridionali Centrali*. In: *St. Com. Geol.* VIII (4), București (1963).
- Ștef, Costea 2006 Ștef Vasile, Costea Marioara (eds.), *Hidrologie Aplicată*. In: Editura Universității “Lucian Blaga”, Sibiu (2006).
- Trufaș, Ștef 1992 Trufaș Valer, Ștef Vasile, *Chimismul râurilor din bazinul hidrografic al Mureșului*. In: *AUB XXXIX*, București (1992).
- Trufaș 1978 Trufaș Valer, *Apele subterane din Munții Sebeșului*. În: *St. Și cercet. de geogr.* 15, București (1978).
- Trufaș, Ștef 1998 Trufaș Valer, Ștef Vasile, *Hidrogeologie*. In: *Facultatea de Geografia Turismului*, Sibiu (1998).
- Vancea 1960 Vancea Augustin, *Neogenul din Bazinul Transilvaniei*. In: Editura Academiei, București (1960).
- Zamfirescu 1997 Zamfirescu Florin, *Elemente de bază în dinamica apelor subteran*. In: Editura Didactică și Pedagogică, București (1997).
- * * * *Harta geologică a RS România, scara 1: 200 000. Foaia Orăștie*. In: Institutul Geologic (1968) București.
- * * * *Geografia României, I., Geografie Fizică*. In: Editura Academiei (1983) București.
- * * * *Geografia României, III., Carpații și Depresiunea Transilvaniei*. In: Editura Academiei (1987) București.

LIST OF ILLUSTRATIONS

- Fig. 1.** Lithological map of Secașul Mare Basin
- Fig. 2.** Râpa Roșie – red sands and slightly cemented clay outcrop (in the Secașelor Plateau)
- Fig. 3.** Crystalline outcrop in the epigenetic sector of the Călnic river (the southern part of the Secașul Mare Basin)
- Fig. 4.** Saline efflorescence at the bottom of the Potca hill in Apoldu de Jos, on the southern slopes
- Fig. 5.** The typology of phreatic water in the Secașul Mare Basin
- Tab. 1.** The variation of porosity depending on the type of rock

LISTA ILUSTRAȚIILOR

- Fig. 1.** Harta litologică a Bazinului Secașul Mare
- Fig. 2.** Râpa Roșie – afloriment de nisipuri roșietice și argile slab cimentate (în Podișul Secașelor)
- Fig. 3.** Afloriment cristalin în sectorul epigenetic al râului Călnic (partea sudică a bazinului Secașului Mare)
- Fig. 4.** Eflorescențe saline la baza dealului Potca la Apoldu de Jos, pe versanți cu expoziție sudică
- Fig. 5.** Tipologia apelor freatice în Bazinul Secașului Mare
- Tab. 1.** Variația porozității în funcție de tipul de roci




Fig. 1. Lithological map of the Secaşul Mare Basin




Fig. 2. Râpa Roșie – red sands and slightly cemented clay outcrop (in the Secaşelor Plateau)




Fig. 3. Crystalline outcrop in the epigenetic sector of the Călnic river
(the southern part of the Secaşul Mare Basin)



Fig. 4. Saline efflorescence the bottom of the Potca hill
in Apoldu de Jos, on the southern slopes

Fig. 5. The typology of phreatic water in the Secaşul Mare Basin



Tab. 1 The variation of porosity depending on the type of rock

Type of rock/ soil	Observations	P%
Soil	The porosity depends on the soil components.	30-50
Clay	<i>Very porous</i> rocks, made up of small particles, with very numerous spaces between them, but so small that the capillary forces retain the water which can no longer flow around the rock – <i>very low permeability</i>	10-50
Sand	The porosity varies to a great extent, according to the purity and cementation level, namely: -uncemented sands – with the highest porosity (30-33%), - the particle content from clay decreases the porosity by 18-24%, - cemented sands – the porosity decreases down to 3-10%.	18-47
Grovel	High porosity increasing by the size of the size of the granules.	
Sandstone	The porosity of the gritstones depends on the nature of the cement (its resistance to the action of the water), of the matrix and of the structure, if they are part of strongly tectonic structures, the splitting degree is high and increases the porosity.	5-15
Limestone	They are not porous rocks, but the tearing discontinuities such as the lithoclasts increase the permeability of the rocks.	2,5-20
Quartzite	The finely grained rocks with a low porosity; generally, the alteration of the silicates contributes to the increase of porosity.	0,5-1

MORPHOLOGICAL AND HYDROLOGICAL DYSFUNCTIONS IN THE URBAN SPACE OF SIBIU MUNICIPALITY

Marioara COSTEA *

Abstract: *This paper emphasizes current dysfunctions of the urban system deriving from the relation between geological substrate-morphodynamic-hydrological processes and proposes a set of geographical space optimization solutions. Extant imbalances are caused by natural factors whose action is intensified by the anthropogenic factor framework. The geospatial arguments were presented with the purpose of advancing an urban development strategy for Sibiu Municipality through the General Urban Plan 2009. Within this framework, we propose a set of measures for the prevention and reduction of the natural dysfunctions, geomorphologic and hydrological in the Sibiu area in order to capitalize in a balanced and durable manner its natural potential. These may be diminished by ensuring the appropriate exploitation of the geographical space, through a set of constant development and hydro-amelioration activities and controlled urban expansion.*

Key words: *geomorphologic and hydrological dysfunctions, urban space, Sibiu.*

Rezumat: *Disfuncționalități geomorfologice și hidrologice în spațiul urban al municipiului Sibiu. Prin lucrarea de față urmărim evidențierea disfuncțiilor actuale ale sistemului urban care derivă din relația substrat geologic – morfodinamică – procese hidrologice și să propunem unele soluții de optimizare a spațiului geografic. Dezechilibrele existente la momentul actual sunt cauzate de factori naturali a căror acțiune este accentuată pe un fond puternic antropizat. Evidențierea unor argumente geo-spațiale temeinic fundamentate s-a realizat în vederea elaborării strategiei de dezvoltare urbană a municipiului Sibiu prin PUG 2009. În acest context propunem unele măsuri de prevenire și diminuare a disfuncționalităților naturale - geomorfologice și hidrologice de pe teritoriul municipiului Sibiu, în vederea unei valorificări echilibrate și durabile a potențialului natural. Acestea pot fi diminuate în condițiile unei exploatare corecte a spațiului geografic, în condițiile unor lucrări susținute de amenajare și hidroameliorare și a expansiunii edilitare controlate.*

Cuvinte cheie: *disfuncționalități geomorfologice și hidrologice, spațiu urban, Sibiu.*

Introduction

The geographical space situated between the administrative limits of the Sibiu Municipality is a highly modified by human action; it is mainly stable, but there are areas characterized by slope imbalance and intensive exploitation. The anthropic transformations on the Sibiu Municipality administrative area have been gradual and in connection with key moments in Sibiu's social and economic development. The human impact side-effects in the Sibiu depression were the transformation of morphometric and morphographic features of the landform through cutting, levelling or soil stripping uncovering and a major impact through the acceleration of geomorphologic processes in certain areas, which rendered those areas hazardous and increased their risk rate. (Costea, Gherasim, 2011, p.24).

The hydrographic network was anthropically altered through rectification, embankment, exploitation of building materials from the riverbed, etc.), the vegetation (massive deforestation, occurrence of ruderal species, extinction of spontaneous species), the soils (change in productivity through liming, drainage etc.) as well as on the climate conditions (by creating complex and basic new topoclimates) (Costea 2002). The most representative dysfunctions are those in the geomorphologic and hydrologic system. They are intricately connected with the geological conditions and space utilization.

Lithologic and structural particularities

Sediment deposits in the analyzed perimeter belong to young structures, being represented by Miocene, Pliocene and Quaternary sediments, differing in terms of their degree of cementation,

* "Lucian Blaga" University of Sibiu, Faculty of Sciences, marioara_costea@yahoo.com

consolidation, resistance to environmental factors. Pannonian deposits occupy most of the area and contain clays, loamy sands, and marl, with a very low degree of cementation. They come into contact with crystalline formations and crop out currently in Gușterița Hill and on the face of the Hârtibaciu Plateau cuesta (Ciobanu *et al.* 2010), whose eastern and north-eastern part fall within the administrative limits of Sibiu Municipality. The Quaternary deposits are the most recent and they cover almost entirely administrative territory. They are represented by terrace deposits and proluvial deposits (dejection cones) from Pleistocene and differ from one another in terms of their genesis, thickness, and granulometric structure (gravels), tending to form conglomerates (sands included in clay mass) covered by a layer of soil with variable thickness, as well as current alluvial deposits (Holocene gravels, sands, silts) very frequent in the meadows of Cibin River and its tributaries.

Dysfunctions caused by the lithological structure and their negative effects upon the urban habitat consist in the following aspects:

- damping and swelling of the clayish substrate affecting the internal balance of the substrate and predispose the Hârtibaciu Plateau slopes and the terrace faces of the Cibin River to gravitation processes such as landslides;
- the friability and weak cementation of the rocks facilitate surface pluviodenudation and erosion processes, with aggressive manifestations such as streaming, ravining and torrential processes, especially on the Gușterița Hill slopes or the faces of the Cibin River terraces (Ciobanu, 2002);
- the high quantity of clayish silts in the alluvial deposits of Cibin River and its tributaries and their impermeability cause the groundwater level to rise in river meadows, with certain permanent or temporary ponding;
- unconsolidated terrace and meadow sediment deposits, associated with hydro-climatic oscillations and the pressure of their weight or of the weight of the buildings above which, in time, cause gravitational re-settlement and suffusion which may have negative effects upon the structure of those buildings;
- the swollen clays which compose the deposits due to over-humidification and the contracting of those clays during droughty periods cause deep fissures in the soil and geological formations; these fissures are access ways, through infiltration, for meteoric waters during rainy periods, which may cause infiltrations and dampness to walls, in the basement of the buildings or even the flooding of some open, unprotected spaces. Moreover, the

water accumulated as aquifer deposits between clayish surface deposits, through the capillarity of the soil, infiltrates in the walls causing dampness.

Geomorphologic potential and relief dynamic

The landforms in the Sibiu Municipality area are characterized by morphometric and morphologic features which disclose, on the one hand, the geological influence and on the other, the present-day modelling conditions. The morphology of the Sibiu Municipality and its peri-urban area belongs to the sub-mountain depression and plateau morphogenetic type, differentiated according to the landforms. At the level of the entire Sibiu Depression, but especially between the administrative limits of the Sibiu Municipality, there is a clear landform asymmetry imposed by Cibin River's tendency to shift position north and north-east; this fact is revealed by an altitudinal decrease on the right-side riverbank, as landform steps (sub-mountain hills, piedmonts, terraces, meadows) and on the left-side riverbank by the Hârtibaciu Plateau gully, affected by gravitational processes (Costea 2007).

Altitudes, in the analyzed sector, fall between 400 m in the Cibin River meadow on the administrative limit with the Șelimbăr Commune and 639 m in South-East Hamba Hill on the Hârtibaciu Plateau cuesta, the average altitude being over 450 m. The 400 m isohypse circumscribes the Cibin River meadow in this sector and insinuates at the basis of the structural steep of the plateau. The density of fragmentation is characterized by values between 0 – 0.5 km/km² in the Cibin River's meadow and terraces and up to 2 – 2.5 km/km² on the front of the Hârtibaciu Plateau cuesta and on the terrace front. The landform energy varies depending on the landform form, with values from 0 - 10 m at the level of the Cibin River's alluvial plain, up to 80 - 140 m at the contact between meadow and the face of the cuesta. The slopes have differentiated values on the analyzed sector, from minimum values of 0 – 3 - 5° at the level of the major bed of Cibin River and its tributaries and on terrace bridges, values of 10 -15 - 35° typical of terrace faces, structural steep and its basic glacis, up to values of over 60° in the perimeter of the clay quarries in the Gușterița Hill. (Costea, Gherasim, 2011, p.19). The most frequent slopes are those ranging from 10 to 30°, values which facilitate pluviodenudation processes and gravitational soil slips (Costea 2002).

The current dynamic of the landform is induced mainly by natural factors: the lithologic and structural contact between the crystalline of and the two mountain units and the sediment deposits

of the Hârtibaci Plateau, climate with oceanic influences as well as the action of autochthonous and allochthonous hydrographic network, on a foundation consisting of gravels, sands, marls, and clays, with harder nuclei. Predominant are slope processes caused by pluviodenudation: surface soil running, streaming, ravinement, torrentiality (Greco 1992, 1996). The erosion, transport, accumulation processes succeed one another in the allochthonous valley of the Cibin River with high flows, which discharged huge quantities of materials as huge piedmont dejection cones - the Cibin River Piedmont and glacises, which were taken on by the autochthonous hydrographic network - Valea Aurie, Valea Săpunului, with variable flows, contributing to the fragmentation of these cumulative structures and to the degradation of the of inter-river surfaces and versants by the regressive retreat of the sources on rank I and II hydrographic basins (Costea 2007, Costea, Gherasim, 2011).

The argillaceous clay component of the terrace deposits is responsible for the presence of aquiferous layers close to the surface and for the water infiltration. On the terrace faces and at their base, especially in extravilan, there are areas with swamps and springs. The third terrace in the Poplaca Pasture, beyond the northern skirt of the Dumbrava forest, is divided by the ravines, gaps, and short or torrential valleys (Valea Urzicarilor), tributaries of the Cibin River upstream Turnișor. These fragment the front of the terrace and deepening through regressive erosion in the bridge of upper terrace. The front of the terrace which makes the connection with the Cibin River meadow is highly degraded (Costea *et al.* 2011) through geomorphologic processes such as landslides and torrential processes (Fig.1).

In the bed of the Cibin River and its right-side tributaries - the Trinkbach creek (Valea Aurie) and Valea Săpunului, in the meandering sectors we witness phenomena such as accumulation of convex riverbanks and erosion of concave riverbanks, the latter undermining slopes and facilitating landslides and collapsing). The meadow of the Cibin River is under the direct incidence of the water flow and versant processes in limitrophe areas (the plateau cuesta), where the most frequent phenomena are superficial slides, collapses, torrentially, ravinement, etc.

Thus, in the Turnișor upstream sector, some of the meanders of the drained Cibin River maintain their connection with the riverbed, taking some of the flow volume when the water level rises, but others are entirely or partially clogged and are not

connected with the river anymore. During rainy periods, these meander loops (rings) may become lacustrine or swamp areas. Nevertheless, the creeks which fragment the Cibin River terraces in this sector feed the meander rings forming ponds and swamps which, given the low slope and the silty substrate, do not drain very easily, preserving dampness and facilitating the development of hydrophytic vegetation. In addition, these torrential organisms discharge dejection cones in the Cibin River meadow which push the thalweg of the Cibin River and redirect processes to the riverbed (Costea 2007).

In addition, in the sector between the confluence with the Rozbav River and the south-eastern limit of the incorporated area has relative altitudes of 1 – 1.5 m and the dynamic manifestations which are typical to the meadow landform are: bank erosion, alluvial accumulations, phreatic moistening, and gleying. The accumulation processes are more active in this sector because on the section upstream-downstream of Gușterița, the Cibin River receives torrential tributaries with sources in the plateau: Valea Fărmândoala, with its tributary Valea Nepîndoala, Valea Pe Remeți. These carry and deposit in the collector the solid materials resulted from fluvial-torrential erosion. The accumulations in the riverbed are located laterally as beaches fixed on the vegetation, or centrally as islets totally or partially fixed on the vegetation, such as those near the Gușterița bridge, or as submerged deposits downstream Gușterița. The bridge legs are obstacles for alluvial deposits and they gradually influence the formation of submerged or emerged deposits on the riverbed.

Morphodynamic natural factors are amplified by anthropic factors, their role in modelling the landforms being very complex, in connection with the amplitude and diversity of the activities performed (Posea *coord.* 1987). Such activities are: agricultural, pastoral, industrial activities, deforestation, river management activities, drainage, plantations, urban expansion and land development for social, economic or amusement activities. Geomorphologic processes have been increased because of deforestation (forests cover today only 20% of the areas previously occupied by forest vegetation) and they affect the depression area especially the piedmonts and structural steeps which delimit the administrative area of the Sibiu Municipality.

Today, the abandonment of agricultural land and the building of sheepfolds around streams or in the middle of a wide field in the northern part of the built-up area are the main causes of soil

degradation through nitrification, of surface erosion and the destruction of the grass. The orchards from Gușterița Hill have been mostly abandoned and thus the area is subjected to moderate degradation but the process might intensify in the long term. Anthropogenic terraces are modelled by current geomorphologic processes which are constantly intensifying and which are reducing land productivity.

Excavations performed through the extraction of clay for brick production in Gușterița Hill (Sibiu) or for the building of the belt highway (in the Șura Mare Hill) are the premises for versant undermining and intensification of processes such as streaming, ravinement, sliding and collapsing and versant degradation. The diluvial contribution due to gravitational processes causes versant agglomeration and the formation of a highly unstable sliding glacis. Moreover, the presence of thick clay layers and their intensive exploitation for industrial use or road construction explain the slopes morphodynamic and the amount of colluvial and proluvial deposits in the riverbeds which fragment the cuesta (the Nepîndoaia Creek, Valea Fărândoala, Valea Pe Remeți).

The road and railway traffic cause certain imbalances due to mechanical stress (such as subsidence, sliding, collapses), the destruction of riverbank protection where hydrographic arteries are accompanied or crossed by roads intensely used (crossing the Cibin River), the building of bridge legs (over the Cibin River and its tributaries) which operate as obstacles for the alluvial deposits (Cibin River, the Gușterița bridge), as well as the undersized bridges and footbridges which cross the hydrographic network and over which there is frequent intense traffic.

Hydrodynamic phenomena - induced dysfunctions

Territorial hydrological dysfunctions are caused by stationary (excess humidity) and active (high-water, freshets, flooding) hydrodynamic processes (Fig.1).

The underground waters located near the surface cause excess humidity especially in river meadows and in the alluvial plain north of Sibiu. The low slope and the substrate facilitate water retention and gleying in the alluvial plain of the Rozbavului River, Valea Popilor and Valea Hamba. Here and there, swamps and marshy pastures have appeared. The draining activities performed in the early 20th century, continued between 1956 and 1975, producing a dense network of channels in this sector, 1.5 to 2 m deep, which contributed to the

drainage of this area and the lowering of the groundwater level. Once certain plots have been resituated to their owners as private property, drainage projects were abandoned. Most of these channels are currently clogged, some were used as garbage pits or debris dumping places, so that in this area, during rainy periods, there is excess humidity and the groundwater level rises (Ștef, Costea 2006).

Regarding the active hydrodynamic processes, when the flow volume is high the meadows of the Cibin River and of its tributaries are vulnerable to flooding and the obstacles in the riverbeds facilitate this phenomenon. The development works carried out on the River Cibin at Gura Râului have restricted these hazardous phenomena by regulating flows and reducing flash floods. However, downstream, the Cibin River receives tributaries which may cause imbalances in the Sibiu Municipality area, due to torrential precipitation and inappropriate use of land (Sandu 1998). Several areas on the Cibin River are exposed to flooding: the Cibin River meadow upstream Turnișor, where garbage and debris clog the major riverbed and increase hydrological risks, the Cibin River meadow in the Gușterița area - the river takes an almost 90° changing its direction; thus, the junction with the Fărândoala creek, the bridge and the obstruction of the flow by islets are major hydrological risk factors.

Even though the Trinkbach creek has been harnessed, sometimes certain estates in the area are flooded. These phenomena are sometimes caused by the fact that accumulations are not performed according to the established parameters, but especially by the fact that this creek has been streamlined, its bed has been concreted and is undersized. When there are flash floods, the water in the minor riverbed infiltrates in the basement of nearby houses in the covered sector.

In conclusion, hydrological risks in the Sibiu Municipality area may be avoided through an appropriate exploitation of existing reservoirs during rainy periods and only if hydroamelioration measures are taken.

Reducing geomorphologic imbalance

As it is described above, geomorphologic imbalance is caused by external agents whose action is accelerated given the favourable geological circumstances and current land use. There are many factors of degradation taking place randomly all over the sector (Grecu, Sandu 2000). The preservation of landform forms as a support for the urban structure and the

decrease in negative effects of current morphodynamic require that the following measures be taken:

- attenuating the intensity of linear and deep erosion in the small hydrographic basins of the Hârtibaci Plateau: Valea Fărmâdoala, Valea Nepîndoaia, Valea Pe Remeți, Valea Hamba and on the ravines and torrents which fragment the faces of the terraces in the Poplaca Pasture through hydrotechnical, forestry and agrotechnical operations so as to transform linear into non-linear flow;
- improving the natural drainage of the substrate through hydro-ameliorative, pedo-ameliorative works (levelling, terracing) in areas affected by landslides in Gușterița Hill, Cocoșului Hill, Gumușel's Hill, Fântâna Rece; we recommend landslide prevention through afforestation and by planting fruit trees;
- intercepting terrace and coast permanent springs through water catchment methods (wells, horizontal pipelines), canalization at terrace base or of the slopes in the plateau;
- eliminating excess water from terrace treads, where possible, by afforestation or grassland vegetation, to reduce moisture;
- stopping bank undermining through bank and slope stabilization operations
- rehabilitation of the areas populated with *Pinus sylvestris* at the source of the Nepîndoaia stream;
- stopping and reducing landslides by building reinforcement walls and buttresses;
- populating the area with small and medium sized species in order to compact the grassland vegetation and reduce surface and linear erosion;
- rehabilitate the orchards from Gușterița Hill in order to consolidate again the ground;
- build natural dams as small fences to reinforce drains, ravines and gaps and reduce deep and regressive erosion;
- ban plowing and reconsider agricultural techniques;
- avoid over-grazing in the torrents catchments (Nepîndoaia, Remețe);
- avoid excavations for communication path and roads at the base of the slopes in the Gușterița area;
- avoid and ban excavations and exports of clay material from the base of slopes in Gușterița Hill;
- ban the expansion of the clay quarry for brick manufacturing at the expense of forest areas;
- ban and stop building operations on slopes to landslides and torrentiality in the Gușterița Hill area and on the terrace faces in incorporated and unincorporated areas of the city (Low 350/2001);

- consolidate the Gușterița Hill front cuesta to prevent new collapses and slides facilitated by the heavy traffic on the belt highway;
- ban the construction of storeyed buildings on the terrace faces in built-up areas;
- ban the overloading of terrace edges with buildings;
- ban the building of housing facilities in the Calea Poplăcii neighbourhood - extension Izlazului in the unstable waste materials warehouse and populating the area with common trees and fruit trees;
- ban deforestation for building purposes in the piedmont area - the Cibin River Piedmont and third terrace as this facilitates landslides.

Hydrological phenomena prevention and reduction

The spatial durable development and evolution of the city of Sibiu requires thorough awareness of the hydrological regime of the area situated between its administrative limits, in unincorporated areas as hydrological flow supply and relieve area, but especially in incorporated areas as reception and transfer areas. The city is subjected to a natural hydrological regime through its hydrographical network (including lakes) which converges in the Cibin River, a river that crosses the city and collects tributaries especially in the eastern part of the city (Ștef, Costea 2006; Stănescu 1995).

The highly developed areas and the sewage transportation systems are elements which require good knowledge of the hydrological drainage regime, especially in built-up areas, where concreted, asphalted surfaces and undersized sewage systems may entail the flooding of the urban area. On the other hand, the fact that the city is located in an area with a moistened substrate and lower slopes, raise hydrogeological issues, issues related to the surfaces drainage towards the stable hydrographic network and the issue of Hydrological risk phenomena caused by hydrological stationary processes.

Within this framework, steps should be taken in order to harness rivers, optimize surface drainage, but also improve pluvial water drainage through the underground urban network and reduce hydrological risks. Thus, our recommendations aim at reducing the risk generated by both stationary and hydrodynamic processes through:

- building drainage canals on humid flat surfaces and rehabilitating the extant ones; the surface drainage systems in Șura Mică and Viile Sibiului and Lazaret neighbourhood should be deepened to the level of the aquiferous deposit;
- building depth drainage systems, performing hydro-ameliorative operations for underground

collection and drainage (wells, horizontal pipelines) to eliminate underground water surplus;

- monitoring the operation of the Gura Râului dam, so as to control the flow;

- harnessing small hydrographic basins and regulating the flows of the tributaries within city limits and regulating the Cibin River within the city limits and at the location where it leaves the city (Low 170/1996);

- building of parks and riverside coppice on the riverbanks of Cibin and its tributaries meant to use and redistribute the water surplus in the meadows;

- developing certain controlled wetlands in Lunca Mare meant to harness the abandoned meanders of Cibin river upstream Turnișor;

- optimizing underground drainage by changing the sewage system size depending on the average quantity of pluvial water and domestic and industrial consumption;

- building marginal canals and surface drainage systems (ditches, gutters), especially on the sloping streets, meant to ensure the transfer of the surplus water to natural collectors or underground drainage systems;

- populating the plateau area with common trees and fruit trees in order to ensure the transfer of pluvial water responsible for flash floods on creeks such as: Hamba, Valea Popilor, Valea Fărmândoala, Valea Pe Remeți;

- taking measures to clear the Cibin River bed at the Gușterița bridge;

- banning the disposal of domestic waste or debris on the riverbanks of Cibin River or its tributaries ; these can be worn in riverbed and can clog the flow; the Trinkbach creek requires special attention, as it may flood the skating rink-Lazaret sector;

- preventing new bridge legs from being built in riverbeds and clearing the riverbeds;

- avoiding building undersized bridges and foot bridges which may be blocked and cause afflux phenomena, thus flooding upstream areas;

- banning the building of facilities in easily flooded areas of the Cibin meadow or that of its tributaries and avoiding the narrowing of minor riverbeds.

Conclusions

The development strategy of the city of Sibiu must aim at fortifying the urban territorial system by harmonizing both natural and anthropic factors for the durable development and optimum use of the land (PATN 1998; Ianoș 2000). The role of this strategy is to ensure environmental protection and preservation, especially to set the limits of anthropic intervention upon the environment through urban expansion and economic activities, so that this intervention might be beneficial to the society.

Therefore, the durable development strategy of the Sibiu Municipality must be based on the efficient and durable use of the natural framework, on the reconciliation between spatial evolution and the favourability and restrictions imposed by the physical-geographical framework, on promoting efficient environmental policies within the context of economic development, on the harmonization of functional zoning and space suitability for certain purposes, as well as on promoting an adequate and efficient management of unincorporated areas (Surd *et. al.* 2005; Low 557/2001).

Acknowledgements. This work was co-financed from the European Social Fund through Sectorial Operational Programme Human Resources Development 2007-2013, project number POSDRU/89/1.5/S/63258 "Postdoctoral school for zootechnical biodiversity and food biotechnology based on the eco-economy and the bio-economy required by eco-san-genesis".

The author also thanks the reviewers for suggestions and language comments and corrections.

REFERENCES

- Ciobanu 2002 Ciobanu Rodica, *Depresiunea Sibiului. Studiu geologic*. In: *Editura Universității „Lucian Blaga”*, 120 p., Sibiu (2002).
- Ciobanu et al. 2010 Ciobanu Rodica, Giușca Roxana, Vonica Ghizela, *Dealul Gușterița. Considerații geologice, geomorfologice și botanice*. In *Brukenthal Acta Musei V.3.*, (2010) Sibiu, p.661 – 677.
- Costea 2002 Costea Marioara, *Tipologia peisajului geografic din Depresiunea Sibiului din perspectivă sistemică*. In *Alpii Transilvaniei 5*, Sibiu (2002), p. 61 – 66.
- Costea 2007 Costea Marioara, *Riscul geografic în aria depresiunilor circumcarpatice interne. Studiu de caz – Depresiunea Sibiului*. In: *Acta Musei Tutovensii II*, Bârlad (2007 a), p. 135- 145.
- Costea et al. 2011 Costea Marioara, Giușcă Roxana, Ciobanu Rodica, *Geomorphologic Risk Modelling of the Sibiu Depression Using Geospatial Surface Analyses*. In: Thakur, J.K.; Singh, S.K.; Ramanathan, A.; Prasad, M.B.K.; Gossel, W. (Eds.), *Geospatial Techniques for Managing Environmental Resources*, Springer & Capital Publishing Company, 280 p., (2012), p. 119 – 139.
- Costea, Gherasim 2011 Costea Marioara, Gherasim Virginia, *Anthropogenic changes in the geomorphological System. A case study: Sibiu Depression (Sibiu–Tâlmăciu sector)*, Transylvania, Romania. In: *Acta Oecologica Carpatica*, IV, Sibiu (2011), p.17 – 26. <http://reviste.ulbsibiu.ro/actaoc/13AOCIV2011Papers.pdf>
- Grecu 1992 Grecu Florina, *Bazinul Hârțibaciului. Elemente de morfohidrografie*. In: *Editura Academiei Române*, 167 p., București (1992).
- Grecu 1996 Grecu Florina, *Expunerea la risc a terenurilor deluroase*. In: *Lucr. A 2-a Conferințe Regionale de Geografie (1994)*, Timișoara (1996), p. 18 – 24.
- Grecu, Sandu 2000 Grecu Florina, Sandu Maria, *The Hârțibaciu Tableland. Geomorphological hazard*. In: *Studia Geom. Carpatho–Balcanica 34*, Bukarest (2000), p. 95 – 105.
- Ianoș 2000 Ianoș Ioan, *Sisteme teritoriale. O abordare geografică*. In: *Editura Tehnică*, București (2000), 197 p.
- Posea (coord), 1987 Posea Grigore (coordonator), *Carpații și Depresiunea Transilvaniei*. In *Geografia României III*, In: *Editura Academiei RSR*, București (1987), 647 p.
- Sandu, 1998 Sandu Maria, *Culoarul depresionar Sibiu-Apold. Studiu geomorfologic*. In: *Editura Academiei*, București (1998), 176 p.
- Surd et al. 2005 Surd Vasile, Bold Ion, Zotic Vasile, Chira Carmen, *Amenajarea teritoriului și infrastructuri tehnice*. In: *Editura Presa Universitară Clujeană*, Cluj-Napoca (2005), 585 p.
- Stănescu 1995 Stănescu Vasile A., *Hidrologie urbană*, In: *Editura Didactică și Pedagogică*, București (1995), 99p.
- Ștef Costea 2006 Ștef Vasile, Costea, Marioara (eds.), *Hidrologie aplicată*. In: *Editura Universității „Lucian Blaga”*, Sibiu (2006), 297 p.
- Planul de Amenajare a Teritoriului Național*, (1998).
- Legea apelor 170/1996*.
- Legea 350/2001 privind amenajarea teritoriului și urbanismul*.
- Legea 557/2001 privind aprobarea PATN, secțiunea a V-a, zone cu risc natural*.

LIST OF ILLUSTRATIONS

Fig. 1. Geomorfodynamic map and risk induced by geomorphological and hydrological processes

A. Landforms and morphodynamic potential;

1. Piedmont areas with low vulnerability to geomorphological risk due to the forestal use;
2. The 3rd terrace of Cîbin river at relative altitude of 40 – 50 - 70 m;
3. The 2nd terrace of Cîbin river at relative altitude of 15 – 25 m;
4. The 1st terrace of Cîbin river at relative altitude of 5 – 10 – 15 m; a. Terrace surfaces with high stability and low-moderate morphodynamic caused by and pluvial denudation and compaction processes; b. Sharp of terraces with high morphodynamic caused by ravening, torrential and landslides processes;
5. Cîbin meadow and alluvial plains of the tributaries with high stability and low-moderate morphodynamic, with common stationary hydrodynamic processes and fluvial processes;
6. Plateau interfluvies with moderate stability and moderate morphodynamic caused by gravitational processes and pluviudenudation;
7. Scarp slope of cuesta with high instability and accentuated morphodynamic caused by active processes developed on the clayey deposits and monoclinical structure, predominant landslides and rock-falls, associated with gullying and torrential erosion;
8. Structural surface with moderate to high stability, with moderate morphodynamic generated by superficial gravitational processes and splash and rill erosion;
9. Deluvio-colluvial glacis and sliding glacis, with moderate stability and moderate morphodynamic, generated by fluvio – torrential processes and landslides, with emphasis risk generated by mechanical stress;
10. Alluvial cones with increased morphodynamic due to poor compaction of deposits;

B. Geomorphological and hydrological processes at risk character. Proposed areas for monitoring - necessary for the control and mitigation

11. Splash and rill erosion;
12. Regressive erosion, ravening and torrential erosion;
13. Gravitational processes (landslides, solifluxions, creep);
14. Processes of compaction induced by mechanical stress from traffic and constructions weight;
15. Riverbeds with alternating fluvial processes (erosion and transport), locally due to anthropic exploitation of sand and gravel;
16. Meandering processes with frequent riverbed mutations due to flow oscillations;
17. Stationary hydrodynamic processes and risk of excess moisture of soil;
18. Areas with high risk to floodings due to overflows or to torrential flows;
19. Processes of infiltration of groundwater, of water from hidden or open river courses, with risk of flooding of buildings basements.

LISTA ILUSTRAȚIILOR

Fig. 1. Harta geomorfodinamică și riscurile induse de procesele geomorfologice și hidrologice

A. Formele de relief și potențialul lor morfodinamic;

1. Suprafețe piemontane cu vulnerabilitate redusă la risc geomorfologic ca urmare a utilizării forestiere;
2. Terasa a 3-a a Cîbinului cu altitudine relativă de 40 – 50 - 70 m;
3. Terasa a 2-a a Cîbinului cu altitudine relativă de 15 – 25 m;
4. Terasa 1-a a Cîbinului cu altitudine relativă de 5 – 10 – 15 m;
a. Poduri de terasă cu stabilitate mare și morfodinamică slab-moderată cauzată de

- pluviodenudare și tasare b. Frunte de terasă cu morfodinamică mare prin procese de ravenare, torențialitate, alunecări;
5. Lunca Cibinului și șesul alluvial al afluenților, cu stabilitate mare și morfodinamică slab – moderată, cu frecvente procese hidrologice staționare și procese morfohidrologice induse de arterele hidrografice;
 6. Culmi interfluviale de podiș cu stabilitate moderată și morfodinamică moderată cauzată de procesele gravitaționale și de pluviodenudare;
 7. Abrupt structural – front de cuestă al Podișului Hârtibaciului cu instabilitate ridicată și morfodinamică accentuată datorită proceselor gravitaționale, torențialității și ravenării, cu risc de reactivare datorat exploatării argilelor;
 8. Suprafață structurală cu stabilitate mare și moderată și morfodinamică moderată generată de procesele gravitaționale superficiale și de pluviodenudare;
 9. Glacis deluvio-coluvial și de alunecare, cu stabilitate moderată și morfodinamică moderată generată de procesele fluvio-torențiale și alunecări de mică profunzime, cu risc de accentuare generat de stresul mecanic;
 10. Conuri de dejecție cu morfodinamică accentuată datorită slabei compactare a depozitelor;

B. Procese geomorfologice și hidrologice cu caracter de risc. Areele propuse pentru monitorizare, se impun măsuri de combatere și diminuare

11. Eroziune în suprafață;
12. Eroziune regresivă, ravenare, torențialitate;
13. Procese gravitaționale (alunecări, solifluxiune, creep);
14. Procese de tasare induse în special de stress mechanic generat de trafic;
15. Procese fluviatile (eroziune și transport), local cu procese de subminare și prăbușire datorate exploatărilor antropice de nisip și pietriș;
16. Procese de meandrare cu frecvente mutații de albie datorate variație scurgerii;
17. Procese hidrologice staționare și risc de supraumezire a terenurilor;
18. Aree cu risc ridicat la inundații datorită revărsărilor și scurgerii torențiale;
19. Procese de infiltrare din pânze freatice, din cursuri de ape libere sau ascunse, cu risc de inundare a subsolurilor.




Fig.1. Geomorfodynamic map and risk induced by geomorphological and hydrological processes

DIODONTIDAE (OSTEICHTHYES) FROM „THE TURNU ROȘU (ROMANIA) EOCENE LIMESTONE” RESERVE

Rodica CIOBANU*

Nicolae TRIF

Abstract. The present study is focused on fossil teeth of the family Diodontidae. The teleostean teeth studied belong to the paleontological collections (Richard Breckner's Collections) of the Natural History Museum from Sibiu and to the private collections (Trif Collections). The identification of the species even at the genus level posed difficulties since the studied teeth are isolated, rather than part of dental apparatus. The main criteria for diagnosis were the morphology of the teeth and the stratigraphical age of rock horizons where the teeth were found. Eleven teeth studied in this paper are referable to *Progymnodon cf. hilgendorfi* (Dames, 1883) and ?*Progymnodon sp.*

Keywords: teeth, Diodontidae, Upper Eocene, Limestone, Turnu Roșu Romania.

Rezumat. Lucrarea de față este focusată pe familia Diodontidae. Dinții de teleosteeni studiați aparțin colecțiilor paleontologice (Colecția Richard Breckner) ale Muzeului de Istorie Naturală din Sibiu și colecțiilor private (Colecția Trif). Identificarea la nivel de gen și specie prezintă dificultăți mai mari la dinții izolați decât la aparatele dentale. Principalul criteriu pentru diagnoză a fost morfologia dinților și vârsta rocilor în care s-au găsit. Cei 11 dinți studiați în lucrare au fost determinați ca apar aparținând la *Progymnodon cf. hilgendorfi* (Dames, 1883) și ?*Progymnodon sp.*

Cuvinte cheie: dinți, Diodontidae, Eocen superior, calcare, Turnu Roșu România.

Introduction

The Eocene fossil fauna from Turnu Roșu (Sibiu) drew the attention of the researcher since the late 18th century. The members of the Transylvanian Society of Natural Sciences, founded in 1849 in Sibiu, were the most active in studying the fossil area Turnu Roșu (Porcești). They pointed out the importance of this habitat not only through the variety of the palaeofauna but also for palaeogeographical reconstructions of the Eocene stratigraphy of the southern Transylvanian Basin (Ciobanu, 2004). In 1850, Ludwig Johann Neugeboren published in Sibiu his first systematic paper on the fossil sharks (teeth), in which the shark species were described for the first time in our country. From 19th century, Neugeboren's paper is the only one, within the current borders of Romania, which describes the shark teeth.

The work appeared shortly after the publication of a fundamental paper for the study of the paleoichthiofauna, which laid the foundation of modern research of fossil fish. I am referring to the work “Recherches sur les poissons fossiles” (1834 - 1844) published by the Swedish researcher Louis

Agassiz. In the preface of the paper Neugeboren underlines the importance of Agassiz's paper to the study he carried on.

If in the 19th century fossil fauna research in Turnu Roșu was at and European level and not only, it is due to the members of the Transylvanian Society of Natural Sciences and Neugeboren was also a member. The majority of the Natural History Museum heritage belonged until November 3rd, 1951 to the Society.

A member of the Society and curator of the paleontological collection was also for a short time Richard Breckner, 7 teeth described in this study belong to his collection.. Considering the importance of the Breckner collection for the Ichthyology – Paleontological study in Romania and not only, I will provide a few brief details regarding Richard Breckner (1900-1979). According to Hienz (1995), Breckner was an art critic, journalist, literary secretary for the theatre and writer. Between 1933- 1938 he was a freelancer making a living by writing scientific papers on the fossil collection from Transylvania. Neither the dictionary nor any other bibliographical source make reference of these

* Brukenthal National Museum,
rodi_ciobanu2005@yahoo.com,
rodica.ciobanu@brukenthalmuseum.ro

about these papers or where the results of his scientific activity were published. Breckner's name appeared more and more often between 1937–1946 in the paper belonging to the Society, in the "Vereinsnachrichten" column, being praised for his work in registering and cataloguing (even re-determining the shark teeth) the collections, especially palaeontological ones. In 1938 Richard Binder, the chairman of the Society, congratulated the diligence and competence of the one who "worked for years on the Porcești tertiary fossils especially the shark teeth". However, there are no details concerning his writing activity or his collection, which we believe, was created during this period (Ciobanu 2007, 2011).

Breckner's palaeontological collection, beside the teleostei fish teeth, contains fossils (molluscs) from Lăpușiu de Sus and 5,000 fish teeth, majority of shark, from Turnu Roșu (Porcești). This collection was acquired in 1954 (according to the museum archives) from Heinrich Breckner (relative of Richard Breckner), a printer from Sibiu.

Material and methods

The paper describes 11 teeth collected from the palaeontological reserve "Calcarele eocene de la Turnu Roșu (Porcești)".

These limestone rich in fauna are part of the Eocene shallow marine sequences lying north of the Făgăraș Mountains, belonging to the southern border of the Transylvanian Basin. Around Turnu Roșu, the Eocene formations emerge like a limestone "patch" area on the north-western ending of the Făgăraș Crystalline (Fig. 1).

The latest concept regarding the stratigraphy of the Turnu Roșu limestone was given by Mészáros (1996) who defined the Turnu Roșu Group, including Valea Nișului and Valea Muntelui formations (Table 1). The faunal analysis underlined the existence of almost all Eocene groups and up to recently they represent the largest deposits bearing isolated fish teeth. Environment reconstructions based on correlation between fossil fauna and recent representatives of the species, are indicating warm tropical-subtropical waters, rich in oxygen and fauna (Mészáros, Ianoliu 1972, 1973; Bucur, Ianoliu 1987; Ciobanu 2006).

Regarding the collecting site of fish fossils remains, we believe that it is Valea Nișului and Valea Caselor, for which there are written references (Fig. 2). Unfortunately, neither Neugeboren, nor other collectors – in our case Breckner – ever mentioned the exact location where they collected the fossils

from. The sample of fish teeth collected in the last few decades is very small compared to the old collections.

In Romania, fossil remains - teeth type «*Diodon*» were described for the Upper Eocene formations from Transylvania.

In the limestone from Cluj, Șuraru, Șuraru (1966) from the outcrops: Pleșca Valley and Baci Quarries they described and illustrated a tooth determined as belonging *Scaroides gatunensis* Toula, 1908. From the Cluj limestone – „upper coarse limestone”, Someș outcrop (area west of Cluj-Napoca), Șuraru *et al.* (1980) described teeth belonging to the species: *Progymnodus cf. hilgendorfi* (Dames), 1883, *Nummopalatus cf. multidentis* (Munster), 1846 and *Scaroides gatunensis* Toula, 1908.

In the year 1987, Șuraru, Șuraru described the Priabonian outcrops from Valea Văratice (Turbuta Formations, Sălaj district) and from outcrops: Pleșca Valley and Baci Quarries (Cluj Napoca area) *Progymnodus cf. hilgendorfi* and *Scaroides gatunensis* species. In the „Cluj limestone”, upper Eocene, Codrea *et al.* (1997) also indicated the presence of the species *Progymnodon cf. hilgendorfi* in a faunal list.

Dica (2003, 2005) made recent observations regarding Diodontidae. In 2003, in the paper where he made the revision of the Diodontidae family from Transylvania, he wrote a review of the bibliographic observations regarding „diodon” teeth. Thus, he shows that Șuraru, Șuraru (1966, 1987) and Șuraru *et al.* (1980) incorrectly assigned the lower jaw to *Scaroides gatunensis* based on Toula's viewpoint and the upper jaw to *Progymnodus cf. hilgendorfi*. In the same paper, Dica specifies based on older material (published) and on new collections that the teeth belong to *Chilomycterus hilgendorfi* and found to four distinct areas, upper Eocene. He pointed out the diagnostic characteristics (for this genus and species) distance between the minute teeth from biting edge and the triturating plates and also the large thin triturating plates. Later, Schulz (2006, p.36, fig.2) cites the species described by Dica (2002, pl.1, figs.1-2) for the *Progymnodon* type.

The first observations on the presence of fish remains (teeth) belonging to the Diodontidae family, in the limestone in Turnu Roșu (the teeth studied in this paper belong to the same outcrop) were made by Șuraru, Șuraru (1966) and assigned to the *Scaroides gatunensis*. The teeth were collected from the Valea Nișului. In the review of the Eocene Diodontids from Transylvania Dica

(2003) pointed out that he collected teeth from the Valea Nişului Formation without indicating the number of items collected and without illustrating them.

Systematic palaeontology

Order Tetraodontiformes Regan, 1929

Suborder Tetraodontoides Berg, 1937

Family Diodontidae Bibron, 1855

Genus *Progymnodon* Dames 1883

1966 *aff. Scaroides gatunensis* Toulou, in Şuraru, Şuraru, p.72-73, 77, figs. 12-14

1980 *Progymnodon cf. hilgendorfi* (Dames, 1883), in Şuraru, Strusievici, Laszlo, p.179, Pl. 1, fig. 4-11.

1987 *Progymnodon cf. hilgendorfi* (Dames, 1883), in Şuraru & Şuraru, p.128-129, Pl.1, fig.1a-b.

1987 *Scaroides gatunensis* Toulou, in Şuraru, Şuraru, p. 129, Pl.1, fig.2a-b.

1997 *Progymnodon cf. hilgendorfi* (Dames, 1883), in Codrea et al., p.39.

2002 *Chilomycterus hilgendorfi* (Dames, 1883), in Dica, p.40-41, Pl.1, fig.1-2

2003 *Chilomycterus hilgendorfi* (Dames, 1883), in Dica, p., Pl.IV, fig.11.

Material: 9 isolated teeth from Breckner collection and 2 recently collected teeth (Trif Collection)

Horizon and locality: Eocene limestone from Turnu Roşu (Porceşti)

Description

Progymnodon hilgendorfi (Dames, 1883)

(Figs.3, 4, 5, 6, 7, 8)

The teeth in figs. 3-8 have rounded to subtriangular anterior border of the package of plates (Fig.3). The connection area of the 2 dental plates is narrow, the two sides of the dental battery are approximately welded. Some of the samples present minute incisors (Figs. 6, 7). The number of triturated plates is difficult to predict because the dental batteries are not preserved wholly. The teeth in figs. 3, 4 and 7 present numerous pores on the occlusal surface, we assume that the enamel from the last triturated plate was not preserved. The teeth in figs. 3, 4 and 5 have the width / length ratio of the tooth battery approximately 2: 1.

?*Progymnodon* sp.

(figs.9, 10, 11, 12)

The teeth from figs. 9, 10, 12 are different from the others in the almost sinusoidal outline of the

trituated plates. In fig. 11 we assume that there are several triturated plates from a dental battery.

Diagnostic features of Diodontid teeth

A diagnostic characteristic of the *Diodon* type dentition is the presence of a dental apparatus made of fused plates, whose triturated surface differs dimensionally and morphologically. The differentiation of various types of the Diodontidae teeth differs from one author to another; they are not very clearly presented and illustrated. Most authors do the differentiation based on criteria pertaining to: the number of plates, how the 2 stacks of plates are welded, the presence or absence of minute teeth in the frontal part, the distance between minute teeth from the biting edges and the triturating plates, and that between the 2 stacks of plates etc.

For example, the difference between *Chilomycterus* and *Diodon* is that in the latter the plates are less extended, a little thicker and more in number. (Dica 2003) Another diagnostic characteristic of this kind is the considerable distance between the minute teeth from the biting edge and the triturated plate.

Regarding the connection between morphology and the positioning of the isolated fossil teeth, as either upper or lower pharyngeal teeth, the authors did not specify clearly the differences.

Another diagnostic element would be, according to some palaeontologists, the presence or absence of teeth on the biting edges. Tyler (1980) (quoted by Dica 2002) believes that this characteristic is not relevant in the diagnosis because the conditions of fossilization, the degree of conservation of the material as well as wear can lead to the loss of marginal teeth,

Tyler (1980) considers that the difference between the *Progymnodon* and *Oligodiodon* represents an evolution in the structure of the dental apparatus. Thus, in the *Oligodiodon* genus, as in the recent genus, the biting edge and the triturating region are separate by a large layer.

Weems (1998) considers that the diagnostic characters for the *Progymnodon* are: “the groove separating the teeth of the trituration plate from the marginal jaw teeth is narrow, and the marginal jaw teeth are relatively large and stout”. In addition, he observed that the width/length ratio of the tooth battery (approximately 2: 1) is quite comparable to the proportions of *Progymnodon hilgendorfi* (Dames, 1883) and defined them as belonging to the same genus.

Schultz (2006, p. 34) made a complex analysis of fossil representatives of the Diodontidae family in the study of the *Oligodiodon* from the Miocene (Badenian) of Styria (Austria). Thus, *Progymnodon*, known only in the Eocene has the dental apparatus made of 2 stacks of triturated plates separated by a wide space. *Oligodiodon* and *Chilomycterus* have fewer triturated plates. The number of triturated plates increases greatly reaching nearly 20 in the *Diodon* known since the Miocene. Regarding the *Scaroides gatulensis* he considers that it belongs to the Diodontidae family and the holotype indicates a synonymy with the *Chilomycterus*. Thus, *Progymnodon* whose triturated plates have a wide median joining area, is indicated only for Eocene and for the *Chilomycterus*, whose triturated plates are united without any space between them, evolved from the Miocene and still exist in the present. Schultz (p. 35) showed that the genera *Progymnodon*, *Oligodiodon*, *Chilomycterus* have great morphological similarities.

Discussions

Few genera of Diodontidae form a compact family of Gymnodontidae, which have changed very little since the Eocene, when they appeared, except a decrease in size, and a slight rearrangement of the tooth dental structure.

The fossil Diodontidae are known through the dental remains that appear frequently in marine deposits and rarely through skeletons (complete skeletons have been found in Eocene from Monte-Bolca, Italy). Quite common in the Tertiary formations, dental remains appear as dental apparatus composed of triturated plates united in the form of a battery. Most of the time this type of fossil teeth is generically referred to “*Diodon*”.

The nomenclature of fossil species and genera is disputed, and the criteria for distinction are unclear. Differentiations between species and genera are not major and can be done only when the fossil material is very rich and varied as species and also from diverse localities. The nomenclature of current species is not clarified as well and there are many synonymies.

Leis (2006) revised the nomenclature and distribution of the Diodontidae and specified that out of 70 nominal species (over 60% of which were described in the 100 years following Linnaeus) only 19 species, belonging to 7 or 8 genera, are valid. There are different opinions regarding the age of the first fossil remains of the Diodontidae family. Most authors consider that the Eocene represents the emergence of the family.

But Weems (1998) brings to attention, for the first time, dental remains – *Progymnodon hilgendorfi* – in formations of Paleocene age. The first reported lower Eocene representative of the Diodontidae family is from India (Kumar and Loyal, 1987). The type of *P.hilgendorfi* is from Middle Eocene of Egypt (Priem 1914).

Based on dental structure, Tavani (1955) differentiated some new genera. Following the study of a rich fossil material, he differentiated different genera of the Diodontidae family by bone mass, present or absence of the biting edge, between teeth and triturated plates. He also considered that *Eodiodon* which does not show small teeth on the biting edges belongs to another family, namely *Eodiodontidae*. Tavani (1955) differentiated 4 genera: *Progymnodon* from Middle and Upper Eocene of Europe, *Kyrtogymnodon* from Pliocene of Europe, *Oligodiodon* from Oligocene and Miocene of Europe and North America and *Diodon* from Miocene of Europe and in modern seas. The evolution from *Progymnodon* to *Oligodiodon* highlights, in the opinion of the author, the change in dental structure, where the biting edge and the triturated teeth are well separated by a thick layer of bone, as in current forms.

Concerning the stratigraphic distribution of Diodontid genera, Schultz (2006, 33) noted that *Progymnodon* occurs only in Eocene, *Oligodiodon* lived from Oligocene to Miocene, and *Chilomycterus* and *Diodon* from Miocene up to the present.

Conclusions

The diodontid fossils described from the Eocene of Romania by Dica (2003) should be referred to *Progymnodon* and not to the *Chilomycterus*.

In regard to the living environment, recent Diodontidae are marine and stenohaline in slightly deeper waters. They feed on invertebrates such as echinoderms, molluscs and crustaceans (Leis, 1984), food and environment that has been reconstructed for the coastal area of the Eocene sea in Turnu Roşu as well (Ciobanu 2006). In terms of biogeographical characteristics, the fish fauna from Turnu Roşu and throughout Transylvania presents tropical indo-pacific features (Şuraru, Şuraru 1966; Ciobanu 2006).

Acknowledgements. I would like to thank Prof. PhD. Kishor Kumar (Wadia Institute of Himalayan Geology, Dehradun, India) for improvements on text, for his useful advice and some stylistic suggestions to improve the manuscript. The

authors are grateful to Ana Maria Mesaroş for constructive comment on our manuscript and to

Ioana Ciobanu for the useful suggestions improving the language for the text.

REFERENCES

- Bucur, Ianoliu 1987 Bucur Ioan, Ianoliu Constantin, *L' Éocène de Turnu Roşu – Porceşti. Contributions sur les algues calcaires*. In: *The Eocene from the Transylvanian Basin*, Cluj Napoca (1987), p. 37-42.
- Ciobanu 2004 Ciobanu Rodica, *Ariile paleontologice protejate ale judeţului Sibiu*. In: *Geocarpatica* IV/4, Sibiu (2004), p. 155-167.
- Ciobanu 2006 Ciobanu Rodica, *Evaluări paleoecologice pe baza patrimoniului muzeal. Studiu de caz: Turnu Roşu*. In: *Marisia. Studii şi materiale* XXVIII, Tg.Mureş (2006), p. 69-79.
- Ciobanu 2007 Ciobanu Rodica, *Naturalists from Sibiu and their fossil collections at the Natural History Museum from Sibiu*. In: *Acta Musei Devensis, Sargetia, Series Scientia naturae* 20, Deva (2007), p. 27-42.
- Codrea et. al. 1997 Codrea Vlad, Vremir Matei, Dica Paul., *Calcarul de Cluj de la Someş-Dig (Cluj-Napoca): Semnificaţii paleoambientale şi impactul activităţilor antropice asupra aflorimentului*. In *Studii şi cercetări* 3, Bistriţa (1997), p. 31-39.
- Dica 2003 Dica Paul, *A review of the Eocene Diodontids and labrids from Transylvania*. In: *Studia Universitatis Babeş-Bolyai, seria Geologie* XLVII/2, Cluj Napoca (2003), p. 37-47.
- Dica 2005 Dica Paul, *Studii paleoambientale in formatiunile paleogene din NV-ul Depresiunii Transilvaniei pe baza asociaţiilor ihtiologice. Raport de cercetare, grant TD 24/15*. In: *Revista de Politică Ştiinţei şi Scientometrie*, Numar Special (2005), p.1-77.
- Kumar et al. 1987 Kumar Kishor, Loyal Raminder, *Eocene ichthyofauna from the Subathu Formation, Nort-Western, Himalaya*. In: *India Journal of the Palaeontological Society of India* 32, (1987), p. 60-84, 3 pls., 102 figs.
- Hienz 1995 Hienz Hermann, *Schriftsteller – Lexicon der Siebenbürger Deutschen Bio – bibliographisches Handbuch für Wissenschaft, Dichtung und Publizistik*, Begründet von Joseph Trausch vorgeführt von Friedrich Schuller und Hermann Hienz, Bd.I (A-C), Köln, Weimar Wien, Böhlau Verlag (1995), 624 p.
- Leis 1984 Leis Jeffrey, *Family Diodontidae in: Fisher, W. (ed.) FAO species identification sheets for fisheries purpose: the Eastern Indian Ocean*. In: *FAO*, Rome (1984), 8 pp. (unpagined)
- Leis 2006 Leis Jeffrey, *Nomenclature and distribution of the species of the porcupine fish family Diodontidae (Pisces, Teleostei)*. In: *Memoires of Museum Victoria* 63/1, (2006), p. 77-90. (<http://www.museum.vic.au/memoirs/index.asp>).
- Mészáros Ianoloiu, 1972 Mészáros Nicolae, Ianoliu Constantin, *Macrofauna eocenă de la Turnu Roşu (I)*. In: *Studii şi comunicări, Muzeul Brukenthal, Ştiinţele Naturii* 17, Sibiu (1972), p. 21-31.
- Mészáros Ianoloiu, 1973 Mészáros Nicolae, Ianoliu Constantin, *Macrofauna eocenă de la Turnu Roşu (II)*. In: *Studii şi comunicări, Muzeul Brukenthal, Ştiinţele Naturii* 18, Sibiu (1973), p. 13-23.
- Mészáros 1996 Mészáros Nicolae, *Stratigrafia regiunii Turnu Roşu – Porceşti*. In: *Convergenţe transilvane* 4, Sibiu (1996), p.43.
- Priem 1914 Priem F., *Sur des Vertébrés du Crétacé et de l'Éocène d'Égypte*. In: *Bulletin de la Socité Géologique de France* 4 série (14), Paris (1914), p.366-382.
- Schultz 2006 Schultz Ortwin, 2006, *Oligiododon, a porcupinefish from the Middle Miocene (Badenian) of Styria, Austria (Diodontidae, Osteichthyes)*, In: *Joannea geol. Paläont.* 8, (2006), p.25-46.
- Şuraru, Şuraru, 1966 Schuraru Nicolae, Şuraru Maria, 1966, *Asupra unor resturi de peşti eocenici din Bazinul Transilvaniei*, In: *Studia Universitatis Babeş-Bolyai, seria geologie-Geografie* fasc.1 Cluj-Napoca (1966), p.69-77
- Şuraru et al. 1980 Şuraru Maria, Strusievici Robert, László Kovaci. *Resturile unor dinţi de teleostei în Eocenul de la Cluj-Napoca*, In: *Studii şi Cercetări de Geologie şi Geofizică* 25, Bucureşti (1980), p.177-181.
- Şuraru, Şuraru 1987 Şuraru Nicolae, Şuraru Maria, 1987, *Neue angaben über einige Fischreste aus dem Eozän des Transylvanien Beckens*, In: *The Eocene from Transylvanian Basin* Cluj-Napoca, p.127-134.
- Tătărâm 1970 Tătărâm Niţa, *Eocenul din sud-vestul Transilvaniei consideraţii biostratigrafice şi paleogeografice*. In: *Analele Universităţii Bucureşti, seria geologie* 19, Sibiu (1970), p.

- 119-141.
- Tavani 1955 Tavani G., *Observazioni su alcuni plectognathi (Gymnodonti)*, In: *Atti della Soc. Toscana Scienzi Naturali*, mem. (A)62, Pisa (1955), p.176-200.
- Tyler 1980 Tyler C. James, *Osteology; Phylogeny and Higher Classification of the fishes of the Order Plectognathi (Tetraodontiformes)*. In: *NOAA Technical report NMFS 434*, Washington (1980, 422 pp.
- Weems 1998 Weems Robert, *Actinopterygian fish remains from the Paleocene of South Carolina*, In: A.E.Sanders (ed.) *Paleobiology of the Williamsburg formation (Black Mingo Group Paleocene) of South Carolina, USA, Transactions of the American Philosophical Society* 88/4, Philadelphia (1998), p.122-146.

LIST OF ILLUSTRATIONS

- Tab. 1.** Stratigraphic succession in the Turnu Roşu Area - Turnu Roşu (Porceşti) Group
- Fig. 1.** Geological map of Turnu Roşu palaeontological reserve (adapted after Tătărâm, 1970)
- Fig. 2.** Eocene Limestone from Turnu Roşu (V. Nişului)
- Fig. 3-8.** *Progymnodon hilgendorfi* (Dames, 1883)
- Fig. 9-11.** *?Progymnodon* sp.

LISTA ILUSTRĂȚIILOR

- Tab. 1.** Succesiunea stratigrafică în arealul Turnu Roşu -a Grupului Turnu Roşu (Porceşti)
- Fig. 1.** Harta geologică a Rezervației paleontologice Turnu Roşu (adptare după Tătărâm, 1970)
- Fig. 2.** Calcarele eocene de la Turnu Roşu (V. Nişului)
- Fig. 3-8.** *Progymnodon hilgendorfi* (Dames, 1883)
- Fig. 9-11.** *?Progymnodon* sp.

Tab. 1. Stratigraphic succession in the Turnu Roşu Area - Turnu Roşu (Porceşti) Group
(Mészáros, 1996, present work)

Valea Nişului Formation	Oligocen basal	Sandstone with fish scale
	Priabonian	Sandstone, calcareous sandstone, limestone with molluscs, selachian and bony fish teeth, crocodilian teeth, sirenian vertebrae, fossilized wood
		Conglomerate with <i>Nummulitesfabianii</i> and <i>Chlamys biarritzensis</i>
Strada Muntelui Formation	Lutetian- Priabonian	Calcare, calcare marnoase cu resturi de selacieni
		Limestone, Marl-limestone with <i>N. millecaput</i> and <i>Campanile</i> , fish teeth, sirenians vertebra
		Limestone with <i>Nummulites laevigatus</i> , <i>Nummulites distans</i> and molluscs
Valea Satului Formation	Cuisian	Microconglomerate
		Calcareous sandstone with molluscs and? shark teeth
		Conglomerate with <i>Ampullinopsis procensis</i>
		Sandstone, conglomerate and marl with <i>Nummulites planulatus</i>
		Clay with foraminifera




Fig. 1. Geological map of Turnu Roşu palaeontological reserve reserve
(adanted after Tătărâm 1970)



Fig. 2. Eocene Limestone from Turnu Roşu (V. Nişului)



Fig. 3. *Progymnodon hilgendorfi* (Dames, 1883)



Fig. 4. *Progymnodon hilgendorfi* (Dames, 1883)




Fig.5. *Progymnodon hilgendorfi* (Dames, 1883)



Fig.6. *Progymnodon hilgendorfi* (Dames, 1883)



Fig.7. *Progygnodon hilgendorfi* (Dames, 1883)



Fig.8. *Progygnodon hilgendorfi* (Dames, 1883)



Fig.9. ?*Progymnodon* sp.



Fig.10. ?*Progymnodon* sp.

SCIENTIST OF HIS TIME – JOHANN MICHAEL ACKNER (1782-1862)

Rodica CIOBANU*

Ackner's complex personality caught the attention of several researchers, both over the past centuries, as well as nowadays (Wollmann 1982, 1983; Schneider & von Killyen 2007; Ciobanu, 2005-2007). The goal of this short communication is to draw attention towards this remarkable Transylvanian personality from Sibiu, as in 2012 we celebrate 230 years from his birth but also we commemorate 150 years since his death.

Ackner played an important role in the evolution of the Transylvanian Society for Natural Sciences from Sibiu in the field of mineralogy and palaeontology. He was an important personality of the Society, distinguishing himself in various fields: education, mineralogy, paleontology, archaeology. He was loved and respected by his contemporaries as a “mentor and teacher of natural sciences and Transylvania's antiquities”. Due to his vast knowledge – according to the ideals of the 18th century Age of Enlightenment, this implied that all the knowledge of the period should be understood - Johann Michael Ackner was a universally appreciated scholar.

Ackner was born on 25th January 1782 in Sighișoara. His father was Georg Ackner, a middle school teacher and later a priest in Meeburger. He attended the gymnasium of Sighișoara and Sibiu. At the age of 21, he left Transylvania to attend the university in Wittenburg.

After finishing his studies in Germany and after making several voyages to Renania, Italy and France, he came back to Sibiu with a great passion for archaeology and geology.

Returning to Transylvania, he did not settle, as was then customary, in his hometown Sighișoara. Instead he accepted a job as professor of philology and archaeology at the Evangelical Gymnasium in Sibiu. During 13 years of activity at the gymnasium, he won the students' attention through his graphic and scientific way of teaching. Conditioned by the lack of teachers J Michael Ackner continued to teach at the gymnasium, even when he was promoted as preacher (archdeacon) in

Sibiu. In 1821, he was chosen by the community from Hammersdorf (today Gușterița) as priest.

There he continued the local chronicle already started in 1709. This contained, among others, short news about bad weather and crop results, notes about the country life adapted to the church calendar, information on the maintenance of the church and the parish, exceptional bad weather, floods and earthquakes are also mentioned, as well as epidemic outbreaks or a locust invasion.

In parallel with his ecclesiastical duties J. Michael Ackner dedicated himself to his various studies related to natural sciences. Gușterița village, situated in the south – east of the city of Sibiu, today a neighbourhood of the city, placed at the bottom of the Pădurea or Gușterița Hill, became famous among palaeontologists due to the fossils discovered here by Ackner, especially the quaternary vertebrates.

If we observe the complete work of the scientist, who was active until the end of his life, three different fields, in which Ackner was active, stand out. After a first stage that represents the study of physics and astronomy a second stage follows, which was devoted mostly to mineralogical, geological and paleontological studies. These were the 30s and 40s of the 19th century. In parallel there were also archaeological studies, which grew in importance in the 50s and were the highlight of his scientific activity until the end of his life. In this paper we will deal with his activity in the field of geology, linked to his activity as collector of minerals and fossils, and especially as member of the Society of Natural Sciences, which founded Natural History Museum – the institution that today houses Ackners's collection.

“The travel notes about a part of the southern Carpathians, that separates Transylvania from Lesser Wallachia” (*Reisebericht über einen Theil der südlichen Karpaten, welche Siebenbürgen von der Kleinen Walachei trennen*) (1838), which later, in 1852, appeared as “Geological-paleontological reports of the Transylvanian border mountains along Small Wallachia” (*Geologisch – paläontologisches Verhältnis des Siebenbürgisches*

* Brukenthal National Museum,
rodi_ciobanu2005@yahoo.com,
rodica.ciobanu@brukenthalmuseum.ro

Gränzgebirges längs der Kleinen Walachei) prove his intense activity in the geology field activity in the field of geology. The report includes an appendix with “a summary on the fossil remains and indications about the formations and mineral deposits found in the mountains along the border of Transylvania and Wallachia”. Of great importance and a gain in knowledge from a mineralogical-geological point of view is also the mentioned journey to the Western Carpathians up to Rodna and through the Transylvanian Plateau. His publications were praised by the scientists of his time.

However, his most appreciated geological studies referred to the surroundings of Sibiu. He explored the cretaceous limestone rock and the schistose sandstone slate from Silverbachtal in Cisnădioara, the area around “Halben Stein”, now protected by law. Ackner's paper “On the formations of sandstone nodules and their spread in the Transylvanian sandstones” (*Über die Sandstein – Kugelbildungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften*) published in *Verhandlungen und Mitteilungen des Siebenbürgischen Vereins für Naturwissenschaften* (1853) is still considered to be a fundamental study. In parallel with his mineralogical-geological studies he was engaged in paleontological studies. Some of these are included in “*Bericht über die zu Holzmengenim Harbachtale gefundenen vorweltlichen Tierreste und verschiedene ähnliche Funde aus Hammersdorf*”, which appeared in 1852 in *Verhandlungen*. The Eocene deposits of shark teeth and molluscs shells from Porcești were also researched but Neugeboren is the one who studied and revaluated them scientifically.

The priest and royal counsellor, Ackner, was known especially through his collection and archaeological researches and was a member of the two Saxon cultural Societies of Sibiu: *Verein für Siebenbürgische Landeskunde* and *Siebenbürgische Naturwissenschaften zu Hermannstadt* (Neugeboren, 1866). Besides being the founder of the latter, he was one of its active members and an assiduous collector. This is proven by the often mentioning of his name in “*Vereinsnachrichten*” connected to the collecting activity of fossils for the Museum.

The collecting activity, started in the 20th decades of the 19th century and represented the materialization of his studies and researches undergone in the field of geology (mineralogy, paleontology). An important paper with paleontological subject is *Contribuția la geognosia*

și paleontologia fosilelor pietrificate (1845) and for the first paper dealing with the ores from Transylvania *Mineralogia Transilvaniei cu observații geognomice* (1855) Ackner received the award of “The Association for Transylvanian Geography” („Asociația pentru Geografia Transilvaniei”).

The papers published in this field – paleontology - did not remain without any echoes. His contemporary L.J. Neugeboren, a member of the same Society having similar scientifically interests, appreciate Ackner's scientific attempts (Neugeboren 1852). The often trips to Cisnădie (Heltau), Cisnădioara (Mischelsberg), around Cluj (Klausenburg) and across the Carpathians (Wollmann 1982) represented occasions for Ackner to collect fossils. One can thus explain the fauna diversity of his collection. Compared to Neugeboren's collection, Ackner's is far richer considering the fossil type specimens and the collecting point such as: Agnita, Bruuiu, Săcădate, Ilimbav, Daia.

The acquisition of the Ackner collection from Hermann Ackner, his eldest son, for the sum of 4,500 florins, was recorded in *Vereinsnachrichten* (1866) in 1866. C. Zekelius, caretaker of the paleontological collection of the Museum mentioned upon taking over the collection that it has a “strong Transylvanian character” and that the items of the Ackner collection were personally collected and obtained from other collectors through exchange or acquisition.

The catalogue of the Ackner collection, already renowned among the Transylvanian naturalists is published in the first issue of the Society journal (1850). By studying the catalogue one may notice that it contained fossil items from almost all of the systematic groups: Plants (Daia, Săcădate, Cornățel, Glâmbocă, Băile Lăpuș), Sponges and corals (Săcele, Zărnești, Brașov, Turnu Roșu, Dobârca), Echinoderms (Turnu Roșu, Cisnădioara, Cluj), Cephalopodes (Cisnădioara, Săcel, Brașov), brachiopods (Brașov, Turnu Roșu), Crustaceans (Turnu Roșu), Bivalves (Turnu Roșu, Dobârca)

In a paper referring to the mineralogical and paleontological collections from Transylvania, Neugeboren (1866), eulogistically talked about the Ackner collection and pointed out that it was bought with the support of the “National Saxon House and of the Societies: “the Transylvanian Society for the Study of Nature from Sibiu” (*Siebenbürgische Verein für Naturwissenschaften zu Hermannstadt*) and “the Society for the Study of Transylvania” (*Verein für Siebenbürgische*

Landeskunde) (Verhandlungen, 1867). In 1867, the duplicates from the Ackner collection were given to the Evangelic School from Sibiu (Verhandlungen, 1867, 1). The collection was enlarged at a short time after the acquisition.

Thus, Neugeboren, as caretaker of the paleontological collections, thanks in a meeting of the Society to Czekelius and Moekesch for the enlargement of the Ackner collection with fossils from the surroundings of Cluj and Alba Iulia (Verhandlungen, 1867:238). Neugeboren mentioned in 1866 that in the Ackner collection there were 3,791 items out of which 1,728 were fossils (geognostical items) in which the following were represented: the "diluvia from Gușterița, the petrified plants from Thalheim – Daia (tertiary), Sebeșul de Sus, Porcești (lower tertiary), Cîsnădioara (chalk formation and Gosau), Rășinari". The collection also included fossils belonging to the 1,315 species, the most valuable being those from Gușterița (mammal bones), from the Hârtibaciu Valley (plants, fossil fish, mammal bones), the Brașov area (certain corals), Săcădate (plants, tertiary fish), Turnu Roșu (Porcești), Cîsnădioara, Buituri, Racoș, Muncelul Mic, Lăpușul de Sus, Cluj (Neugeboren, 1866).

The fossils discovered by Ackner in Gusterita Hill, while he was still a priest, drew the attention of other paleontologists towards this area. At the end of the 19th and the beginning of the 20th century the studies dedicated to the geological potential of Gușterița mentioned the fossil specimens found in the Ackner collection. The fossil area from Turnu Roșu is known, especially, due to the works and collection of Neugeboren, while Gușterița is renowned because of Ackner's discoveries and collections. The naturalist Ackner, unlike Neugeboren, did not work, outside volunteering as member of the Society, inside the museum, and obtained special performance in the field of archaeology. Neugeboren carried out a few studies related exclusively to the geological aspects and in his ad notations did not lose the opportunity to related to incorrect determinations, point to the fact that time was not on his side in going deep into this field. However, Neugeboren did not lose the opportunity to praise Ackner's collecting activity, mentioning that through his collection several fossil groups were completed and dedicated him a species of bivalve *Cardium acknerii* (Neugeboren 1851)

From the Vereinsnachrichten column records of the Verhandlungen journal one might notice the

continuous increase of the Ackner collection after its acquisition by the museum. It was only in 1891, when Sachsenheim, the caretaker of the museum's collection during that period, worked on recording the collection. (Verhandlungen, 1891, XV). Unfortunately, this register was not kept and an inventory book made by Breckner at the beginning of the 20th century and kept at the Natural History Museum, does not point out the data referring to the Ackner collection but only the existence of isolated pieces donated by Ackner. The entire collection, as museum entity, used to exist in the museum before 1955. We conclude this from the fact that Ilie (1955) makes references to the Ackner collection when he presents the Pontian and the Pleistocene from Gușterița, in the geological research of the Alba-Iulia-Sibiu-Făgăraș-Rupea area, namely in the stratigraphic descriptions. Currently, the Ackner collection does no longer exist in its original form, but only as isolated pieces among the "*Old collection of the museum*" also called "the Society Collection".

Johann Michael Ackner's scientific achievements enjoyed all the recognition of his contemporaries. Thus, in April 1850 he became corresponding member of the Institute of Archeology of Rome. A year later Ackner was named a member of the Leopold-Caroline Academy of Naturalists from Breslau. He also became a corresponding member of the Society of Scholar from Cracow, as well as of the Central Committee for Research and Maintenance of Architectural Monuments and correspondent of the Geological Imperial Royal Institute of Vienna. As a sign of great recognition in 1854 he was honoured with the Gold Cross of Merit and in 1858 appointed to the Imperial Council. He was a member in the leading committee in both associations for regional studies and natural sciences. Ackner was also a honorary member of "Astra", but unfortunately he was unable to receive his diploma.

Although the collection does no longer exist as Ackner left it, important items from this collection are presented and scientifically rendered. Had there been drawings of the fossil items, the reconstitution of the entire collection would have been possible.

Acknowledgements. The authors are grateful to Ana-Maria Mesaroș for constructive comment on our manuscript and to Ioana Ciobanu for the useful suggestions improving the language for the text.

REFERENCES

- Ciobanu, 2005 - Ciobanu Rodica, *Naturalists from Sibiu and their fossil collections at the Natural History Museum from Sibiu*. In: *Sargetia, Acta Mus. Dev. Ser. Sci. Nat.* XX, Deva (2005 – 2007), p. 27 – 41.
- Ilie 1955 Ilie Mircea, *Cercetări geologice în Bazinul Transilvaniei (Regiunea Alba Iulia – Sibiu Făgăraș – Rupea)*. In: *Anuarul Comitetului Geologic* 37, București (1955), p.335.
- Neugeboren, 1851 Neugeboren Johann Ludwig, *Zur vortweltlichen Conchiliogie Siebenbürgens*. In: *Verhandlungen und Mitteilungen Siebenbürgische Verein für Naturwissenschaftlichen zu Hermannstadt* 2, Sibiu (1851), p. 4-9.
- Neugeboren1852 Neugeboren Johann Ludwig, *Literärische Notiz über M.J.Ackner's Monographie „Geologisch paläontologisches Verhältniss des siebenbürgischen Gränzgebirges längs der kleinen Walachei“ im Archiv des vereins für siebenbürgens Landeskunde*, In: *Verhandlungen und Mitteilungen Siebenbürgische Verein für Naturwissenschaftlichen zu Hermannstadt* 2, Sibiu (1851), p. 25-30.
- Neugeboren1866 Neugeboren Johann Ludwig, *Notizen über Sammlungen siebenbürgischer Mineralien*. In: *Archiv des Vereins für siebenbürgische Landeskunde, neue folge* 7 (1-2), Sibiu (1866), p. 389-390.
- Schneider, von Killyen 2007 Schneider Erika, Killyen v. Hansgeorg, *Johann Michael Ackner (1782-1862) Universalgelehrter seiner Zeit*. In: *Naturforscher in hermannstadt. Vorläufer, Gründer & Förderer des Siebenbürgischen Vereins für Naturwissenschaften*, Schneider Erika, Killyen v. Hansgeorg, Scheneider Eckbert (eds.), Sibiu (2007), p.31-37.
- Wollmann 1982 Wollmann Volker, *Johann Michael Ackner (1782-1862). Leben und Werk*, Dacia Verlag, Cluj-Napoca (1982), p. 19-30, 51-78.
- Wollmann 1983 Wollmann Volker, *Briefe zur Geschichte der siebenbürgischen Altertumskunde*, Kriterion Verlag, București (1983), p. 281.



Fig.1. Johann Michael Ackner (1782-1862) – with a mammoth femur collected from Gușterița (Sibiu), currently exhibited in Natural History Museum from Sibiu

Rodica CIOBANU*

The geomorphologic analysis implies a complex study of the relief, the genesis and evolution of the landforms in their complexity, of the typology of geomorphologic processes that take place and of the impact that these geomorphologic processes have on the other components of the environment.

The results of any geomorphologic analysis are materialized in graphic and textual representations, which become geomorphologic maps that are realized through a complex geomorphologic mapping process.

In the present day context, when the technological development and the volume of information has grown substantially in all economic and scientific fields, the analysis of the relief, as main component of landscape, relies more and more on computer technology, with possibilities of fast processing, and on modern methods of complex analysis accomplished with the help of specialized software, which become more advanced and cheaper by the day.

The “Geomorphologic mapping and analysis” work was drawn up for the purpose of providing synthesized and systematic documentary teaching aids, which are necessary for the mastering of geomorphologic mapping, as well as for the assimilation of relief analysis knowledge.

The content of this volume dwells on general notions concerning the relief research methodology, the principles of graphic representation and geomorphologic mapping, as well as the content and structure of topographical and geological maps, used as basis for geomorphologic mapping and geomorphologic maps as end product.

The first part of the paper introduces several documented and scientifically augmented research methods used in geomorphology: general methods (inspection, comparison, analysis, prognosis) and specific ones (morphometric and morphographic

methods). This part also describes graphic representation methods of statistic data for describing, comparing, causal and functional explaining of the different morphologic surfaces, morphologic indicators or geomorphologic phenomena, as well as cartographic representation methods for landforms.

The paper especially addresses geography students, but also those who study ecology or related specializations. In order to assure a faster documentation for students from other specializations of the environmental science field or for different experts who studied topography and geology, the author resorted to a detailed presentation of the basic elements used in geomorphologic mapping.

Particular emphasis is given to the use of topographic maps as mathematical support in creating geomorphologic maps, to highlighting the characteristics of the standardized and adopted cartographical projections for topographic maps, to the presentation of the sharing system on sheets and the creation of the map sheets’ nomenclature, to the study of the content elements and to the meaning and ways of representation of the conventional signs.

The work also addresses the role of geological substrate in morphogenesis and presents the mapping principles and methods of the petrographic and structural relief, the analysis of the forms and how to interpret maps.

The geological analysis in interpreting structural tectonic and lithological forms (the geochronological and morphochronological scale, specific research methods applicable in geomorphology), types of rocks and their role in relief shaping, the petrographic and structural relief characteristics, the creation stages for morphostructural and morpholithological maps are all scientifically documented by the author and comply with the requirements of higher education. The difficulty of creating and interpreting them is supported by concrete examples.

*Brukenthal National Museum,
rodi_ciobanu2005@yahoo.com,
rodica.ciobanu@brukenthalmuseum.ro

Special attention is given to morphometrical and morphometrical maps, by presenting the methods of calculation and representation of the relief fragmentation density and of the energy of the relief, the methods for determining the terrain's declivity and for creating the map of the cliffs and of the slopes' exhibition, the principle and methods of compiling a morphographic map. The theoretical and methodological content is completed and argued with a wide range of case studies and mapping examples accompanied by competent explanations.

Synthetically representations are an important part of the work: the general morphologic map, the actual map of the geomorphologic processes and the geomorphologic risk map. The creation and interpretation of these maps requires a particular experience in cartographic representation and relief analysis, which is the reason for which the author presents map patterns and case studies.

The last part covers the study of modern digital mapping technology, summarizing principles, structure and methods of creating a geographical database and how to work with the program Map

Maker and Surfer, in order to create digital geomorphologic maps. The presented topic of digital mapping, besides the scientific approach, has also a practical-applicability role for the bachelor and master students in their development of practical skills regarding the effective use of the mapping software when creating databases and maps.

The work is characterized by a lofty scientific level and by an appropriate approach towards the challenges of mapping and geomorphologic analysis. The basis for preparing the paper was the author's experience, alongside a rich bibliographical material.

Environmental science bachelor and master students, as well as those interested in the mastery and understanding of the mapping and geomorphologic analysis fields can find in each chapter not only the theoretical presentation of the problem, but also practical applicability of cartographic representations and the applicative context of the geomorphologic analysis.

THE MIKLOS SZEKERES DONATION TO THE MALACOLOGICAL COLLECTION FROM THE NATURAL HISTORY MUSEUM IN SIBIU

Ana-Maria MESAROS*

Abstract. In 2012 an important donation was made by Miklos Szekeres to the Natural History Museum from Sibiu (Brukenthal National Museum). Szekeres donated to the Malacological Collection 9 paratypes belonging to 6 subspecies of the genus *Alopi*a (H. et A. Adams). The museum received copies of the papers published by Szekeres including the original descriptions for each subspecies. The paratypes were collected from Romania between 1998 and 2008. To each specimen an inventory numbers was given and included in the museum Malacological collections.

Keywords: paratypes, donation, genus *Alopi*a

Rezumat. În 2012 o importantă donație a fost făcută de către Miklos Szekeres Muzeului de Istorie Naturală din Sibiu (Muzeul National Brukenthal). Szekeres a donat colecției de Malacologie a muzeului 9 paratipi aparținând la 6 subspecii ale genului *Alopi*a (H. et A. Adams). Muzeul a primit o dată cu donația și descrierea originală a specimenelor publicate de către Szekeres. Paratipii au fost colectați din România între anii 1998 – 2008. Exemplarele au primit numere de inventar și au fost incluse în colecțiile malacologice ale muzeului.

Cuvinte cheie: paratipi, donație, genul *Alopi*a

Introduction

Miklos Szekeres, scientific advisor at the Biological Center of the Hungarian Academy of Sciences, donated in March 2012 to the Natural History Museum (part of the National Brukenthal Museum) from Sibiu 9 paratypes belonging to 6 subspecies of the genus *Alopi*a (H. et A. Adams). The specimens were collected by scientific collaborators of Szekeres from Romania.

Each specimen entering the museum collection was measured, photographed and included in the DOCPAT 2000 museum data base with all the information sent by the donor: the description, collecting area and date, the names of the collectors. According to the cultural heritage categories in Romania section natural heritage these specimens can be classified at thesaurus (the highest category).

The paratypes donated by Szekeres, following the accounting note number 1 from 03.04.2012, are:

- inventory number 209.708, *Alopi*a *glorifica sarkanyi* Szekeres, collected from Mateiaș Mountains, East - Northeast of Câmpulung, 1220 meters altitude, collected by L. Németh, 10.09.2003;
- inventory number 209.709, *Alopi*a *glorifica valeriae* Szekeres, collected from Dâmbovița

- Valley, 3 kilometers Northwest of Cetățeni, 550 meters altitude, collected by L. Németh, 21.07.2006;
- inventory number 209.710, *Alopi*a *glorifica valeriae* Szekeres, collected from Dâmbovița Valley, 3 kilometers Northwest of Cetățeni, 550 meters altitude, collected by L. Németh, 21.07.2006;
- inventory number 209.711, *Alopi*a *subcosticollis majorosi* Szekeres, collected from Holocene cave sediment near the entrance of Peștera Polovragi, 670 meters, collected by G. Majoros and L. Németh, 22. 07.1998;
- inventory number 209.712, *Alopi*a *subcosticollis tamasorum* Szekeres, collected from Cerna Valley, Bobotu Gorge North of Herculan Baths, 380 meters, collected by T. Deli, T. Domokos, J. Lennert and M. Szekeres, 14.08.2006;
- inventory number 209.713, *Alopi*a *glorifica deceptans* Deli & Szekeres, collected from Piatra Craiului Mountains, Seacă Pietrelor Valley, 2 kilometers North of Brusturet Lodge, 1150 meters altitude, collected by L. Németh, 2-3.09.2005;
- inventory number 209.714, *Alopi*a *glorifica deceptans* Deli & Szekeres, collected from Piatra Craiului Mountains, Seacă Pietrelor Valley, 2 kilometers North of Brusturet Lodge, 1150 meters altitude, collected by L. Németh, 2-3.09.2005;
- inventory number 209.715 *Alopi*a *grossuana nemethi* Deli & Szekeres, collected from Vâlcău Mountains, Sohodol Valley, 8 kilometers North of

* Brukenthal National Museum,
ana.mesaros@yahoo.com, ana.mesaros@brukenthalmuseum.ro

the valley entrance, 450 meters altitude, collected by T. Deli, É. Horváth, J. Lennert, B. Páll-Gergely and P. Subai, 03.05.2008;

- inventory number 209.716 *Alopi* *grossuana nemethi* Deli & Szekeres, collected from Vâlcan Mountains, Sohodol Valley, 8 kilometers North of the valley entrance, 450 meters altitude, collected by T. Deli, É. Horváth, J. Lennert, B. Páll-Gergely and P. Subai, 03.05.2008.

The special *Alopi* collections from the Natural History Museum in Sibiu have been of interest to European scientific researchers because of the large number of species included. The majority of specimens are endemic to Romania, but the most important parts of the collections are the type specimens and varieties of some *Alopi* species collected by Eduard Albert Bielz (1827 – 1898), Moritz von Kimakowicz (1849 – 1921) and later his son Richard von Kimakowicz (1876 – 1973).

The specimens collected by these international renowned scientists are part of the Natural History Museum from Sibiu heritage. The specimens were described in the journal of the Transylvanian Society for Natural Sciences (*Verhandlungen und Mittheilungen des siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt*), the museum's founding Society.

The collected materials included 18 new species, for that period, belonging to the *Alopi* genus. From the 18 holotypes Corocleanu (1968, 422 – 424) studied eight of these species. Also Corocleanu (1977, 213 – 223) published the 18.324 specimens belonging to the 2 genres *Herilla* (H. et A. Adams, 1855) and *Alopi* (H. et

A. Adams, 1855), in total 17 species and 43 varieties from the Bielz Collection.

The Special *Alopi* Kimakowicz Collection counts 73.321 specimens (from which 37.190 are doubles) collected and identified by Moritz and Richard von Kimakowicz. Bought Moritz von Kimakowicz and his son dedicated their studies in the field to this genus (Coan *et al.* 2011, 4). Because of the large number of specimens and the general scientific organization this collection is unique in our country. Corocleanu (1987, 275 – 295) published a list of 34.878 specimen of *Alopi* subspecies found in the Kimakowicz Collection, collected from Romania. The conclusion of the study was that there are 62 species and subspecies of the genus *Alopi* in the Kimakowicz Special *Alopi* Collection.

Even with this large number of species and subspecies, in our museum collections have not been identified until today specimens belonging to the 6 subspecies donated by Szekeres. Thru this donation the value of the Malacological Collections from the Natural History Museum in Sibiu has increased and once again it is confirmed that these collections are not only a national scientific heritage but an international one.

The Natural History Museum in Sibiu is profound honored to have received this donation from Miklos Szekeres.

The donation could not have been possible without the help of Ioan Sîrbu PhD, „Lucian Blaga” University, whom we would like to thank for his contribution.

REFERENCES

- | | |
|-------------------------|--|
| Coan <i>et al.</i> 2011 | Coan Eugene, Kabat Alan, Petit Richard, 2,400 years of malacology, 8th edition, In: <i>American Malacological Society</i> (2011) p. 4, |
| Corocleanu 1968 | Corocleanu Ileana, <i>Tipurile câtorva specii și varietăți ale genului Alopi H. et A. Adams 1855 păstrate în colecția de moluște a Muzeului Brukenthal</i> . In: <i>Revista Muzeelor</i> , 5(V), București (1968), p. 422 – 424. |
| Corocleanu 1977 | Corocleanu Ileana, <i>Alopiinele (Mollusca Gastropoda Pulmonata, Fam. Clausiliidae) în colecțiile Muzeului de Istorie Naturală din Sibiu (Partea I-a)</i> . In: <i>Muzeul Brukenthal, Studii și Comunicări, Științe Naturale</i> 21, (1977) Sibiu, p. 213 – 223. |
| Corocleanu 1987 | Corocleanu, Ileana, <i>Clausiliidae (Gen. Alopi) din Colecțiile Muzeului de Istorie Naturală din Sibiu (partea a II-a)</i> . In: <i>Complexul Muzeal Sibiu</i> 1, Sibiu (1987), p. 275 – 295. |

REVIEWERS FOR
BRUKENTHAL ACTA MUSEI VII.3

The Natural History Museum of Sibiu staff is deeply grateful to the following specialists who gave of their time to review manuscripts submitted in 2012, for publication in *Brukenthal Acta Musei VII.3*:

AGATHE Sabine, University of Salzburg, Austria
BENEDEK Klara, Sapientia Hungarian University of Transilvania, Târgu Mureş, Romania
BERGER Helmut, Consulting Engineering Office for Ecology, Salzburg, Austria
BUCŞA Corneliu, Lucian Blaga University of Sibiu, Romania
BURNAZ Silvia, Museum of Dacian and Roman Civilization, Deva, Romania
CANTOR Maria, University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, Romania
CHIMIŞLIU Cornelia, Oltenia Museum of Craiova, Romania
CHOBANOV Dragan, Bulgarian Academy of Science, Bulgaria
DINCĂ Vlad, University of Stockholm, Sweden
ESCHNER Anita, Natural History Museum Vienna, Austria
FEHÉR Zoltán, Hungarian Natural History Museum Budapest, Hungary
FRINK József Pál, Cheile Bicazului – Hăşmaş National Park, Romania
GIUŞCĂ Roxana, American Sentinel University, U.S.A
GOLBAN Dorina, Țării Crişurilor Museum, Oradea, Romania
HOLUŠA Jaroslav, Czech University of Life Sciences, Czech Republic
IORGU Ionuț Ștefan, Grigore Antipa National Museum of Natural History, Bucharest, Romania
KISHOR Kumar, Wadia Institute of Himalayan Geology, Dehradun, India
LE BOULENGÉ Éric, University of Leuven, Belgium
LUPU Gabriel, Danube Delta National Institute, Tulcea, Romania
MOGA Cosmin, Ecotransilvania Environmental and Education Association, Romania
MURARIU Dumitru, Grigore Antipa National Museum of Natural History, Bucharest, Romania
OȚEL Vasile, Danube Delta Research Institute, Romania
ÖLLERER Kinga, Institute of Biology of the Romanian Academy, Bucharest, Romania
PÁLL-GERGELY Barna, Shinshu University, Matsumoto, Japan
RÁKOSY László, Babes-Bolyai University of Cluj-Napoca, Romania
RIESE Heike, Senckenberg Museum für Naturkunde of Görlitz, Germany
STANCA-MOISE Cristina, Lucian Blaga University of Sibiu, Romania
TAMAȘ Mircea, University of Medicine and Pharmacy of Cluj-Napoca, Romania
TODERAȘ Teodor, Faculty of Geography and Tourism of Sibiu, Romania
VONICA Ghizela, Brukenthal National Museum, Romania
XU Henglong, Institute of Evolution and Marine Biodiversity, Ocean University of China, China

